



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 10:45 AM GMT

PDB ID : 3MZS
Title : Crystal Structure of Cytochrome P450 CYP11A1 in complex with 22-hydroxy-cholesterol
Authors : Stout, C.D.; Annalora, A.; Mast, N.; Pikuleva, I.
Deposited on : 2010-05-12
Resolution : 2.50 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	3553 (2.50-2.50)
Clashscore	102246	4242 (2.50-2.50)
Ramachandran outliers	100387	4156 (2.50-2.50)
Sidechain outliers	100360	4158 (2.50-2.50)
RSRZ outliers	91569	3562 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	486	<div> <div>5%</div> <div>52% 40% . . .</div> </div>
1	B	486	<div> <div>7%</div> <div>46% 43% 8% . .</div> </div>
1	C	486	<div> <div>8%</div> <div>50% 40% 6% . .</div> </div>
1	D	486	<div> <div>7%</div> <div>46% 42% 8% . .</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	IPA	A	502	-	-	-	X
4	IPA	B	502	-	-	X	X
4	IPA	C	502	-	-	X	X
4	IPA	D	502	-	-	-	X

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 16012 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cholesterol side-chain cleavage enzyme.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	470	Total	C	N	O	S	0	0	0
			3904	2541	664	683	16			
1	B	470	Total	C	N	O	S	0	0	0
			3904	2541	664	683	16			
1	C	470	Total	C	N	O	S	0	0	0
			3904	2541	664	683	16			
1	D	470	Total	C	N	O	S	0	0	0
			3904	2541	664	683	16			

There are 24 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MET	-	EXPRESSION TAG	UNP P00189
A	2	ALA	-	EXPRESSION TAG	UNP P00189
A	483	HIS	-	EXPRESSION TAG	UNP P00189
A	484	HIS	-	EXPRESSION TAG	UNP P00189
A	485	HIS	-	EXPRESSION TAG	UNP P00189
A	486	HIS	-	EXPRESSION TAG	UNP P00189
B	1	MET	-	EXPRESSION TAG	UNP P00189
B	2	ALA	-	EXPRESSION TAG	UNP P00189
B	483	HIS	-	EXPRESSION TAG	UNP P00189
B	484	HIS	-	EXPRESSION TAG	UNP P00189
B	485	HIS	-	EXPRESSION TAG	UNP P00189
B	486	HIS	-	EXPRESSION TAG	UNP P00189
C	1	MET	-	EXPRESSION TAG	UNP P00189
C	2	ALA	-	EXPRESSION TAG	UNP P00189
C	483	HIS	-	EXPRESSION TAG	UNP P00189
C	484	HIS	-	EXPRESSION TAG	UNP P00189
C	485	HIS	-	EXPRESSION TAG	UNP P00189
C	486	HIS	-	EXPRESSION TAG	UNP P00189
D	1	MET	-	EXPRESSION TAG	UNP P00189
D	2	ALA	-	EXPRESSION TAG	UNP P00189
D	483	HIS	-	EXPRESSION TAG	UNP P00189

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
D	484	HIS	-	EXPRESSION TAG	UNP P00189
D	485	HIS	-	EXPRESSION TAG	UNP P00189
D	486	HIS	-	EXPRESSION TAG	UNP P00189

- Molecule 2 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula: $C_{34}H_{32}FeN_4O_4$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total 43	C 34	Fe 1	N 4	O 4	0	0
2	B	1	Total 43	C 34	Fe 1	N 4	O 4	0	0
2	C	1	Total 43	C 34	Fe 1	N 4	O 4	0	0
2	D	1	Total 43	C 34	Fe 1	N 4	O 4	0	0

- Molecule 3 is (3ALPHA,8ALPHA,22R)-CHOLEST-5-ENE-3,22-DIOL (three-letter code: HC9) (formula: $C_{27}H_{46}O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			29	27	2		
3	B	1	Total	C	O	0	0
			29	27	2		
3	C	1	Total	C	O	0	0
			29	27	2		
3	D	1	Total	C	O	0	0
			29	27	2		

- Molecule 4 is ISOPROPYL ALCOHOL (three-letter code: IPA) (formula: C_3H_8O).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 4 3 1	0	0
4	B	1	Total C O 4 3 1	0	0
4	C	1	Total C O 4 3 1	0	0
4	D	1	Total C O 4 3 1	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	25	Total O 25 25	0	0
5	B	15	Total O 15 15	0	0
5	C	27	Total O 27 27	0	0
5	D	25	Total O 25 25	0	0


3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Cholesterol side-chain cleavage enzyme




• Molecule 1: Cholesterol side-chain cleavage enzyme







• Molecule 1: Cholesterol side-chain cleavage enzyme



• Molecule 1: Cholesterol side-chain cleavage enzyme





4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	109.45Å 94.63Å 113.50Å 90.00° 89.96° 90.00°	Depositor
Resolution (Å)	60.01 – 2.50 60.56 – 2.50	Depositor EDS
% Data completeness (in resolution range)	94.9 (60.01-2.50) 99.3 (60.56-2.50)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	0.07	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.29 (at 2.51Å)	Xtriage
Refinement program	REFMAC 5.5.0072	Depositor
R, R_{free}	0.266 , 0.281 0.267 , 0.279	Depositor DCC
R_{free} test set	3990 reflections (5.25%)	DCC
Wilson B-factor (Å ²)	84.1	Xtriage
Anisotropy	0.692	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 39.2	EDS
Estimated twinning fraction	0.546 for H, K, L 0.454 for -H, -K, L 0.000 for -l,k,h 0.276 for h,-k,-l 0.000 for -l,-k,-h	Xtriage
Reported twinning fraction	0.546 for H, K, L 0.454 for -H, -K, L	Depositor
L-test for twinning ²	$\langle L \rangle = 0.52$, $\langle L^2 \rangle = 0.35$	Xtriage
Outliers	1 of 80275 reflections (0.001%)	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	16012	wwPDB-VP
Average B, all atoms (Å ²)	65.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.36% of the height of the origin peak. No significant pseudotranslation is detected.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: HEM, HC9, IPA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.99	5/4014 (0.1%)	0.89	4/5435 (0.1%)
1	B	0.89	1/4014 (0.0%)	0.84	7/5435 (0.1%)
1	C	0.98	2/4014 (0.0%)	0.89	4/5435 (0.1%)
1	D	0.93	2/4014 (0.0%)	0.89	7/5435 (0.1%)
All	All	0.95	10/16056 (0.1%)	0.88	22/21740 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	0	1
1	C	0	1
1	D	0	1
All	All	0	4

The worst 5 of 10 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	289	VAL	CA-CB	-9.40	1.35	1.54
1	C	260	PRO	N-CA	7.47	1.59	1.47
1	A	293	SER	CB-OG	7.21	1.51	1.42
1	D	156	PHE	C-N	6.92	1.50	1.34
1	A	458	PHE	CE2-CZ	6.07	1.48	1.37

The worst 5 of 22 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	156	PHE	CA-C-O	10.56	142.28	120.10
1	D	156	PHE	O-C-N	-9.99	106.71	122.70
1	B	347	ARG	NE-CZ-NH2	8.85	124.73	120.30
1	A	385	ARG	NE-CZ-NH2	-7.67	116.47	120.30
1	D	261	GLY	N-CA-C	-7.15	95.23	113.10

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	259	TYR	Peptide
1	B	259	TYR	Peptide
1	C	259	TYR	Peptide
1	D	259	TYR	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3904	0	3910	185	0
1	B	3904	0	3910	273	0
1	C	3904	0	3910	207	0
1	D	3904	0	3910	225	0
2	A	43	0	30	4	0
2	B	43	0	30	4	0
2	C	43	0	30	5	0
2	D	43	0	30	5	0
3	A	29	0	46	1	0
3	B	29	0	46	2	0
3	C	29	0	45	5	0
3	D	29	0	45	1	0
4	A	4	0	8	1	0
4	B	4	0	8	8	0
4	C	4	0	8	4	0
4	D	4	0	8	2	0
5	A	25	0	0	1	0
5	B	15	0	0	1	0
5	C	27	0	0	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	D	25	0	0	4	0
All	All	16012	0	15974	875	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 27.

The worst 5 of 875 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:128:ASN:HB3	1:B:260:PRO:CG	1.22	1.61
1:B:128:ASN:HB3	1:B:260:PRO:CD	1.31	1.60
1:B:128:ASN:CB	1:B:260:PRO:CG	1.83	1.55
1:B:128:ASN:CB	1:B:260:PRO:HG2	1.42	1.45
1:D:156:PHE:O	1:D:157:VAL:HG23	1.47	1.12

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	468/486 (96%)	390 (83%)	66 (14%)	12 (3%)	7	10
1	B	468/486 (96%)	391 (84%)	59 (13%)	18 (4%)	4	5
1	C	468/486 (96%)	391 (84%)	59 (13%)	18 (4%)	4	5
1	D	468/486 (96%)	376 (80%)	77 (16%)	15 (3%)	5	6
All	All	1872/1944 (96%)	1548 (83%)	261 (14%)	63 (3%)	5	6

5 of 63 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	258	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	404	LYS
1	B	33	SER
1	B	255	GLU
1	B	327	ASP

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	426/440 (97%)	398 (93%)	28 (7%)	21	38
1	B	426/440 (97%)	383 (90%)	43 (10%)	9	17
1	C	426/440 (97%)	389 (91%)	37 (9%)	13	24
1	D	426/440 (97%)	378 (89%)	48 (11%)	7	13
All	All	1704/1760 (97%)	1548 (91%)	156 (9%)	11	21

5 of 156 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	475	PHE
1	C	256	PHE
1	D	400	ARG
1	C	19	ASN
1	C	120	VAL

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 49 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	319	ASN
1	C	45	GLN
1	D	237	ASN
1	B	366	GLN
1	C	138	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

12 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	HEM	A	500	1,3	30,50,50	3.80	14 (46%)	24,82,82	2.93	11 (45%)
3	HC9	A	501	2	32,32,32	1.02	0	49,50,50	1.69	10 (20%)
4	IPA	A	502	-	3,3,3	0.86	0	3,3,3	0.49	0
2	HEM	B	500	1,3	30,50,50	3.53	13 (43%)	24,82,82	2.71	13 (54%)
3	HC9	B	501	2	32,32,32	0.82	0	49,50,50	1.57	7 (14%)
4	IPA	B	502	-	3,3,3	0.71	0	3,3,3	0.83	0
2	HEM	C	500	1,3	30,50,50	3.88	14 (46%)	24,82,82	2.80	10 (41%)
3	HC9	C	501	2	32,32,32	0.82	0	49,50,50	1.65	13 (26%)
4	IPA	C	502	-	3,3,3	0.67	0	3,3,3	0.50	0
2	HEM	D	500	1,3	30,50,50	3.67	15 (50%)	24,82,82	2.83	10 (41%)
3	HC9	D	501	2	32,32,32	1.15	3 (9%)	49,50,50	2.23	15 (30%)
4	IPA	D	502	-	3,3,3	0.73	0	3,3,3	0.58	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means

no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HEM	A	500	1,3	-	0/10/54/54	0/0/8/8
3	HC9	A	501	2	-	0/13/71/71	0/4/4/4
4	IPA	A	502	-	-	0/0/0/0	0/0/0/0
2	HEM	B	500	1,3	-	0/10/54/54	0/0/8/8
3	HC9	B	501	2	-	0/13/71/71	0/4/4/4
4	IPA	B	502	-	-	0/0/0/0	0/0/0/0
2	HEM	C	500	1,3	-	0/10/54/54	0/0/8/8
3	HC9	C	501	2	-	0/13/71/71	0/4/4/4
4	IPA	C	502	-	-	0/0/0/0	0/0/0/0
2	HEM	D	500	1,3	-	0/10/54/54	0/0/8/8
3	HC9	D	501	2	-	0/13/71/71	0/4/4/4
4	IPA	D	502	-	-	0/0/0/0	0/0/0/0

The worst 5 of 59 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	500	HEM	C2C-C1C	-11.53	1.30	1.52
2	C	500	HEM	C2C-C1C	-11.12	1.31	1.52
2	A	500	HEM	C2C-C1C	-10.78	1.32	1.52
2	B	500	HEM	C2C-C1C	-9.92	1.33	1.52
2	A	500	HEM	C3B-CAB	-9.90	1.32	1.51

The worst 5 of 89 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	500	HEM	C3B-C4B-CHC	-8.68	110.94	123.16
2	A	500	HEM	C3B-C4B-CHC	-8.62	111.02	123.16
2	C	500	HEM	C3B-C4B-CHC	-7.53	112.57	123.16
2	B	500	HEM	C3B-C4B-CHC	-6.60	113.87	123.16
3	D	501	HC9	C12-C13-C14	-5.46	98.42	107.31

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

12 monomers are involved in 42 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	500	HEM	4	0
3	A	501	HC9	1	0
4	A	502	IPA	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	500	HEM	4	0
3	B	501	HC9	2	0
4	B	502	IPA	8	0
2	C	500	HEM	5	0
3	C	501	HC9	5	0
4	C	502	IPA	4	0
2	D	500	HEM	5	0
3	D	501	HC9	1	0
4	D	502	IPA	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	470/486 (96%)	0.50	25 (5%) 30 34	34, 65, 84, 96	0
1	B	470/486 (96%)	0.59	34 (7%) 18 20	39, 67, 87, 97	0
1	C	470/486 (96%)	0.53	37 (7%) 15 17	35, 65, 84, 93	0
1	D	470/486 (96%)	0.54	33 (7%) 19 22	32, 67, 86, 94	0
All	All	1880/1944 (96%)	0.54	129 (6%) 20 22	32, 66, 85, 97	0

The worst 5 of 129 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	266	LEU	6.1
1	C	263	LEU	5.5
1	B	154	GLY	4.4
1	B	184	LEU	4.3
1	A	446	VAL	3.8

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
4	IPA	B	502	4/4	0.94	0.52	6.59	60,61,62,63	0
4	IPA	C	502	4/4	0.96	0.51	5.29	60,61,61,63	0
4	IPA	A	502	4/4	0.94	0.52	5.01	49,49,50,51	0
4	IPA	D	502	4/4	0.89	0.40	4.42	59,59,61,65	0
3	HC9	A	501	29/29	0.95	0.27	1.48	41,49,58,60	0
3	HC9	B	501	29/29	0.94	0.27	1.34	44,53,64,68	0
3	HC9	D	501	29/29	0.96	0.23	0.87	39,48,56,61	0
3	HC9	C	501	29/29	0.97	0.23	0.66	44,54,61,66	0
2	HEM	C	500	43/43	0.97	0.18	-0.34	37,50,56,59	0
2	HEM	D	500	43/43	0.97	0.18	-0.43	18,41,56,62	0
2	HEM	A	500	43/43	0.98	0.16	-0.79	25,43,54,57	0
2	HEM	B	500	43/43	0.96	0.17	-0.81	34,47,55,60	0

6.5 Other polymers [i](#)

There are no such residues in this entry.