



# Full wwPDB NMR Structure Validation Report ⓘ

Sep 27, 2016 – 08:30 PM EDT

PDB ID : 2N4Q  
Title : Solution NMR structure of CBX8 in complex with AF9 (CBX8-AF9)  
Authors : Kuntimaddi, A.; Leach, B.I.; Bushweller, J.H.  
Deposited on : 2015-06-30

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

Cyrange	:	Kirchner and Güntert (2011)
NmrClust	:	Kelley et al. (1996)
MolProbity	:	4.02b-467
Mogul	:	unknown
Percentile statistics	:	20151230.v01 (using entries in the PDB archive December 30th 2015)
RCI	:	v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV	:	Wang et al. (2010)
ShiftChecker	:	rb-20027939
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	rb-20027939


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*SOLUTION NMR*

The overall completeness of chemical shifts assignment is 80%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	114402	11133
Ramachandran outliers	111179	9975
Sidechain outliers	111093	9958

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	24	
2	B	70	

## 2 Ensemble composition and analysis

This entry contains 10 models. Model 2 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *lowest energy*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:332-A:343, B:503-B:562 (72)	0.24	2

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 1 clusters and 4 single-model clusters were found.

Cluster number	Models
1	1, 2, 3, 5, 6, 9
Single-model clusters	4; 7; 8; 10

### 3 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 1532 atoms, of which 771 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called Chromobox protein homolog 8.

Mol	Chain	Residues	Atoms					Trace
1	A	24	Total	C	H	N	O	0
			368	111	187	34	36	

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	326	THR	-	EXPRESSION TAG	UNP Q9HC52

- Molecule 2 is a protein called Protein AF-9.

Mol	Chain	Residues	Atoms					Trace
2	B	70	Total	C	H	N	O	S
			1164	363	584	101	113	3
								0

There is a discrepancy between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
B	499	MET	-	EXPRESSION TAG	UNP P42568

## 4 Residue-property plots


### 4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA and DNA chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: Chromobox protein homolog 8



- Molecule 2: Protein AF-9




### 4.2 Scores per residue for each member of the ensemble


Colouring as in section 4.1 above.

#### 4.2.1 Score per residue for model 1

- Molecule 1: Chromobox protein homolog 8




- Molecule 2: Protein AF-9




### 4.2.2 Score per residue for model 2 (medoid)

- Molecule 1: Chromobox protein homolog 8




- Molecule 2: Protein AF-9




### 4.2.3 Score per residue for model 3

- Molecule 1: Chromobox protein homolog 8




- Molecule 2: Protein AF-9




### 4.2.4 Score per residue for model 4

- Molecule 1: Chromobox protein homolog 8




- Molecule 2: Protein AF-9




### 4.2.5 Score per residue for model 5

- Molecule 1: Chromobox protein homolog 8




- Molecule 2: Protein AF-9




### 4.2.6 Score per residue for model 6

- Molecule 1: Chromobox protein homolog 8




- Molecule 2: Protein AF-9




### 4.2.7 Score per residue for model 7

- Molecule 1: Chromobox protein homolog 8




- Molecule 2: Protein AF-9




### 4.2.8 Score per residue for model 8

- Molecule 1: Chromobox protein homolog 8




- Molecule 2: Protein AF-9




### 4.2.9 Score per residue for model 9

- Molecule 1: Chromobox protein homolog 8




- Molecule 2: Protein AF-9




### 4.2.10 Score per residue for model 10

- Molecule 1: Chromobox protein homolog 8



- Molecule 2: Protein AF-9





## 5 Refinement protocol and experimental data overview

The models were refined using the following method: *simulated annealing*.

Of the 100 calculated structures, 10 were deposited, based on the following criterion: *structures with the lowest energy*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
X-PLOR NIH	structure solution	
X-PLOR NIH	refinement	

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	2n4q_cs.cif
Number of chemical shift lists	1
Total number of shifts	1036
Number of shifts mapped to atoms	1036
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	80%

No validations of the models with respect to experimental NMR restraints is performed at this time.

## 6 Model quality [i](#)

### 6.1 Standard geometry [i](#)

There are no covalent bond-length or bond-angle outliers.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 6.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	92	112	112	6±2
2	B	510	518	514	13±3
All	All	6020	6300	6260	170

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 14.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:335:ALA:HB2	2:B:511:HIS:NE2	0.75	1.96	2	7
1:A:335:ALA:HB2	2:B:511:HIS:CE1	0.75	2.16	8	3
2:B:523:LEU:O	2:B:527:VAL:HG23	0.66	1.89	2	6
1:A:337:ILE:HD11	2:B:545:PHE:CE1	0.66	2.26	7	4
1:A:341:ARG:NH2	2:B:523:LEU:HD23	0.64	2.08	9	1
2:B:553:THR:HG23	2:B:554:THR:N	0.62	2.08	8	1
1:A:333:LEU:HD23	2:B:547:LEU:CB	0.61	2.25	8	3
2:B:504:LEU:O	2:B:508:VAL:HG23	0.59	1.97	6	8
2:B:548:CYS:SG	2:B:549:SER:N	0.59	2.76	8	1
2:B:511:HIS:CE1	2:B:547:LEU:HD11	0.55	2.37	6	1
2:B:528:ASN:N	2:B:528:ASN:ND2	0.55	2.53	6	1
2:B:553:THR:HG23	2:B:554:THR:H	0.55	1.61	8	1
2:B:551:ASP:O	2:B:553:THR:N	0.54	2.41	7	2

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
2:B:551:ASP:O	2:B:555:VAL:HG23	0.53	2.03	8	2
1:A:343:LEU:N	1:A:343:LEU:HD22	0.53	2.18	2	5
2:B:514:LEU:O	2:B:517:LEU:N	0.52	2.43	4	1
1:A:341:ARG:HH22	2:B:523:LEU:HD23	0.52	1.62	9	1
2:B:546:ASP:OD1	2:B:547:LEU:N	0.52	2.42	5	2
1:A:343:LEU:N	1:A:343:LEU:CD2	0.51	2.73	2	4
2:B:547:LEU:CD2	2:B:547:LEU:N	0.51	2.74	4	2
2:B:518:ARG:O	2:B:520:ARG:N	0.50	2.45	10	1
1:A:343:LEU:HD22	1:A:343:LEU:N	0.50	2.20	4	5
2:B:539:THR:N	2:B:542:THR:O	0.50	2.44	1	10
2:B:531:GLU:OE1	2:B:536:PHE:CZ	0.50	2.65	3	1
2:B:546:ASP:OD1	2:B:546:ASP:N	0.50	2.41	9	1
1:A:343:LEU:CD2	1:A:343:LEU:N	0.50	2.75	4	6
2:B:538:ILE:O	2:B:538:ILE:HG23	0.50	2.07	3	5
2:B:545:PHE:CD1	2:B:545:PHE:N	0.49	2.80	8	2
2:B:547:LEU:HD22	2:B:547:LEU:N	0.49	2.22	4	2
2:B:553:THR:OG1	2:B:554:THR:N	0.49	2.46	10	2
2:B:548:CYS:SG	2:B:548:CYS:O	0.49	2.71	9	3
2:B:550:LEU:HD12	2:B:554:THR:CB	0.49	2.37	3	1
2:B:550:LEU:HD12	2:B:554:THR:HB	0.49	1.85	3	1
2:B:553:THR:CG2	2:B:554:THR:N	0.49	2.76	8	1
2:B:551:ASP:C	2:B:553:THR:N	0.49	2.66	7	2
1:A:339:VAL:O	1:A:341:ARG:N	0.48	2.46	8	6
2:B:545:PHE:N	2:B:545:PHE:CD1	0.48	2.80	3	1
2:B:556:ARG:NH1	2:B:559:GLN:NE2	0.48	2.61	9	1
2:B:544:ASP:OD1	2:B:545:PHE:N	0.48	2.46	5	1
2:B:556:ARG:NH1	2:B:559:GLN:HE22	0.48	2.05	9	1
2:B:527:VAL:HG13	2:B:543:PHE:CE2	0.48	2.43	8	1
1:A:333:LEU:HD23	2:B:547:LEU:HB2	0.48	1.86	8	1
1:A:333:LEU:HD13	2:B:508:VAL:HG22	0.48	1.84	9	2
2:B:519:GLU:N	2:B:519:GLU:OE1	0.48	2.47	1	1
2:B:551:ASP:N	2:B:551:ASP:OD1	0.47	2.46	1	3
2:B:551:ASP:OD1	2:B:551:ASP:N	0.47	2.48	7	2
1:A:339:VAL:C	1:A:341:ARG:N	0.47	2.68	3	7
2:B:546:ASP:O	2:B:549:SER:OG	0.45	2.34	6	3
2:B:529:LEU:O	2:B:532:GLU:N	0.45	2.50	3	1
2:B:538:ILE:HG23	2:B:538:ILE:O	0.45	2.11	8	2
2:B:506:GLU:OE1	2:B:559:GLN:NE2	0.45	2.49	2	1
2:B:504:LEU:HD11	2:B:548:CYS:SG	0.45	2.51	4	1
2:B:552:LYS:O	2:B:556:ARG:CG	0.44	2.66	2	1
2:B:552:LYS:H	2:B:552:LYS:CD	0.44	2.23	4	1

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
2:B:552:LYS:CD	2:B:552:LYS:H	0.44	2.25	3	2
2:B:507:LEU:HD22	2:B:547:LEU:HB3	0.44	1.89	4	1
2:B:546:ASP:OD1	2:B:548:CYS:N	0.44	2.48	10	1
2:B:528:ASN:HD22	2:B:528:ASN:N	0.43	2.10	6	1
2:B:527:VAL:O	2:B:531:GLU:OE2	0.43	2.36	7	1
2:B:531:GLU:OE1	2:B:536:PHE:CE2	0.43	2.71	3	1
2:B:551:ASP:O	2:B:554:THR:OG1	0.43	2.36	3	1
2:B:550:LEU:O	2:B:551:ASP:O	0.43	2.37	6	1
1:A:336:ARG:CG	1:A:336:ARG:O	0.43	2.67	8	1
1:A:335:ALA:O	2:B:544:ASP:OD1	0.43	2.37	5	1
2:B:537:HIS:O	2:B:544:ASP:O	0.43	2.37	5	5
2:B:556:ARG:CD	2:B:556:ARG:C	0.43	2.87	10	1
2:B:514:LEU:O	2:B:516:THR:N	0.42	2.52	4	1
2:B:514:LEU:C	2:B:516:THR:N	0.42	2.72	7	2
1:A:333:LEU:CD1	2:B:508:VAL:HG13	0.42	2.45	4	1
1:A:339:VAL:C	1:A:341:ARG:H	0.42	2.17	3	3
1:A:339:VAL:HG23	1:A:340:ALA:N	0.41	2.31	3	1
2:B:504:LEU:HD12	2:B:504:LEU:N	0.41	2.30	10	1
2:B:546:ASP:OD2	2:B:549:SER:CB	0.41	2.69	2	1
2:B:547:LEU:C	2:B:549:SER:H	0.41	2.19	4	1
2:B:553:THR:CG2	2:B:554:THR:H	0.41	2.27	8	1
2:B:537:HIS:O	2:B:544:ASP:OD1	0.41	2.39	10	1
2:B:525:GLN:OE1	2:B:525:GLN:N	0.40	2.54	1	1
2:B:535:HIS:ND1	2:B:549:SER:OG	0.40	2.51	3	1
2:B:551:ASP:C	2:B:553:THR:H	0.40	2.20	7	1
1:A:333:LEU:HD23	2:B:547:LEU:HB3	0.40	1.93	8	1

## 6.3 Torsion angles ⓘ

### 6.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	12/24 (50%)	11±0 (88±4%)	1±0 (11±4%)	0±0 (1±3%)	29	74
2	B	60/70 (86%)	55±1 (92±2%)	4±1 (7±1%)	1±0 (1±1%)	24	71
All	All	720/940 (77%)	660 (92%)	53 (7%)	7 (1%)	24	71

All 4 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
2	B	519	GLU	3
2	B	552	LYS	2
1	A	340	ALA	1
2	B	551	ASP	1

### 6.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	10/19 (53%)	9±1 (89±9%)	1±1 (11±9%)	12	56
2	B	59/67 (88%)	43±2 (72±4%)	16±2 (28±4%)	2	21
All	All	690/860 (80%)	516 (75%)	174 (25%)	3	25

All 44 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
2	B	542	THR	10
2	B	552	LYS	10
2	B	511	HIS	9
2	B	539	THR	8
2	B	529	LEU	8
2	B	546	ASP	8
2	B	535	HIS	7
2	B	537	HIS	6
2	B	548	CYS	6
2	B	554	THR	6
2	B	538	ILE	6
2	B	514	LEU	5
2	B	551	ASP	5
2	B	505	ASP	5
2	B	560	SER	5
2	B	540	ASN	4
2	B	556	ARG	4
2	B	544	ASP	4

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Models (Total)
2	B	528	ASN	4
2	B	521	HIS	4
2	B	512	ARG	4
1	A	337	ILE	4
2	B	518	ARG	3
1	A	333	LEU	3
2	B	557	LYS	3
2	B	541	THR	3
2	B	520	ARG	3
2	B	517	LEU	3
1	A	332	SER	3
2	B	525	GLN	2
2	B	559	GLN	2
2	B	553	THR	2
2	B	519	GLU	2
2	B	503	TYR	2
2	B	523	LEU	2
2	B	510	LEU	1
2	B	550	LEU	1
1	A	339	VAL	1
2	B	516	THR	1
2	B	532	GLU	1
2	B	531	GLU	1
2	B	509	GLU	1
2	B	515	MET	1
2	B	561	TYR	1

### 6.3.3 RNA ⓘ

There are no RNA molecules in this entry.

### 6.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 6.6 Ligand geometry

There are no ligands in this entry.

## 6.7 Other polymers

There are no such molecules in this entry.

## 6.8 Polymer linkage issues

There are no chain breaks in this entry.

## 7 Chemical shift validation

The completeness of assignment taking into account all chemical shift lists is 80% for the well-defined parts and 77% for the entire structure.

### 7.1 Chemical shift list 1

File name: 2n4q\_cs.cif

Chemical shift list name: *assigned\_chem\_shift\_list\_1*

#### 7.1.1 Bookkeeping

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	1036
Number of shifts mapped to atoms	1036
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	0

#### 7.1.2 Chemical shift referencing

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction $\pm$ precision, ppm	Suggested action
$^{13}\text{C}_\alpha$	93	$-0.15 \pm 0.11$	None needed ( $< 0.5$ ppm)
$^{13}\text{C}_\beta$	88	$0.28 \pm 0.07$	None needed ( $< 0.5$ ppm)
$^{13}\text{C}'$	85	$-0.24 \pm 0.16$	None needed ( $< 0.5$ ppm)
$^{15}\text{N}$	88	$-0.02 \pm 0.59$	None needed ( $< 0.5$ ppm)

#### 7.1.3 Completeness of resonance assignments

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 80%, i.e. 773 atoms were assigned a chemical shift out of a possible 972. 9 out of 17 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	$^1\text{H}$	$^{13}\text{C}$	$^{15}\text{N}$
Backbone	355/358 (99%)	143/143 (100%)	141/144 (98%)	71/71 (100%)
Sidechain	360/539 (67%)	202/312 (65%)	158/199 (79%)	0/28 (0%)

*Continued on next page...*



Continued from previous page...

	Total	<sup>1</sup> H	<sup>13</sup> C	<sup>15</sup> N
Aromatic	58/75 (77%)	30/39 (77%)	28/28 (100%)	0/8 (0%)
Overall	773/972 (80%)	375/494 (76%)	327/371 (88%)	71/107 (66%)

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 77%, i.e. 924 atoms were assigned a chemical shift out of a possible 1201. 9 out of 17 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	<sup>1</sup> H	<sup>13</sup> C	<sup>15</sup> N
Backbone	440/464 (95%)	174/185 (94%)	178/188 (95%)	88/91 (97%)
Sidechain	426/662 (64%)	237/385 (62%)	189/244 (77%)	0/33 (0%)
Aromatic	58/75 (77%)	30/39 (77%)	28/28 (100%)	0/8 (0%)
Overall	924/1201 (77%)	441/609 (72%)	395/460 (86%)	88/132 (67%)


#### 7.1.4 Statistically unusual chemical shifts [i](#)

There are no statistically unusual chemical shifts.

#### 7.1.5 Random Coil Index (RCI) plots [i](#)

The images below report *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition.

Random coil index (RCI) for chain A:



Random coil index (RCI) for chain B:

