



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 10:53 AM GMT

PDB ID : 3N87
Title : Crystal structure of 3-dehydroquinate dehydratase from Mycobacterium tuberculosis in complex with inhibitor 3
Authors : Dias, M.V.B.; Snee, W.C.; Bromfield, K.M.; Payne, R.; Palaninathan, S.K.; Ciulli, A.; Howard, N.I.; Abell, C.; Sacchettini, J.C.; Blundell, T.L.
Deposited on : 2010-05-27
Resolution : 2.40 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	2919 (2.40-2.40)
Clashscore	102246	3407 (2.40-2.40)
Ramachandran outliers	100387	3351 (2.40-2.40)
Sidechain outliers	100360	3352 (2.40-2.40)
RSRZ outliers	91569	2928 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	147	<div> <div>2%</div> <div> <div></div> <div>73%</div> <div>20%</div> <div>• •</div> </div> </div>
1	B	147	<div> <div></div> <div> <div>78%</div> <div>16%</div> <div>• •</div> </div> </div>
1	C	147	<div> <div>4%</div> <div> <div></div> <div>72%</div> <div>20%</div> <div>• • •</div> </div> </div>
1	D	147	<div> <div>6%</div> <div> <div></div> <div>71%</div> <div>19%</div> <div>5%</div> <div>• • •</div> </div> </div>
1	E	147	<div> <div>4%</div> <div> <div></div> <div>69%</div> <div>22%</div> <div>5%</div> <div>•</div> </div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	147	
1	G	147	
1	H	147	
1	I	147	
1	J	147	
1	K	147	
1	L	147	
1	M	147	
1	N	147	
1	O	147	
1	P	147	
1	Q	147	
1	R	147	
1	S	147	
1	T	147	
1	U	147	
1	V	147	
1	W	147	
1	X	147	

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 27701 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 3-dehydroquinate dehydratase.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	141	Total	C	N	O	S	0	0	0
			1065	671	193	200	1			
1	B	141	Total	C	N	O	S	0	0	0
			1071	674	196	200	1			
1	C	141	Total	C	N	O	S	0	0	0
			1065	670	196	198	1			
1	D	141	Total	C	N	O	S	0	0	0
			1069	672	196	200	1			
1	E	141	Total	C	N	O	S	0	0	0
			1071	674	196	200	1			
1	F	141	Total	C	N	O	S	0	1	0
			1071	675	195	200	1			
1	G	141	Total	C	N	O	S	0	0	0
			1071	674	196	200	1			
1	H	141	Total	C	N	O	S	0	0	0
			1071	674	196	200	1			
1	I	141	Total	C	N	O	S	0	0	0
			1071	674	196	200	1			
1	J	141	Total	C	N	O	S	0	0	0
			1065	670	196	198	1			
1	K	141	Total	C	N	O	S	0	0	0
			1061	669	193	198	1			
1	L	141	Total	C	N	O	S	0	0	0
			1071	674	196	200	1			
1	M	141	Total	C	N	O	S	0	0	0
			1071	674	196	200	1			
1	N	141	Total	C	N	O	S	0	1	0
			1077	678	198	200	1			
1	O	141	Total	C	N	O	S	0	0	0
			1071	674	196	200	1			
1	P	141	Total	C	N	O	S	0	0	0
			1071	674	196	200	1			

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	Q	141	Total	C	N	O	S	0	0	0
			1065	671	193	200	1			
1	R	141	Total	C	N	O	S	0	0	0
			1059	667	193	198	1			
1	S	141	Total	C	N	O	S	0	0	0
			1071	674	196	200	1			
1	T	141	Total	C	N	O	S	0	0	0
			1071	674	196	200	1			
1	U	141	Total	C	N	O	S	0	0	0
			1071	674	196	200	1			
1	V	141	Total	C	N	O	S	0	0	0
			1067	672	196	198	1			
1	W	141	Total	C	N	O	S	0	0	0
			1065	671	193	200	1			
1	X	141	Total	C	N	O	S	0	0	0
			1061	669	193	198	1			

- Molecule 2 is (1R,4R,5R)-1,4,5-TRIHIDROXY-3-[3-(PHENYLCARBONYL)PHENYL]CYCLOHEX-2-ENE-1-CARBOXYLIC ACID (three-letter code: N87) (formula: C₂₀H₁₈O₆).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			26	20	6		
2	B	1	Total	C	O	0	0
			26	20	6		
2	C	1	Total	C	O	0	0
			26	20	6		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	D	1	Total	C	O	0	0
			26	20	6		
2	E	1	Total	C	O	0	0
			26	20	6		
2	F	1	Total	C	O	0	0
			26	20	6		
2	G	1	Total	C	O	0	0
			26	20	6		
2	H	1	Total	C	O	0	0
			26	20	6		
2	I	1	Total	C	O	0	0
			26	20	6		
2	J	1	Total	C	O	0	0
			26	20	6		
2	K	1	Total	C	O	0	0
			26	20	6		
2	L	1	Total	C	O	0	0
			26	20	6		
2	M	1	Total	C	O	0	0
			26	20	6		
2	N	1	Total	C	O	0	0
			26	20	6		
2	O	1	Total	C	O	0	0
			26	20	6		
2	P	1	Total	C	O	0	0
			26	20	6		
2	Q	1	Total	C	O	0	0
			26	20	6		
2	R	1	Total	C	O	0	0
			26	20	6		
2	S	1	Total	C	O	0	0
			26	20	6		
2	T	1	Total	C	O	0	0
			26	20	6		
2	U	1	Total	C	O	0	0
			26	20	6		
2	V	1	Total	C	O	0	0
			26	20	6		
2	W	1	Total	C	O	0	0
			26	20	6		
2	X	1	Total	C	O	0	0
			26	20	6		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	53	Total O 53 53	0	0
3	B	90	Total O 90 90	0	0
3	C	59	Total O 59 59	0	0
3	D	54	Total O 54 54	0	0
3	E	78	Total O 78 78	0	0
3	F	60	Total O 60 60	0	0
3	G	66	Total O 66 66	0	0
3	H	59	Total O 59 59	0	0
3	I	48	Total O 48 48	0	0
3	J	70	Total O 70 70	0	0
3	K	55	Total O 55 55	0	0
3	L	49	Total O 49 49	0	0
3	M	57	Total O 57 57	0	0
3	N	54	Total O 54 54	0	0
3	O	53	Total O 53 53	0	0
3	P	67	Total O 67 67	0	0
3	Q	50	Total O 50 50	0	0
3	R	63	Total O 63 63	0	0
3	S	66	Total O 66 66	0	0
3	T	73	Total O 73 73	0	0
3	U	59	Total O 59 59	0	0

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	V	61	Total 61	O 61	0	0
3	W	43	Total 43	O 43	0	0
3	X	48	Total 48	O 48	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: 3-dehydroquinase dehydratase




- Molecule 1: 3-dehydroquinase dehydratase




- Molecule 1: 3-dehydroquinase dehydratase




- Molecule 1: 3-dehydroquinase dehydratase




- Molecule 1: 3-dehydroquinase dehydratase




- Molecule 1: 3-dehydroquinase dehydratase




- Molecule 1: 3-dehydroquinase dehydratase




- Molecule 1: 3-dehydroquinase dehydratase



- Molecule 1: 3-dehydroquinase dehydratase




- Molecule 1: 3-dehydroquinase dehydratase






• Molecule 1: 3-dehydroquinase dehydratase




• Molecule 1: 3-dehydroquinase dehydratase




• Molecule 1: 3-dehydroquinase dehydratase



• Molecule 1: 3-dehydroquinase dehydratase




• Molecule 1: 3-dehydroquinase dehydratase




THR


- Molecule 1: 3-dehydroquinate dehydratase




- Molecule 1: 3-dehydroquinate dehydratase




- Molecule 1: 3-dehydroquinate dehydratase




- Molecule 1: 3-dehydroquinate dehydratase



- Molecule 1: 3-dehydroquinate dehydratase




- Molecule 1: 3-dehydroquinate dehydratase




THR


- Molecule 1: 3-dehydroquinase dehydratase



- Molecule 1: 3-dehydroquinase dehydratase



- Molecule 1: 3-dehydroquinase dehydratase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	91.02Å 96.78Å 128.11Å 78.36° 81.24° 78.58°	Depositor
Resolution (Å)	36.22 – 2.40 36.22 – 2.40	Depositor EDS
% Data completeness (in resolution range)	95.4 (36.22-2.40) 89.9 (36.22-2.40)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	0.07	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.23 (at 2.39Å)	Xtriage
Refinement program	REFMAC 5.5.0102	Depositor
R, R_{free}	0.191 , 0.274 0.194 , 0.274	Depositor DCC
R_{free} test set	7761 reflections (5.25%)	DCC
Wilson B-factor (Å ²)	31.9	Xtriage
Anisotropy	0.154	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 63.3	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 155522 reflections	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	27701	wwPDB-VP
Average B, all atoms (Å ²)	33.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.46% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: N87

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.68	0/1083	0.81	1/1476 (0.1%)
1	B	0.84	1/1089 (0.1%)	0.90	1/1483 (0.1%)
1	C	0.77	0/1082	0.89	3/1473 (0.2%)
1	D	0.76	0/1087	0.83	0/1480
1	E	0.75	0/1089	0.87	0/1483
1	F	0.79	0/1094	0.87	1/1491 (0.1%)
1	G	0.82	0/1089	0.90	0/1483
1	H	0.79	0/1089	0.89	1/1483 (0.1%)
1	I	0.72	0/1089	0.81	0/1483
1	J	0.84	0/1082	0.92	1/1473 (0.1%)
1	K	0.73	0/1079	0.84	1/1471 (0.1%)
1	L	0.74	0/1089	0.83	0/1483
1	M	0.76	0/1089	0.88	2/1483 (0.1%)
1	N	0.77	0/1100	0.90	1/1498 (0.1%)
1	O	0.75	0/1089	0.83	0/1483
1	P	0.84	0/1089	0.94	3/1483 (0.2%)
1	Q	0.79	0/1083	0.86	0/1476
1	R	0.79	0/1076	0.92	2/1466 (0.1%)
1	S	0.84	0/1089	0.92	1/1483 (0.1%)
1	T	0.83	0/1089	0.94	2/1483 (0.1%)
1	U	0.83	0/1089	0.83	0/1483
1	V	0.86	1/1085 (0.1%)	0.97	0/1478
1	W	0.74	0/1083	0.80	0/1476
1	X	0.71	0/1079	0.83	0/1471
All	All	0.78	2/26081 (0.0%)	0.88	20/35525 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	E	0	2
1	H	0	1
1	I	0	2
1	Q	0	2
1	U	0	1
1	V	0	1
1	X	0	1
All	All	0	11

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	V	113	ARG	CG-CD	5.75	1.66	1.51
1	B	133	TYR	CE2-CZ	5.13	1.45	1.38

The worst 5 of 20 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	3	LEU	CA-CB-CG	7.50	132.55	115.30
1	R	21	PRO	N-CA-CB	7.04	111.75	103.30
1	J	21	PRO	N-CA-CB	6.99	111.69	103.30
1	N	32	LEU	CA-CB-CG	6.59	130.47	115.30
1	M	43	LEU	CA-CB-CG	6.53	130.31	115.30

There are no chirality outliers.

5 of 11 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	25	GLY	Peptide
1	E	16	LEU	Peptide
1	E	25	GLY	Peptide
1	H	142	GLU	Peptide
1	I	22	ALA	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1065	0	1068	21	0
1	B	1071	0	1079	17	0
1	C	1065	0	1069	18	0
1	D	1069	0	1072	38	0
1	E	1071	0	1079	42	0
1	F	1071	0	1071	24	0
1	G	1071	0	1079	35	0
1	H	1071	0	1079	27	0
1	I	1071	0	1079	21	0
1	J	1065	0	1069	38	0
1	K	1061	0	1064	30	0
1	L	1071	0	1079	21	0
1	M	1071	0	1079	23	0
1	N	1077	0	1082	32	0
1	O	1071	0	1079	30	0
1	P	1071	0	1079	25	0
1	Q	1065	0	1068	19	0
1	R	1059	0	1058	32	0
1	S	1071	0	1079	28	0
1	T	1071	0	1079	12	0
1	U	1071	0	1079	24	0
1	V	1067	0	1075	39	0
1	W	1065	0	1068	16	0
1	X	1061	0	1064	28	0
2	A	26	0	17	1	0
2	B	26	0	17	1	0
2	C	26	0	17	4	0
2	D	26	0	17	0	0
2	E	26	0	17	3	0
2	F	26	0	17	1	0
2	G	26	0	17	7	0
2	H	26	0	17	0	0
2	I	26	0	17	5	0
2	J	26	0	17	2	0
2	K	26	0	17	1	0
2	L	26	0	17	8	0
2	M	26	0	17	6	0
2	N	26	0	17	2	0
2	O	26	0	17	3	0
2	P	26	0	17	0	0
2	Q	26	0	17	3	0
2	R	26	0	17	3	0
2	S	26	0	17	4	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	T	26	0	17	0	0
2	U	26	0	17	5	0
2	V	26	0	17	4	0
2	W	26	0	17	2	0
2	X	26	0	17	0	0
3	A	53	0	0	2	0
3	B	90	0	0	7	0
3	C	59	0	0	1	0
3	D	54	0	0	2	0
3	E	78	0	0	2	0
3	F	60	0	0	3	0
3	G	66	0	0	5	0
3	H	59	0	0	2	0
3	I	48	0	0	0	0
3	J	70	0	0	10	0
3	K	55	0	0	3	0
3	L	49	0	0	1	0
3	M	57	0	0	0	0
3	N	54	0	0	2	0
3	O	53	0	0	7	0
3	P	67	0	0	2	0
3	Q	50	0	0	2	0
3	R	63	0	0	1	0
3	S	66	0	0	3	0
3	T	73	0	0	0	0
3	U	59	0	0	1	0
3	V	61	0	0	1	0
3	W	43	0	0	1	0
3	X	48	0	0	3	0
All	All	27701	0	26184	647	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

The worst 5 of 647 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:F:21:PRO:HA	1:F:22:ALA:O	1.46	1.13
2:I:147:N87:H22	2:I:147:N87:H23	1.19	1.12
2:M:147:N87:H23	2:M:147:N87:H22	1.13	1.11
1:Q:38:ARG:HG2	1:Q:38:ARG:HH11	1.11	1.10

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:J:18:ARG:HH11	1:J:18:ARG:HG3	0.93	1.10

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	139/147 (95%)	126 (91%)	11 (8%)	2 (1%)	14	19
1	B	139/147 (95%)	130 (94%)	9 (6%)	0	100	100
1	C	139/147 (95%)	128 (92%)	7 (5%)	4 (3%)	6	5
1	D	139/147 (95%)	128 (92%)	8 (6%)	3 (2%)	8	9
1	E	139/147 (95%)	127 (91%)	9 (6%)	3 (2%)	8	9
1	F	140/147 (95%)	132 (94%)	4 (3%)	4 (3%)	6	5
1	G	139/147 (95%)	131 (94%)	6 (4%)	2 (1%)	14	19
1	H	139/147 (95%)	132 (95%)	7 (5%)	0	100	100
1	I	139/147 (95%)	132 (95%)	6 (4%)	1 (1%)	26	38
1	J	139/147 (95%)	130 (94%)	5 (4%)	4 (3%)	6	5
1	K	139/147 (95%)	124 (89%)	11 (8%)	4 (3%)	6	5
1	L	139/147 (95%)	131 (94%)	5 (4%)	3 (2%)	8	9
1	M	139/147 (95%)	134 (96%)	5 (4%)	0	100	100
1	N	140/147 (95%)	122 (87%)	13 (9%)	5 (4%)	4	3
1	O	139/147 (95%)	126 (91%)	12 (9%)	1 (1%)	26	38
1	P	139/147 (95%)	130 (94%)	7 (5%)	2 (1%)	14	19
1	Q	139/147 (95%)	127 (91%)	8 (6%)	4 (3%)	6	5
1	R	139/147 (95%)	123 (88%)	11 (8%)	5 (4%)	4	3
1	S	139/147 (95%)	130 (94%)	5 (4%)	4 (3%)	6	5

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	T	139/147 (95%)	132 (95%)	6 (4%)	1 (1%)	26	38
1	U	139/147 (95%)	126 (91%)	9 (6%)	4 (3%)	6	5
1	V	139/147 (95%)	125 (90%)	6 (4%)	8 (6%)	2	1
1	W	139/147 (95%)	129 (93%)	8 (6%)	2 (1%)	14	19
1	X	139/147 (95%)	126 (91%)	10 (7%)	3 (2%)	8	9
All	All	3338/3528 (95%)	3081 (92%)	188 (6%)	69 (2%)	9	10

5 of 69 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	19	ARG
1	C	20	GLU
1	C	21	PRO
1	D	20	GLU
1	D	21	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	109/115 (95%)	102 (94%)	7 (6%)	22	34
1	B	110/115 (96%)	104 (94%)	6 (6%)	27	42
1	C	108/115 (94%)	101 (94%)	7 (6%)	21	33
1	D	109/115 (95%)	97 (89%)	12 (11%)	8	10
1	E	110/115 (96%)	103 (94%)	7 (6%)	22	34
1	F	110/115 (96%)	106 (96%)	4 (4%)	42	63
1	G	110/115 (96%)	100 (91%)	10 (9%)	12	17
1	H	110/115 (96%)	99 (90%)	11 (10%)	9	14
1	I	110/115 (96%)	102 (93%)	8 (7%)	17	27
1	J	108/115 (94%)	99 (92%)	9 (8%)	14	21
1	K	108/115 (94%)	100 (93%)	8 (7%)	17	26

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	L	110/115 (96%)	102 (93%)	8 (7%)	17	27
1	M	110/115 (96%)	98 (89%)	12 (11%)	8	11
1	N	111/115 (96%)	102 (92%)	9 (8%)	15	22
1	O	110/115 (96%)	101 (92%)	9 (8%)	14	21
1	P	110/115 (96%)	100 (91%)	10 (9%)	12	17
1	Q	109/115 (95%)	102 (94%)	7 (6%)	22	34
1	R	107/115 (93%)	100 (94%)	7 (6%)	21	33
1	S	110/115 (96%)	101 (92%)	9 (8%)	14	21
1	T	110/115 (96%)	103 (94%)	7 (6%)	22	34
1	U	110/115 (96%)	103 (94%)	7 (6%)	22	34
1	V	109/115 (95%)	93 (85%)	16 (15%)	4	4
1	W	109/115 (95%)	101 (93%)	8 (7%)	17	27
1	X	108/115 (94%)	99 (92%)	9 (8%)	14	21
All	All	2625/2760 (95%)	2418 (92%)	207 (8%)	15	23

5 of 207 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	L	54	SER
1	N	70	GLU
1	W	20	GLU
1	L	143	HIS
1	M	131	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 36 such sidechains are listed below:

Mol	Chain	Res	Type
1	M	131	GLN
1	P	63	HIS
1	X	64	GLN
1	O	63	HIS
1	P	101	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

24 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	N87	A	147	-	22,28,28	0.84	0	21,41,41	1.43	2 (9%)
2	N87	B	147	-	22,28,28	0.68	0	21,41,41	1.09	1 (4%)
2	N87	C	147	-	22,28,28	1.20	2 (9%)	21,41,41	0.85	1 (4%)
2	N87	D	147	-	22,28,28	1.66	5 (22%)	21,41,41	2.24	8 (38%)
2	N87	E	147	-	22,28,28	0.75	0	21,41,41	0.92	1 (4%)
2	N87	F	147	-	22,28,28	1.14	2 (9%)	21,41,41	0.92	0
2	N87	G	147	-	22,28,28	0.59	0	21,41,41	1.10	0
2	N87	H	147	-	22,28,28	0.90	1 (4%)	21,41,41	1.76	6 (28%)
2	N87	I	147	-	22,28,28	0.89	0	21,41,41	0.88	1 (4%)
2	N87	J	147	-	22,28,28	1.04	1 (4%)	21,41,41	0.70	0
2	N87	K	147	-	22,28,28	1.49	3 (13%)	21,41,41	1.56	3 (14%)
2	N87	L	147	-	22,28,28	1.56	3 (13%)	21,41,41	0.97	0
2	N87	M	147	-	22,28,28	1.30	2 (9%)	21,41,41	0.84	1 (4%)
2	N87	N	147	-	22,28,28	1.22	2 (9%)	21,41,41	1.47	3 (14%)
2	N87	O	147	-	22,28,28	1.02	0	21,41,41	1.12	2 (9%)
2	N87	P	147	-	22,28,28	1.14	1 (4%)	21,41,41	1.78	4 (19%)
2	N87	Q	147	-	22,28,28	0.70	0	21,41,41	1.66	7 (33%)
2	N87	R	147	-	22,28,28	1.03	0	21,41,41	1.21	3 (14%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	N87	S	147	-	22,28,28	1.07	2 (9%)	21,41,41	0.74	0
2	N87	T	147	-	22,28,28	0.72	0	21,41,41	1.50	4 (19%)
2	N87	U	147	-	22,28,28	1.00	0	21,41,41	1.14	1 (4%)
2	N87	V	147	-	22,28,28	1.21	3 (13%)	21,41,41	1.38	4 (19%)
2	N87	W	147	-	22,28,28	1.36	3 (13%)	21,41,41	1.14	1 (4%)
2	N87	X	147	-	22,28,28	1.51	3 (13%)	21,41,41	1.39	3 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	N87	A	147	-	-	0/12/36/36	0/3/3/3
2	N87	B	147	-	-	0/12/36/36	0/3/3/3
2	N87	C	147	-	-	0/12/36/36	0/3/3/3
2	N87	D	147	-	-	0/12/36/36	0/3/3/3
2	N87	E	147	-	-	0/12/36/36	0/3/3/3
2	N87	F	147	-	-	0/12/36/36	0/3/3/3
2	N87	G	147	-	-	0/12/36/36	0/3/3/3
2	N87	H	147	-	-	0/12/36/36	0/3/3/3
2	N87	I	147	-	-	0/12/36/36	0/3/3/3
2	N87	J	147	-	-	0/12/36/36	0/3/3/3
2	N87	K	147	-	-	0/12/36/36	0/3/3/3
2	N87	L	147	-	-	0/12/36/36	0/3/3/3
2	N87	M	147	-	-	0/12/36/36	0/3/3/3
2	N87	N	147	-	-	0/12/36/36	0/3/3/3
2	N87	O	147	-	-	0/12/36/36	0/3/3/3
2	N87	P	147	-	-	0/12/36/36	0/3/3/3
2	N87	Q	147	-	-	0/12/36/36	0/3/3/3
2	N87	R	147	-	-	0/12/36/36	0/3/3/3
2	N87	S	147	-	-	0/12/36/36	0/3/3/3
2	N87	T	147	-	-	0/12/36/36	0/3/3/3
2	N87	U	147	-	-	0/12/36/36	0/3/3/3
2	N87	V	147	-	-	0/12/36/36	0/3/3/3
2	N87	W	147	-	-	0/12/36/36	0/3/3/3
2	N87	X	147	-	-	0/12/36/36	0/3/3/3

The worst 5 of 33 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	H	147	N87	C4-C6	2.03	1.52	1.50
2	P	147	N87	C13-C7	2.06	1.51	1.48
2	D	147	N87	C15-C16	2.06	1.53	1.49
2	N	147	N87	C13-C7	2.10	1.51	1.48
2	S	147	N87	C6-C7	2.10	1.36	1.34

The worst 5 of 56 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	147	N87	O26-C16-C17	-5.25	111.73	120.12
2	P	147	N87	O26-C16-C17	-4.45	113.01	120.12
2	H	147	N87	O26-C16-C17	-3.83	114.00	120.12
2	Q	147	N87	C25-C13-C7	-3.74	117.11	121.08
2	U	147	N87	C25-C13-C7	-3.35	117.53	121.08

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

19 monomers are involved in 65 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	147	N87	1	0
2	B	147	N87	1	0
2	C	147	N87	4	0
2	E	147	N87	3	0
2	F	147	N87	1	0
2	G	147	N87	7	0
2	I	147	N87	5	0
2	J	147	N87	2	0
2	K	147	N87	1	0
2	L	147	N87	8	0
2	M	147	N87	6	0
2	N	147	N87	2	0
2	O	147	N87	3	0
2	Q	147	N87	3	0
2	R	147	N87	3	0
2	S	147	N87	4	0
2	U	147	N87	5	0
2	V	147	N87	4	0
2	W	147	N87	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	141/147 (95%)	0.01	3 (2%) 67 66	25, 37, 56, 68	2 (1%)
1	B	141/147 (95%)	-0.22	0 100 100	13, 26, 41, 50	1 (0%)
1	C	141/147 (95%)	0.10	6 (4%) 39 40	18, 30, 53, 62	0
1	D	141/147 (95%)	0.21	9 (6%) 23 23	20, 35, 54, 65	0
1	E	141/147 (95%)	0.12	6 (4%) 39 40	19, 31, 53, 64	0
1	F	141/147 (95%)	0.15	7 (4%) 32 33	20, 31, 55, 63	0
1	G	141/147 (95%)	-0.26	1 (0%) 89 88	17, 28, 46, 62	1 (0%)
1	H	141/147 (95%)	-0.29	1 (0%) 89 88	13, 30, 45, 53	0
1	I	141/147 (95%)	0.09	6 (4%) 39 40	21, 33, 57, 68	0
1	J	141/147 (95%)	-0.20	2 (1%) 78 77	17, 27, 49, 63	0
1	K	141/147 (95%)	0.06	7 (4%) 32 33	13, 34, 55, 61	0
1	L	141/147 (95%)	-0.03	7 (4%) 32 33	20, 35, 50, 58	0
1	M	141/147 (95%)	-0.01	6 (4%) 39 40	22, 32, 52, 58	0
1	N	141/147 (95%)	-0.02	6 (4%) 39 40	24, 35, 54, 63	0
1	O	141/147 (95%)	0.04	6 (4%) 39 40	20, 30, 54, 67	0
1	P	141/147 (95%)	-0.28	0 100 100	9, 27, 42, 56	0
1	Q	141/147 (95%)	-0.08	4 (2%) 56 55	20, 30, 49, 60	0
1	R	141/147 (95%)	0.04	7 (4%) 32 33	21, 31, 54, 62	0
1	S	141/147 (95%)	-0.25	5 (3%) 48 48	13, 26, 45, 59	2 (1%)
1	T	141/147 (95%)	-0.36	0 100 100	17, 27, 42, 54	1 (0%)
1	U	141/147 (95%)	-0.00	5 (3%) 48 48	17, 29, 51, 59	1 (0%)
1	V	141/147 (95%)	0.03	4 (2%) 56 55	20, 29, 51, 63	1 (0%)
1	W	141/147 (95%)	-0.03	6 (4%) 39 40	20, 36, 56, 62	1 (0%)
1	X	141/147 (95%)	0.04	7 (4%) 32 33	22, 33, 54, 64	1 (0%)

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å ²)	Q<0.9
All	All	3384/3528 (95%)	-0.05	111 (3%)	50	50	9, 31, 54, 68	11 (0%)

The worst 5 of 111 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	R	22	ALA	6.2
1	E	22	ALA	5.1
1	X	20	GLU	5.0
1	J	23	VAL	4.3
1	F	21	PRO	4.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	N87	A	147	26/26	0.92	0.17	0.83	36,43,59,59	0
2	N87	T	147	26/26	0.95	0.14	0.65	28,35,47,48	0
2	N87	E	147	26/26	0.93	0.17	0.49	33,41,60,61	0
2	N87	M	147	26/26	0.90	0.17	0.48	31,44,59,60	0
2	N87	D	147	26/26	0.89	0.18	0.39	27,38,55,57	0
2	N87	X	147	26/26	0.86	0.20	0.35	39,51,58,59	0
2	N87	O	147	26/26	0.90	0.17	0.32	40,46,54,55	0
2	N87	I	147	26/26	0.88	0.18	0.30	35,46,66,68	0
2	N87	Q	147	26/26	0.91	0.18	0.22	34,40,49,51	0
2	N87	R	147	26/26	0.90	0.18	0.19	35,45,63,64	0
2	N87	K	147	26/26	0.89	0.17	0.19	29,42,56,57	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	N87	N	147	26/26	0.89	0.17	0.16	26,38,55,57	0
2	N87	F	147	26/26	0.91	0.18	0.08	36,42,55,55	0
2	N87	C	147	26/26	0.91	0.17	0.06	26,40,58,59	0
2	N87	V	147	26/26	0.93	0.16	0.05	27,43,55,57	0
2	N87	L	147	26/26	0.91	0.15	0.01	27,36,47,49	0
2	N87	W	147	26/26	0.93	0.15	-0.01	31,42,59,60	0
2	N87	J	147	26/26	0.93	0.16	-0.02	28,39,56,58	0
2	N87	B	147	26/26	0.96	0.13	-0.18	23,28,37,38	0
2	N87	U	147	26/26	0.94	0.15	-0.27	24,38,57,59	0
2	N87	G	147	26/26	0.95	0.13	-0.32	31,39,50,51	0
2	N87	H	147	26/26	0.96	0.11	-0.33	18,29,30,32	0
2	N87	S	147	26/26	0.95	0.12	-0.34	17,30,48,49	0
2	N87	P	147	26/26	0.97	0.10	-0.95	15,24,28,29	0

6.5 Other polymers [i](#)

There are no such residues in this entry.