



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 07:09 PM GMT

PDB ID : 4NSQ
Title : Crystal structure of PCAF
Authors : Lin, J.Y.; Cai, Y.F.
Deposited on : 2013-11-28
Resolution : 2.31 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.31 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	4425 (2.34-2.30)
Clashscore	102246	5057 (2.34-2.30)
Ramachandran outliers	100387	5008 (2.34-2.30)
Sidechain outliers	100360	5007 (2.34-2.30)
RSRZ outliers	91569	4432 (2.34-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	190	<div> <div>6%</div> <div>54%</div> <div>24%</div> <div>•</div> <div>17%</div> </div>
1	B	190	<div> <div>6%</div> <div>55%</div> <div>25%</div> <div>••</div> <div>17%</div> </div>
1	C	190	<div> <div>13%</div> <div>54%</div> <div>23%</div> <div>5%</div> <div>17%</div> </div>
1	D	190	<div> <div>14%</div> <div>48%</div> <div>27%</div> <div>7%</div> <div>17%</div> </div>

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 5296 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Histone acetyltransferase KAT2B.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	157	Total	C	N	O	S	0	0	0
			1276	836	213	219	8			
1	B	157	Total	C	N	O	S	0	0	0
			1276	836	213	219	8			
1	C	157	Total	C	N	O	S	0	0	0
			1276	836	213	219	8			
1	D	157	Total	C	N	O	S	0	0	0
			1276	836	213	219	8			

There are 96 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	469	MET	-	EXPRESSION TAG	UNP Q92831
A	470	GLY	-	EXPRESSION TAG	UNP Q92831
A	471	SER	-	EXPRESSION TAG	UNP Q92831
A	472	SER	-	EXPRESSION TAG	UNP Q92831
A	473	HIS	-	EXPRESSION TAG	UNP Q92831
A	474	HIS	-	EXPRESSION TAG	UNP Q92831
A	475	HIS	-	EXPRESSION TAG	UNP Q92831
A	476	HIS	-	EXPRESSION TAG	UNP Q92831
A	477	HIS	-	EXPRESSION TAG	UNP Q92831
A	478	HIS	-	EXPRESSION TAG	UNP Q92831
A	479	SER	-	EXPRESSION TAG	UNP Q92831
A	480	SER	-	EXPRESSION TAG	UNP Q92831
A	481	GLY	-	EXPRESSION TAG	UNP Q92831
A	482	LEU	-	EXPRESSION TAG	UNP Q92831
A	483	VAL	-	EXPRESSION TAG	UNP Q92831
A	484	PRO	-	EXPRESSION TAG	UNP Q92831
A	485	ARG	-	EXPRESSION TAG	UNP Q92831
A	486	GLY	-	EXPRESSION TAG	UNP Q92831
A	487	SER	-	EXPRESSION TAG	UNP Q92831
A	488	HIS	-	EXPRESSION TAG	UNP Q92831
A	489	MET	-	EXPRESSION TAG	UNP Q92831

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	490	ALA	-	EXPRESSION TAG	UNP Q92831
A	491	SER	-	EXPRESSION TAG	UNP Q92831
A	492	LYS	-	EXPRESSION TAG	UNP Q92831
B	469	MET	-	EXPRESSION TAG	UNP Q92831
B	470	GLY	-	EXPRESSION TAG	UNP Q92831
B	471	SER	-	EXPRESSION TAG	UNP Q92831
B	472	SER	-	EXPRESSION TAG	UNP Q92831
B	473	HIS	-	EXPRESSION TAG	UNP Q92831
B	474	HIS	-	EXPRESSION TAG	UNP Q92831
B	475	HIS	-	EXPRESSION TAG	UNP Q92831
B	476	HIS	-	EXPRESSION TAG	UNP Q92831
B	477	HIS	-	EXPRESSION TAG	UNP Q92831
B	478	HIS	-	EXPRESSION TAG	UNP Q92831
B	479	SER	-	EXPRESSION TAG	UNP Q92831
B	480	SER	-	EXPRESSION TAG	UNP Q92831
B	481	GLY	-	EXPRESSION TAG	UNP Q92831
B	482	LEU	-	EXPRESSION TAG	UNP Q92831
B	483	VAL	-	EXPRESSION TAG	UNP Q92831
B	484	PRO	-	EXPRESSION TAG	UNP Q92831
B	485	ARG	-	EXPRESSION TAG	UNP Q92831
B	486	GLY	-	EXPRESSION TAG	UNP Q92831
B	487	SER	-	EXPRESSION TAG	UNP Q92831
B	488	HIS	-	EXPRESSION TAG	UNP Q92831
B	489	MET	-	EXPRESSION TAG	UNP Q92831
B	490	ALA	-	EXPRESSION TAG	UNP Q92831
B	491	SER	-	EXPRESSION TAG	UNP Q92831
B	492	LYS	-	EXPRESSION TAG	UNP Q92831
C	469	MET	-	EXPRESSION TAG	UNP Q92831
C	470	GLY	-	EXPRESSION TAG	UNP Q92831
C	471	SER	-	EXPRESSION TAG	UNP Q92831
C	472	SER	-	EXPRESSION TAG	UNP Q92831
C	473	HIS	-	EXPRESSION TAG	UNP Q92831
C	474	HIS	-	EXPRESSION TAG	UNP Q92831
C	475	HIS	-	EXPRESSION TAG	UNP Q92831
C	476	HIS	-	EXPRESSION TAG	UNP Q92831
C	477	HIS	-	EXPRESSION TAG	UNP Q92831
C	478	HIS	-	EXPRESSION TAG	UNP Q92831
C	479	SER	-	EXPRESSION TAG	UNP Q92831
C	480	SER	-	EXPRESSION TAG	UNP Q92831
C	481	GLY	-	EXPRESSION TAG	UNP Q92831
C	482	LEU	-	EXPRESSION TAG	UNP Q92831
C	483	VAL	-	EXPRESSION TAG	UNP Q92831

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	484	PRO	-	EXPRESSION TAG	UNP Q92831
C	485	ARG	-	EXPRESSION TAG	UNP Q92831
C	486	GLY	-	EXPRESSION TAG	UNP Q92831
C	487	SER	-	EXPRESSION TAG	UNP Q92831
C	488	HIS	-	EXPRESSION TAG	UNP Q92831
C	489	MET	-	EXPRESSION TAG	UNP Q92831
C	490	ALA	-	EXPRESSION TAG	UNP Q92831
C	491	SER	-	EXPRESSION TAG	UNP Q92831
C	492	LYS	-	EXPRESSION TAG	UNP Q92831
D	469	MET	-	EXPRESSION TAG	UNP Q92831
D	470	GLY	-	EXPRESSION TAG	UNP Q92831
D	471	SER	-	EXPRESSION TAG	UNP Q92831
D	472	SER	-	EXPRESSION TAG	UNP Q92831
D	473	HIS	-	EXPRESSION TAG	UNP Q92831
D	474	HIS	-	EXPRESSION TAG	UNP Q92831
D	475	HIS	-	EXPRESSION TAG	UNP Q92831
D	476	HIS	-	EXPRESSION TAG	UNP Q92831
D	477	HIS	-	EXPRESSION TAG	UNP Q92831
D	478	HIS	-	EXPRESSION TAG	UNP Q92831
D	479	SER	-	EXPRESSION TAG	UNP Q92831
D	480	SER	-	EXPRESSION TAG	UNP Q92831
D	481	GLY	-	EXPRESSION TAG	UNP Q92831
D	482	LEU	-	EXPRESSION TAG	UNP Q92831
D	483	VAL	-	EXPRESSION TAG	UNP Q92831
D	484	PRO	-	EXPRESSION TAG	UNP Q92831
D	485	ARG	-	EXPRESSION TAG	UNP Q92831
D	486	GLY	-	EXPRESSION TAG	UNP Q92831
D	487	SER	-	EXPRESSION TAG	UNP Q92831
D	488	HIS	-	EXPRESSION TAG	UNP Q92831
D	489	MET	-	EXPRESSION TAG	UNP Q92831
D	490	ALA	-	EXPRESSION TAG	UNP Q92831
D	491	SER	-	EXPRESSION TAG	UNP Q92831
D	492	LYS	-	EXPRESSION TAG	UNP Q92831

- Molecule 2 is COENZYME A (three-letter code: COA) (formula: C₂₁H₃₆N₇O₁₆P₃S).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
2	A	1	Total 48	C 21	N 7	O 16	P 3	S 1	0	0
2	B	1	Total 48	C 21	N 7	O 16	P 3	S 1	0	0
2	C	1	Total 48	C 21	N 7	O 16	P 3	S 1	0	0
2	D	1	Total 48	C 21	N 7	O 16	P 3	S 1	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Histone acetyltransferase KAT2B




• Molecule 1: Histone acetyltransferase KAT2B




• Molecule 1: Histone acetyltransferase KAT2B





● Molecule 1: Histone acetyltransferase KAT2B



4 Data and refinement statistics

Property	Value	Source
Space group	P 43	Depositor
Cell constants a, b, c, α , β , γ	65.57Å 65.57Å 187.62Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	45.26 – 2.31 45.26 – 2.31	Depositor EDS
% Data completeness (in resolution range)	(Not available) (45.26-2.31) 88.1 (45.26-2.31)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.68 (at 2.32Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.7.1_743)	Depositor
R, R_{free}	0.229 , 0.275 0.250 , 0.302	Depositor DCC
R_{free} test set	1988 reflections (6.96%)	DCC
Wilson B-factor (Å ²)	57.5	Xtriage
Anisotropy	0.487	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 58.3	EDS
Estimated twinning fraction	0.065 for h,-k,-l	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Outliers	0 of 32806 reflections	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	5296	wwPDB-VP
Average B, all atoms (Å ²)	81.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.04% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: COA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.71	0/1309	0.61	0/1760
1	B	0.78	2/1309 (0.2%)	0.59	0/1760
1	C	0.68	0/1309	0.57	0/1760
1	D	0.58	0/1309	0.55	0/1760
All	All	0.69	2/5236 (0.0%)	0.58	0/7040

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	559	CYS	CB-SG	5.40	1.91	1.82
1	B	570	GLU	CG-CD	5.16	1.59	1.51

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1276	0	1297	44	0
1	B	1276	0	1297	46	0
1	C	1276	0	1297	38	0
1	D	1276	0	1297	74	0
2	A	48	0	32	3	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	48	0	32	4	0
2	C	48	0	32	8	0
2	D	48	0	32	10	0
All	All	5296	0	5316	205	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 19.

The worst 5 of 205 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:651:ASN:HB2	1:B:652:PRO:CD	1.42	1.49
1:A:651:ASN:HB2	1:A:652:PRO:CD	1.59	1.32
1:C:651:ASN:HB2	1:C:652:PRO:CD	1.62	1.29
1:A:525:GLN:O	1:A:527:PRO:HD3	1.28	1.25
1:B:651:ASN:CB	1:B:652:PRO:HD2	1.67	1.22

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	153/190 (80%)	146 (95%)	5 (3%)	2 (1%)	15	14
1	B	153/190 (80%)	145 (95%)	6 (4%)	2 (1%)	15	14
1	C	153/190 (80%)	147 (96%)	6 (4%)	0	100	100
1	D	153/190 (80%)	145 (95%)	7 (5%)	1 (1%)	26	31
All	All	612/760 (80%)	583 (95%)	24 (4%)	5 (1%)	24	27

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	651	ASN
1	A	526	LEU
1	B	565	SER
1	A	651	ASN
1	D	582	VAL

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	139/168 (83%)	118 (85%)	21 (15%)	3	3
1	B	139/168 (83%)	128 (92%)	11 (8%)	15	19
1	C	139/168 (83%)	118 (85%)	21 (15%)	3	3
1	D	139/168 (83%)	117 (84%)	22 (16%)	3	3
All	All	556/672 (83%)	481 (86%)	75 (14%)	5	4

5 of 75 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	494	ILE
1	C	553	ARG
1	D	626	ILE
1	C	502	SER
1	C	523	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	508	ASN
1	C	597	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	COA	A	700	-	40,50,50	2.18	7 (17%)	50,75,75	2.13	11 (22%)
2	COA	B	700	-	40,50,50	2.12	8 (20%)	50,75,75	2.18	14 (28%)
2	COA	C	700	-	40,50,50	2.17	8 (20%)	50,75,75	2.32	10 (20%)
2	COA	D	700	-	40,50,50	2.12	8 (20%)	50,75,75	2.21	14 (28%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	COA	A	700	-	-	0/44/64/64	0/3/3/3
2	COA	B	700	-	-	0/44/64/64	0/3/3/3
2	COA	C	700	-	-	0/44/64/64	0/3/3/3
2	COA	D	700	-	-	0/44/64/64	0/3/3/3

The worst 5 of 31 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	700	COA	C2B-C3B	-5.13	1.41	1.53
2	C	700	COA	C2B-C3B	-4.98	1.41	1.53

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	700	COA	C2B-C3B	-4.88	1.42	1.53
2	D	700	COA	C2B-C3B	-4.80	1.42	1.53
2	A	700	COA	C3B-C4B	-3.38	1.43	1.52

The worst 5 of 49 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	700	COA	N3A-C2A-N1A	-11.32	120.23	128.89
2	D	700	COA	N3A-C2A-N1A	-11.11	120.39	128.89
2	B	700	COA	N3A-C2A-N1A	-10.11	121.15	128.89
2	A	700	COA	N3A-C2A-N1A	-10.11	121.15	128.89
2	C	700	COA	P2A-O3A-P1A	-5.17	118.21	132.73

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 25 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	700	COA	3	0
2	B	700	COA	4	0
2	C	700	COA	8	0
2	D	700	COA	10	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	157/190 (82%)	0.90	12 (7%) 17 24	53, 72, 104, 122	0
1	B	157/190 (82%)	0.94	11 (7%) 19 27	24, 69, 100, 134	0
1	C	157/190 (82%)	1.02	25 (15%) 3 4	53, 82, 114, 144	0
1	D	157/190 (82%)	1.39	26 (16%) 2 4	54, 87, 126, 158	0
All	All	628/760 (82%)	1.06	74 (11%) 6 10	24, 77, 115, 158	0

The worst 5 of 74 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	652	PRO	12.7
1	D	650	LEU	11.3
1	D	651	ASN	10.0
1	A	652	PRO	7.7
1	B	502	SER	7.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	COA	C	700	48/48	0.91	0.22	0.95	60,82,122,136	0
2	COA	B	700	48/48	0.89	0.19	0.40	59,78,99,112	0
2	COA	A	700	48/48	0.92	0.19	0.01	64,77,98,103	0
2	COA	D	700	48/48	0.89	0.19	-0.01	62,94,114,121	0

6.5 Other polymers [i](#)

There are no such residues in this entry.