



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 11:08 AM GMT

PDB ID : 3O7X
Title : Crystal structure of human Hili PAZ domain
Authors : Tian, Y.; Simanshu, D.K.; Ma, J.-B.; Patel, D.J.
Deposited on : 2010-08-01
Resolution : 2.92 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.92 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1643 (2.94-2.90)
Clashscore	102246	1871 (2.94-2.90)
Ramachandran outliers	100387	1824 (2.94-2.90)
Sidechain outliers	100360	1826 (2.94-2.90)
RSRZ outliers	91569	1650 (2.94-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	140	<div> <div>%</div> <div> <div></div> <div>61%</div> <div>16%</div> <div>• •</div> <div>17%</div> </div> </div>
1	B	140	<div> <div>%</div> <div> <div></div> <div>53%</div> <div>19%</div> <div>•</div> <div>26%</div> </div> </div>
1	C	140	<div> <div></div> <div> <div></div> <div>59%</div> <div>15%</div> <div>•</div> <div>24%</div> </div> </div>
1	D	140	<div> <div>%</div> <div> <div></div> <div>57%</div> <div>15%</div> <div>6%</div> <div>•</div> <div>21%</div> </div> </div>

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 3587 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Piwi-like protein 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	116	Total	C	N	O	S	0	0	0
			950	598	159	187	6			
1	B	104	Total	C	N	O	S	0	0	0
			856	543	140	168	5			
1	C	107	Total	C	N	O	S	0	0	0
			872	552	143	172	5			
1	D	110	Total	C	N	O	S	0	0	0
			900	571	150	174	5			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	386	SER	-	EXPRESSION TAG	UNP Q8TC59
B	386	SER	-	EXPRESSION TAG	UNP Q8TC59
C	386	SER	-	EXPRESSION TAG	UNP Q8TC59
D	386	SER	-	EXPRESSION TAG	UNP Q8TC59


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	3	Total	O	0	0
			3	3		
2	B	1	Total	O	0	0
			1	1		
2	C	1	Total	O	0	0
			1	1		
2	D	4	Total	O	0	0
			4	4		


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Piwi-like protein 2




• Molecule 1: Piwi-like protein 2



• Molecule 1: Piwi-like protein 2



• Molecule 1: Piwi-like protein 2



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	40.53Å 142.95Å 226.51Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	44.38 – 2.92 44.39 – 2.92	Depositor EDS
% Data completeness (in resolution range)	90.2 (44.38-2.92) 90.3 (44.39-2.92)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	5.89 (at 2.90Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.4_153)	Depositor
R, R_{free}	0.246 , 0.293 0.244 , 0.288	Depositor DCC
R_{free} test set	1341 reflections (9.96%)	DCC
Wilson B-factor (Å ²)	60.1	Xtriage
Anisotropy	0.712	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 45.1	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 14056 reflections	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	3587	wwPDB-VP
Average B, all atoms (Å ²)	75.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.59% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.31	0/969	0.93	3/1312 (0.2%)
1	B	0.29	0/873	0.49	1/1182 (0.1%)
1	C	0.31	0/889	0.61	2/1204 (0.2%)
1	D	0.28	0/917	0.54	2/1240 (0.2%)
All	All	0.30	0/3648	0.67	8/4938 (0.2%)

There are no bond length outliers.

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	421	ARG	NE-CZ-NH1	-20.61	110.00	120.30
1	A	421	ARG	NE-CZ-NH2	18.07	129.34	120.30
1	C	421	ARG	NE-CZ-NH2	-10.58	115.01	120.30
1	A	421	ARG	CD-NE-CZ	9.78	137.29	123.60
1	C	421	ARG	NE-CZ-NH1	8.91	124.76	120.30
1	D	421	ARG	NE-CZ-NH1	5.69	123.14	120.30
1	D	421	ARG	NE-CZ-NH2	-5.69	117.46	120.30
1	B	421	ARG	CB-CG-CD	5.35	125.52	111.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	950	0	910	17	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	856	0	824	21	0
1	C	872	0	833	16	0
1	D	900	0	869	23	0
2	A	3	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	4	0	0	0	0
All	All	3587	0	3436	72	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (72) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:480:GLN:HG3	1:D:481:ASP:H	1.19	1.06
1:D:479:ARG:HG2	1:D:479:ARG:HH11	1.34	0.92
1:D:480:GLN:HE21	1:D:480:GLN:HA	1.53	0.72
1:B:401:ASN:HD21	1:C:404:HIS:HD2	1.37	0.72
1:D:480:GLN:HG3	1:D:481:ASP:N	2.02	0.69
1:A:481:ASP:O	1:A:483:HIS:N	2.24	0.69
1:A:437:THR:HG23	1:A:439:LYS:H	1.58	0.69
1:B:437:THR:HG23	1:B:439:LYS:H	1.58	0.68
1:D:437:THR:HG23	1:D:439:LYS:H	1.59	0.67
1:C:437:THR:HG23	1:C:439:LYS:H	1.59	0.67
1:A:483:HIS:CD2	1:A:483:HIS:N	2.64	0.65
1:D:479:ARG:HH11	1:D:479:ARG:CG	2.08	0.63
1:B:401:ASN:HD21	1:C:404:HIS:CD2	2.17	0.62
1:D:479:ARG:O	1:D:480:GLN:HB2	2.01	0.61
1:A:482:ASN:O	1:A:483:HIS:C	2.42	0.58
1:A:412:LEU:HD23	1:A:413:LEU:HD13	1.88	0.55
1:D:412:LEU:HD23	1:D:413:LEU:HD13	1.89	0.55
1:D:452:PHE:O	1:D:456:TYR:HD1	1.90	0.54
1:B:411:LYS:HG2	1:B:411:LYS:O	2.07	0.54
1:D:480:GLN:O	1:D:481:ASP:CB	2.55	0.54
1:B:452:PHE:O	1:B:456:TYR:HD1	1.91	0.54
1:B:412:LEU:HD23	1:B:413:LEU:HD13	1.90	0.54
1:B:413:LEU:HD21	1:C:390:VAL:HG13	1.89	0.53
1:A:452:PHE:O	1:A:456:TYR:HD1	1.92	0.53
1:C:452:PHE:O	1:C:456:TYR:HD1	1.91	0.52
1:C:412:LEU:HD23	1:C:413:LEU:HD13	1.91	0.52

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:479:ARG:HG2	1:D:479:ARG:NH1	2.14	0.51
1:D:433:ASP:HB2	1:D:473:ILE:HD11	1.92	0.51
1:B:420:THR:HB	1:B:422:TYR:CE2	2.47	0.49
1:A:433:ASP:HB2	1:A:473:ILE:HD11	1.94	0.49
1:A:420:THR:HB	1:A:422:TYR:CE2	2.47	0.49
1:B:387:ASN:OD1	1:B:387:ASN:O	2.31	0.48
1:C:433:ASP:HB2	1:C:473:ILE:HD11	1.94	0.48
1:D:457:SER:O	1:D:461:GLY:HA2	2.14	0.48
1:B:400:GLN:C	1:B:402:LYS:H	2.16	0.47
1:D:420:THR:HB	1:D:422:TYR:CE2	2.49	0.47
1:D:426:THR:O	1:D:426:THR:HG23	2.15	0.47
1:C:420:THR:HB	1:C:422:TYR:CE2	2.49	0.46
1:B:433:ASP:HB2	1:B:473:ILE:HD11	1.96	0.46
1:D:413:LEU:HD12	1:D:413:LEU:HA	1.77	0.46
1:C:459:ASN:HD22	1:C:459:ASN:N	2.14	0.46
1:B:457:SER:O	1:B:461:GLY:HA2	2.16	0.45
1:B:401:ASN:ND2	1:C:404:HIS:HD2	2.07	0.45
1:C:413:LEU:HD12	1:C:413:LEU:HA	1.78	0.45
1:D:480:GLN:CG	1:D:481:ASP:H	2.02	0.44
1:A:457:SER:O	1:A:461:GLY:HA2	2.16	0.44
1:D:459:ASN:HD22	1:D:459:ASN:N	2.15	0.44
1:A:426:THR:HG23	1:A:426:THR:O	2.16	0.44
1:B:466:GLU:HG3	1:C:391:LEU:HD21	1.99	0.44
1:A:480:GLN:O	1:A:482:ASN:N	2.51	0.44
1:C:457:SER:O	1:C:461:GLY:HA2	2.16	0.44
1:A:459:ASN:N	1:A:459:ASN:HD22	2.14	0.43
1:D:418:VAL:HG12	1:D:499:PHE:O	2.18	0.43
1:B:397:ILE:O	1:B:401:ASN:HB2	2.19	0.43
1:B:411:LYS:HB3	1:B:411:LYS:HE2	1.76	0.42
1:C:418:VAL:HG12	1:C:499:PHE:O	2.19	0.42
1:B:389:CYS:HB3	1:B:392:ASP:OD2	2.19	0.42
1:B:413:LEU:HA	1:B:413:LEU:HD12	1.79	0.42
1:A:387:ASN:O	1:A:388:ASP:HB2	2.20	0.42
1:B:459:ASN:N	1:B:459:ASN:HD22	2.17	0.42
1:A:485:MET:HG2	1:A:486:LEU:N	2.35	0.42
1:D:480:GLN:CG	1:D:481:ASP:N	2.74	0.42
1:D:397:ILE:HA	1:D:397:ILE:HD13	1.82	0.41
1:A:480:GLN:O	1:A:487:LEU:HD13	2.19	0.41
1:D:479:ARG:C	1:D:487:LEU:HD22	2.41	0.41
1:B:426:THR:O	1:B:426:THR:HG23	2.18	0.41
1:D:476:PRO:O	1:D:477:SER:HB3	2.21	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:426:THR:HG23	1:C:426:THR:O	2.20	0.41
1:A:389:CYS:HB3	1:A:392:ASP:OD2	2.21	0.40
1:B:395:HIS:O	1:B:398:TYR:HB3	2.22	0.40
1:A:391:LEU:O	1:A:391:LEU:HD12	2.21	0.40
1:C:444:MET:HE3	1:C:450:ILE:HG21	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	114/140 (81%)	95 (83%)	14 (12%)	5 (4%)	3	11
1	B	100/140 (71%)	90 (90%)	9 (9%)	1 (1%)	19	53
1	C	103/140 (74%)	95 (92%)	8 (8%)	0	100	100
1	D	106/140 (76%)	91 (86%)	9 (8%)	6 (6%)	2	6
All	All	423/560 (76%)	371 (88%)	40 (10%)	12 (3%)	6	23

All (12) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	482	ASN
1	A	483	HIS
1	D	477	SER
1	D	480	GLN
1	D	476	PRO
1	D	478	GLU
1	A	481	ASP
1	D	389	CYS
1	A	388	ASP
1	A	489	GLY

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	490	GLU
1	D	390	VAL

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	108/131 (82%)	99 (92%)	9 (8%)	14	38
1	B	98/131 (75%)	92 (94%)	6 (6%)	23	55
1	C	99/131 (76%)	94 (95%)	5 (5%)	29	64
1	D	101/131 (77%)	91 (90%)	10 (10%)	10	29
All	All	406/524 (78%)	376 (93%)	30 (7%)	17	44

All (30) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	413	LEU
1	A	418	VAL
1	A	431	ASP
1	A	459	ASN
1	A	477	SER
1	A	481	ASP
1	A	483	HIS
1	A	485	MET
1	A	486	LEU
1	B	391	LEU
1	B	401	ASN
1	B	413	LEU
1	B	418	VAL
1	B	431	ASP
1	B	459	ASN
1	C	387	ASN
1	C	413	LEU
1	C	418	VAL
1	C	431	ASP

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	459	ASN
1	D	399	GLN
1	D	401	ASN
1	D	413	LEU
1	D	418	VAL
1	D	421	ARG
1	D	431	ASP
1	D	459	ASN
1	D	479	ARG
1	D	480	GLN
1	D	487	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (16) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	416	ASN
1	A	459	ASN
1	A	480	GLN
1	A	482	ASN
1	A	483	HIS
1	B	416	ASN
1	B	459	ASN
1	C	387	ASN
1	C	404	HIS
1	C	416	ASN
1	C	459	ASN
1	D	399	GLN
1	D	401	ASN
1	D	416	ASN
1	D	459	ASN
1	D	480	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	116/140 (82%)	0.25	1 (0%) 85 84	50, 74, 93, 106	0
1	B	104/140 (74%)	0.42	2 (1%) 70 66	50, 74, 94, 110	0
1	C	107/140 (76%)	0.34	0 100 100	53, 72, 98, 127	0
1	D	110/140 (78%)	0.40	1 (0%) 85 84	50, 77, 103, 119	0
All	All	437/560 (78%)	0.35	4 (0%) 85 84	50, 74, 98, 127	0

All (4) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	422	TYR	2.2
1	A	400	GLN	2.1
1	D	413	LEU	2.1
1	B	424	ASN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers

There are no such residues in this entry.