



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 07:15 PM GMT

PDB ID : 4OBR
Title : Crystal structure of human alpha-L-iduronidase complex with alpha-L-iduronic acid
Authors : Bie, H.Y.; Yin, J.; He, X.; Kermode, A.R.; James, M.N.G.
Deposited on : 2014-01-07
Resolution : 2.46 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 2.46 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	4776 (2.50-2.42)
Clashscore	102246	1030 (2.48-2.44)
Ramachandran outliers	100387	1024 (2.48-2.44)
Sidechain outliers	100360	1024 (2.48-2.44)
RSRZ outliers	91569	4787 (2.50-2.42)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	627	 3% 90% 5%
1	B	627	 % 93%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	NAG	A	902	-	-	-	X
5	GOL	A	909	-	-	-	X
5	GOL	A	910	-	-	-	X
5	GOL	A	911	-	-	-	X
5	GOL	B	913	-	-	-	X
6	TLA	A	912	-	-	-	X
7	IDR	A	913	-	-	-	X
7	IDR	B	921	-	-	-	X

2 Entry composition [i](#)

There are 11 unique types of molecules in this entry. The entry contains 10073 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Alpha-L-iduronidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	595	Total	C	N	O	S	0	2	0
			4746	3036	854	842	14			
1	B	606	Total	C	N	O	S	0	5	0
			4854	3099	881	860	14			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	33	GLN	HIS	SEE REMARK 999	UNP P35475
A	63	PRO	GLN	SEE REMARK 999	UNP P35475
A	105	GLN	ARG	SEE REMARK 999	UNP P35475
B	33	GLN	HIS	SEE REMARK 999	UNP P35475
B	63	PRO	GLN	SEE REMARK 999	UNP P35475
B	105	GLN	ARG	SEE REMARK 999	UNP P35475

- Molecule 2 is SUGAR (N-ACETYL-D-GLUCOSAMINE) (three-letter code: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	B	1	Total	C	N	O	0	0
			14	8	1	5		


- Molecule 3 is a polymer of unknown type called SUGAR (2-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	2	Total	C	N	O	0	0
			28	16	2	10		
3	B	2	Total	C	N	O	0	0
			28	16	2	10		

- Molecule 4 is a polymer of unknown type called SUGAR (5-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	5	Total	C	N	O	0	0
			61	34	2	25		

- Molecule 5 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			6	3	3		
5	A	1	Total	C	O	0	0
			6	3	3		

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			6	3	3		
5	B	1	Total	C	O	0	0
			6	3	3		
5	B	1	Total	C	O	0	0
			6	3	3		
5	B	1	Total	C	O	0	0
			6	3	3		
5	B	1	Total	C	O	0	0
			6	3	3		
5	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 6 is L(+)-TARTARIC ACID (three-letter code: TLA) (formula: C₄H₆O₆).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			10	4	6		
6	B	1	Total	C	O	0	0
			10	4	6		
6	B	1	Total	C	O	0	0
			10	4	6		

- Molecule 7 is SUGAR (L-IDURONIC ACID) (three-letter code: IDR) (formula: C₆H₁₀O₇).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	1	Total	C	O	0	0
			13	6	7		
7	B	1	Total	C	O	0	0
			13	6	7		

- Molecule 8 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	B	1	Total	Cl	0	0
			1	1		
8	A	1	Total	Cl	0	0
			1	1		

- Molecule 9 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
9	A	1	Total	O	S	0	0
			5	4	1		

- Molecule 10 is a polymer of unknown type called SUGAR (9-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
10	B	9	Total	C	N	O	0	0
			105	58	2	45		


- Molecule 11 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
11	A	37	Total	O	0	0
			37	37		
11	B	69	Total	O	0	0
			69	69		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Alpha-L-iduronidase



• Molecule 1: Alpha-L-iduronidase



4 Data and refinement statistics

Property	Value	Source
Space group	H 3	Depositor
Cell constants a, b, c, α , β , γ	258.94Å 258.94Å 71.53Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	48.94 – 2.46 48.94 – 2.46	Depositor EDS
% Data completeness (in resolution range)	97.8 (48.94-2.46) 97.8 (48.94-2.46)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.09	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.53 (at 2.45Å)	Xtriage
Refinement program	REFMAC 5.7.0029	Depositor
R, R_{free}	0.202 , 0.239 0.204 , 0.239	Depositor DCC
R_{free} test set	3175 reflections (5.26%)	DCC
Wilson B-factor (Å ²)	36.6	Xtriage
Anisotropy	0.697	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 27.5	EDS
Estimated twinning fraction	0.016 for h,-h-k,-l	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 63496 reflections	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	10073	wwPDB-VP
Average B, all atoms (Å ²)	40.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.09% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, BMA, NAG, CL, IDR, TLA, SO4, MAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.37	0/4891	0.58	0/6682
1	B	0.38	0/5002	0.59	0/6832
All	All	0.37	0/9893	0.59	0/13514

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4746	0	4640	12	0
1	B	4854	0	4754	13	0
2	A	14	0	13	0	0
2	B	14	0	13	0	0
3	A	28	0	25	0	0
3	B	28	0	25	0	0
4	A	61	0	52	0	0
5	A	18	0	24	0	0
5	B	36	0	48	6	0
6	A	10	0	4	1	0
6	B	20	0	8	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
7	A	13	0	9	0	0
7	B	13	0	9	0	0
8	A	1	0	0	0	0
8	B	1	0	0	0	0
9	A	5	0	0	0	0
10	B	105	0	88	0	0
11	A	37	0	0	0	0
11	B	69	0	0	0	0
All	All	10073	0	9712	25	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (25) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:450:PRO:O	1:A:531:ARG:NH1	2.33	0.61
1:A:127:LEU:HD22	1:A:171:HIS:CE1	2.43	0.54
1:B:75:ALA:HB2	5:B:917:GOL:H32	1.90	0.52
1:A:276:GLU:OE1	1:A:329:HIS:NE2	2.40	0.51
1:A:127:LEU:N	1:A:127:LEU:HD23	2.26	0.50
1:B:356:HIS:HB2	5:B:913:GOL:C3	2.43	0.49
1:A:401:LEU:HD21	1:A:424:ALA:HB2	1.95	0.48
1:B:530:LEU:HB3	1:B:534:SER:OG	2.13	0.48
1:B:314:ALA:HA	1:B:360:PHE:O	2.14	0.47
1:B:471:TYR:HA	1:B:539:HIS:O	2.15	0.47
1:B:555:ARG:NH2	5:B:917:GOL:O2	2.41	0.46
1:A:315:ASP:HB2	1:A:478:ASN:OD1	2.16	0.46
1:A:273:LEU:HD11	1:A:333:LEU:HD13	1.97	0.46
1:B:434:TRP:CH2	1:B:436:ALA:HB2	2.51	0.45
1:A:588:ASP:OD1	1:A:588:ASP:N	2.50	0.45
1:A:245:THR:HB	6:A:912:TLA:O4	2.17	0.45
1:B:127:LEU:CD2	1:B:171:HIS:CE1	3.01	0.44
1:A:383:ARG:HG2	1:A:387:LEU:HD23	2.01	0.43
1:B:604:LEU:C	1:B:604:LEU:HD23	2.40	0.42
1:B:356:HIS:HB2	5:B:913:GOL:H32	2.00	0.42
1:B:75:ALA:HB2	5:B:917:GOL:C3	2.50	0.42
1:A:434:TRP:CH2	1:A:436:ALA:HB2	2.56	0.41
1:B:224:SER:OG	5:B:918:GOL:H32	2.21	0.41
1:A:584:GLN:OE1	1:A:619:ARG:NH2	2.47	0.41
1:B:188:PHE:CE1	1:B:193:MET:HG2	2.55	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	589/627 (94%)	564 (96%)	25 (4%)	0	100	100
1	B	605/627 (96%)	584 (96%)	20 (3%)	1 (0%)	52	64
All	All	1194/1254 (95%)	1148 (96%)	45 (4%)	1 (0%)	56	71

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	247	PHE

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	505/527 (96%)	495 (98%)	10 (2%)	63	78
1	B	516/527 (98%)	507 (98%)	9 (2%)	68	81
All	All	1021/1054 (97%)	1002 (98%)	19 (2%)	65	79

All (19) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	41	TRP
1	A	69	ASP

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	239	ARG
1	A	245	THR
1	A	336	ASN
1	A	375	ARG
1	A	553	ARG
1	A	588	ASP
1	A	598	LYS
1	A	639	LEU
1	B	41	TRP
1	B	64	TYR
1	B	69	ASP
1	B	195	MET
1	B	337	THR
1	B	375	ARG
1	B	555	ARG
1	B	590	LYS
1	B	623	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	33	GLN
1	A	125	GLN
1	A	147	GLN
1	A	171	HIS
1	A	358	HIS
1	B	171	HIS
1	B	226	HIS
1	B	281	GLN
1	B	603	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates

18 carbohydrates are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	NAG	A	902	1,3	14,14,15	0.58	0	15,19,21	0.86	1 (6%)
3	NAG	A	903	3	14,14,15	0.55	0	15,19,21	0.84	0
4	NAG	A	904	1,4	14,14,15	0.61	0	15,19,21	1.02	1 (6%)
4	NAG	A	905	4	14,14,15	0.59	0	15,19,21	1.19	2 (13%)
4	BMA	A	906	4	11,11,12	0.38	0	14,15,17	0.53	0
4	MAN	A	907	4	11,11,12	0.51	0	14,15,17	1.08	1 (7%)
4	MAN	A	908	4	11,11,12	0.43	0	14,15,17	1.10	1 (7%)
3	NAG	B	902	1,3	14,14,15	0.49	0	15,19,21	1.31	2 (13%)
3	NAG	B	903	3	14,14,15	0.56	0	15,19,21	0.87	0
10	NAG	B	904	1,10	14,14,15	0.70	0	15,19,21	1.01	0
10	NAG	B	905	10	14,14,15	0.75	1 (7%)	15,19,21	0.87	0
10	BMA	B	906	10	11,11,12	0.57	0	14,15,17	0.99	1 (7%)
10	MAN	B	907	10	11,11,12	0.65	0	14,15,17	1.25	2 (14%)
10	MAN	B	908	10	11,11,12	0.48	0	14,15,17	1.13	1 (7%)
10	MAN	B	909	10	11,11,12	0.44	0	14,15,17	1.36	2 (14%)
10	MAN	B	910	10	11,11,12	0.53	0	14,15,17	0.81	0
10	MAN	B	911	10	11,11,12	0.48	0	14,15,17	1.31	2 (14%)
10	MAN	B	912	10	11,11,12	0.67	0	14,15,17	1.23	1 (7%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	A	902	1,3	-	0/6/23/26	0/1/1/1
3	NAG	A	903	3	-	0/6/23/26	0/1/1/1
4	NAG	A	904	1,4	-	0/6/23/26	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	A	905	4	-	0/6/23/26	0/1/1/1
4	BMA	A	906	4	-	0/2/19/22	0/1/1/1
4	MAN	A	907	4	-	0/2/19/22	0/1/1/1
4	MAN	A	908	4	-	0/2/19/22	0/1/1/1
3	NAG	B	902	1,3	-	0/6/23/26	0/1/1/1
3	NAG	B	903	3	-	0/6/23/26	0/1/1/1
10	NAG	B	904	1,10	-	0/6/23/26	0/1/1/1
10	NAG	B	905	10	-	0/6/23/26	0/1/1/1
10	BMA	B	906	10	-	0/2/19/22	0/1/1/1
10	MAN	B	907	10	-	0/2/19/22	0/1/1/1
10	MAN	B	908	10	-	0/2/19/22	0/1/1/1
10	MAN	B	909	10	-	0/2/19/22	0/1/1/1
10	MAN	B	910	10	-	0/2/19/22	0/1/1/1
10	MAN	B	911	10	-	0/2/19/22	0/1/1/1
10	MAN	B	912	10	-	0/2/19/22	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
10	B	905	NAG	O5-C1	-2.03	1.40	1.43

All (17) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
10	B	911	MAN	C1-C2-C3	-3.06	105.92	109.54
4	A	905	NAG	O4-C4-C3	-2.46	104.81	110.34
10	B	909	MAN	C1-C2-C3	-2.26	106.87	109.54
3	B	902	NAG	O7-C7-C8	-2.08	118.25	122.06
3	A	902	NAG	C1-O5-C5	2.03	114.83	112.25
10	B	906	BMA	C1-C2-C3	2.17	112.11	109.54
10	B	907	MAN	C1-C2-C3	2.33	112.30	109.54
4	A	905	NAG	C1-O5-C5	2.63	115.58	112.25
4	A	907	MAN	C1-O5-C5	2.81	115.81	112.25
10	B	911	MAN	O5-C1-C2	2.82	115.43	110.86
10	B	907	MAN	C1-O5-C5	2.83	115.83	112.25
4	A	904	NAG	C1-O5-C5	2.85	115.86	112.25
4	A	908	MAN	O5-C1-C2	3.19	116.04	110.86
10	B	912	MAN	C1-C2-C3	3.24	113.37	109.54
10	B	909	MAN	O5-C1-C2	3.61	116.70	110.86
10	B	908	MAN	C1-O5-C5	3.67	116.90	112.25
3	B	902	NAG	C1-O5-C5	3.89	117.19	112.25

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.6 Ligand geometry

Of 19 ligands modelled in this entry, 2 are monoatomic - leaving 17 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	NAG	A	901	1	14,14,15	0.54	0	15,19,21	1.39	1 (6%)
5	GOL	A	909	-	5,5,5	0.25	0	5,5,5	0.28	0
5	GOL	A	910	-	5,5,5	0.30	0	5,5,5	0.17	0
5	GOL	A	911	-	5,5,5	0.27	0	5,5,5	0.51	0
6	TLA	A	912	-	3,9,9	0.56	0	6,12,12	1.11	0
7	IDR	A	913	-	10,13,13	0.82	0	15,19,19	2.01	5 (33%)
9	SO4	A	915	-	4,4,4	0.43	0	6,6,6	0.06	0
2	NAG	B	901	1	14,14,15	0.45	0	15,19,21	0.79	0
5	GOL	B	913	-	5,5,5	0.30	0	5,5,5	0.37	0
5	GOL	B	914	-	5,5,5	0.21	0	5,5,5	0.29	0
5	GOL	B	915	-	5,5,5	0.32	0	5,5,5	0.22	0
5	GOL	B	916	-	5,5,5	0.36	0	5,5,5	0.40	0
5	GOL	B	917	-	5,5,5	0.27	0	5,5,5	0.47	0
5	GOL	B	918	-	5,5,5	0.25	0	5,5,5	0.39	0
6	TLA	B	919	-	3,9,9	0.35	0	6,12,12	1.31	1 (16%)
6	TLA	B	920	-	3,9,9	0.45	0	6,12,12	0.60	0
7	IDR	B	921	-	10,13,13	0.82	0	15,19,19	1.98	4 (26%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	A	901	1	-	0/6/23/26	0/1/1/1
5	GOL	A	909	-	-	0/4/4/4	0/0/0/0
5	GOL	A	910	-	-	0/4/4/4	0/0/0/0
5	GOL	A	911	-	-	0/4/4/4	0/0/0/0
6	TLA	A	912	-	-	0/4/12/12	0/0/0/0
7	IDR	A	913	-	-	0/0/24/24	0/1/1/1
9	SO4	A	915	-	-	0/0/0/0	0/0/0/0
2	NAG	B	901	1	-	0/6/23/26	0/1/1/1
5	GOL	B	913	-	-	0/4/4/4	0/0/0/0
5	GOL	B	914	-	-	0/4/4/4	0/0/0/0
5	GOL	B	915	-	-	0/4/4/4	0/0/0/0
5	GOL	B	916	-	-	0/4/4/4	0/0/0/0
5	GOL	B	917	-	-	0/4/4/4	0/0/0/0
5	GOL	B	918	-	-	0/4/4/4	0/0/0/0
6	TLA	B	919	-	-	0/4/12/12	0/0/0/0
6	TLA	B	920	-	-	0/4/12/12	0/0/0/0
7	IDR	B	921	-	-	0/0/24/24	0/1/1/1

There are no bond length outliers.

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	B	919	TLA	C1-C2-C3	-2.50	108.21	113.35
7	B	921	IDR	O1-C1-O5	-2.21	104.21	110.25
7	A	913	IDR	O5-C5-C4	2.31	112.72	108.76
7	A	913	IDR	O5-C1-C2	2.47	113.74	109.80
7	A	913	IDR	C4-C3-C2	2.51	115.48	110.79
7	B	921	IDR	C4-C3-C2	2.52	115.50	110.79
7	A	913	IDR	C1-O5-C5	2.87	116.44	112.22
7	B	921	IDR	C1-O5-C5	3.05	116.70	112.22
2	A	901	NAG	C1-O5-C5	3.88	117.17	112.25
7	B	921	IDR	C1-C2-C3	4.41	116.98	110.43
7	A	913	IDR	C1-C2-C3	4.51	117.13	110.43

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	A	912	TLA	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	B	913	GOL	2	0
5	B	917	GOL	3	0
5	B	918	GOL	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	595/627 (94%)	-0.02	19 (3%) 51 54	25, 39, 67, 92	0
1	B	606/627 (96%)	-0.19	9 (1%) 76 78	23, 33, 58, 87	0
All	All	1201/1254 (95%)	-0.11	28 (2%) 64 66	23, 35, 64, 92	0

All (28) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	589	GLY	7.6
1	A	338	THR	5.6
1	A	336	ASN	5.2
1	A	339	SER	4.2
1	B	588	ASP	3.7
1	A	64	TYR	3.4
1	B	590	LYS	3.3
1	A	585	PHE	3.3
1	B	641	VAL	3.2
1	A	594	PRO	3.2
1	A	639	LEU	3.2
1	A	190	ASN	2.9
1	A	627	ALA	2.9
1	A	337	THR	2.8
1	B	639	LEU	2.7
1	A	186	HIS	2.6
1	B	338	THR	2.6
1	A	189	ASP	2.5
1	A	635	PRO	2.5
1	A	611	THR	2.4
1	A	187	ASP	2.2
1	B	186	HIS	2.2
1	A	408	ALA	2.2
1	B	229	PRO	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	587	GLN	2.1
1	B	426	ARG	2.1
1	A	411	VAL	2.1
1	A	521	GLY	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	NAG	A	902	14/15	0.91	0.21	2.07	46,53,60,70	0
10	NAG	B	905	14/15	0.95	0.14	0.12	32,35,39,40	0
10	NAG	B	904	14/15	0.97	0.13	-0.19	33,34,36,38	0
10	BMA	B	906	11/12	0.94	0.12	-	42,47,53,55	0
4	MAN	A	908	11/12	0.86	0.29	-	76,82,87,87	0
10	MAN	B	912	11/12	0.85	0.23	-	71,74,77,77	0
4	NAG	A	905	14/15	0.95	0.13	-	47,50,53,57	0
3	NAG	B	903	14/15	0.72	0.51	-	79,84,91,95	0
10	MAN	B	907	11/12	0.87	0.14	-	63,70,78,79	0
10	MAN	B	910	11/12	0.72	0.26	-	84,93,95,95	0
10	MAN	B	908	11/12	0.53	0.35	-	80,84,87,90	0
3	NAG	B	902	14/15	0.92	0.22	-	47,53,61,70	0
3	NAG	A	903	14/15	0.82	0.37	-	78,82,88,88	0
4	BMA	A	906	11/12	0.95	0.15	-	57,64,71,72	0
10	MAN	B	909	11/12	0.61	0.23	-	71,85,89,93	0
4	MAN	A	907	11/12	0.89	0.27	-	68,73,75,76	0
4	NAG	A	904	14/15	0.94	0.15	-	44,46,49,52	0
10	MAN	B	911	11/12	0.93	0.12	-	53,56,59,65	0

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
7	IDR	A	913	13/13	0.89	0.23	5.58	49,54,58,59	0
7	IDR	B	921	13/13	0.91	0.17	4.22	42,53,58,60	0
5	GOL	A	909	6/6	0.84	0.29	4.01	52,55,56,58	0
5	GOL	B	913	6/6	0.87	0.17	3.54	44,44,44,45	0
5	GOL	A	910	6/6	0.89	0.21	3.41	63,64,65,66	0
5	GOL	A	911	6/6	0.86	0.26	2.25	47,49,51,53	0
6	TLA	A	912	10/10	0.87	0.17	2.15	55,58,59,62	0
5	GOL	B	914	6/6	0.92	0.16	0.84	40,41,42,44	0
5	GOL	B	916	6/6	0.86	0.15	0.61	46,49,50,52	0
5	GOL	B	915	6/6	0.90	0.11	-0.30	49,51,52,52	0
5	GOL	B	917	6/6	0.93	0.13	-0.46	36,36,37,37	0
6	TLA	B	920	10/10	0.89	0.16	-0.56	78,84,86,88	0
8	CL	A	914	1/1	0.98	0.11	-0.99	37,37,37,37	0
6	TLA	B	919	10/10	0.94	0.13	-1.28	37,40,46,46	0
8	CL	B	922	1/1	1.00	0.08	-1.78	32,32,32,32	0
9	SO4	A	915	5/5	0.90	0.15	-	82,82,85,88	0
5	GOL	B	918	6/6	0.79	0.31	-	67,69,72,76	0
2	NAG	B	901	14/15	0.94	0.19	-	44,49,52,55	0
2	NAG	A	901	14/15	0.86	0.22	-	71,75,76,77	0

6.5 Other polymers [i](#)

There are no such residues in this entry.