



# wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 07:19 PM GMT

PDB ID : 4OID  
Title : Structural and kinetic bases for the metal preference of the M18 aminopeptidase from *Pseudomonas aeruginosa*  
Authors : Nguyen, D.D.; Pandian, R.; Kim, D.D.; Ha, S.C.; Yoon, H.J.; Kim, K.S.; Yun, K.H.; Kim, J.H.; Kim, K.K.  
Deposited on : 2014-01-19  
Resolution : 2.30 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	3852 (2.30-2.30)
Clashscore	102246	4452 (2.30-2.30)
Ramachandran outliers	100387	4410 (2.30-2.30)
Sidechain outliers	100360	4409 (2.30-2.30)
RSRZ outliers	91569	3857 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	429	<div> <div>7%</div> <div>78% 14% • 5%</div> </div>
1	B	429	<div> <div>6%</div> <div>79% 13% •• 5%</div> </div>
1	C	429	<div> <div>7%</div> <div>78% 15% • 5%</div> </div>
1	D	429	<div> <div>5%</div> <div>82% 11% • 5%</div> </div>

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 13228 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Probable M18 family aminopeptidase 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	407	Total	C	N	O	S	0	0	0
			3151	1979	577	587	8			
1	B	407	Total	C	N	O	S	0	0	0
			3152	1978	578	588	8			
1	C	407	Total	C	N	O	S	0	0	0
			3153	1979	578	588	8			
1	D	407	Total	C	N	O	S	0	0	0
			3153	1979	578	588	8			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	236	ALA	ASP	ENGINEERED MUTATION	UNP Q9HYZ3
B	236	ALA	ASP	ENGINEERED MUTATION	UNP Q9HYZ3
C	236	ALA	ASP	ENGINEERED MUTATION	UNP Q9HYZ3
D	236	ALA	ASP	ENGINEERED MUTATION	UNP Q9HYZ3


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	146	Total	O	0	0
			146	146		
2	B	167	Total	O	0	0
			167	167		
2	C	153	Total	O	0	0
			153	153		
2	D	153	Total	O	0	0
			153	153		


### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Probable M18 family aminopeptidase 2




- Molecule 1: Probable M18 family aminopeptidase 2






- Molecule 1: Probable M18 family aminopeptidase 2





● Molecule 1: Probable M18 family aminopeptidase 2



## 4 Data and refinement statistics

Property	Value	Source
Space group	H 3	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	133.61Å 133.61Å 320.96Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	31.88 – 2.30 31.88 – 2.30	Depositor EDS
% Data completeness (in resolution range)	98.5 (31.88-2.30) 98.5 (31.88-2.30)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.23 (at 2.31Å)	Xtriage
Refinement program	REFMAC 5.6.0117	Depositor
R, $R_{free}$	0.157 , 0.207 0.162 , 0.210	Depositor DCC
$R_{free}$ test set	4678 reflections (5.28%)	DCC
Wilson B-factor (Å <sup>2</sup> )	27.4	Xtriage
Anisotropy	0.010	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.39 , 54.2	EDS
Estimated twinning fraction	0.012 for -h,1/3*h-1/3*k-1/3*l,-4/3*h-8/3*k+1/3*l 0.014 for -1/3*h+1/3*k+1/3*l,-k,8/3*h+4/3*k+1/3*l 0.019 for -2/3*h-1/3*k-1/3*l,-1/3*h-2/3*k+1/3*l,-4/3*h+4/3*k+1/3*l 0.008 for 1/3*h+2/3*k-1/3*l,-k,-8/3*h-4/3*k-1/3*l 0.018 for -1/3*h-2/3*k+1/3*l,-2/3*h-1/3*k-1/3*l,4/3*h-4/3*k-1/3*l 0.012 for -h,2/3*h+1/3*k+1/3*l,4/3*h+8/3*k-1/3*l 0.030 for h,-h-k,-l	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 93257 reflections	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	13228	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	31.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.35% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup> Intensities estimated from amplitudes.

<sup>2</sup> Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.96	0/3215	1.02	7/4361 (0.2%)
1	B	0.98	1/3217 (0.0%)	1.06	13/4366 (0.3%)
1	C	0.94	2/3218 (0.1%)	1.08	19/4368 (0.4%)
1	D	0.96	1/3218 (0.0%)	1.03	8/4368 (0.2%)
All	All	0.96	4/12868 (0.0%)	1.05	47/17463 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2
1	B	0	2
All	All	0	4

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	51	ARG	N-CA	6.54	1.59	1.46
1	C	401	HIS	N-CA	-5.31	1.35	1.46
1	D	305	SER	CB-OG	-5.12	1.35	1.42
1	B	107	GLU	CD-OE2	5.09	1.31	1.25

The worst 5 of 47 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	77	ARG	NE-CZ-NH2	-12.22	114.19	120.30
1	D	77	ARG	NE-CZ-NH2	-12.21	114.20	120.30
1	B	51	ARG	NE-CZ-NH2	-10.57	115.02	120.30
1	C	77	ARG	NE-CZ-NH1	9.79	125.19	120.30
1	C	68	ARG	NE-CZ-NH1	9.32	124.96	120.30

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	180	LEU	Peptide
1	A	425	SER	Peptide
1	B	183	GLY	Peptide
1	B	45	HIS	Peptide

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3151	0	3094	50	0
1	B	3152	0	3092	60	0
1	C	3153	0	3095	47	0
1	D	3153	0	3095	48	0
2	A	146	0	0	7	0
2	B	167	0	0	11	0
2	C	153	0	0	5	0
2	D	153	0	0	5	0
All	All	13228	0	12376	200	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

The worst 5 of 200 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:2:ARG:HB3	1:C:3:ALA:CB	1.65	1.26
1:C:2:ARG:CB	1:C:3:ALA:HB3	1.68	1.22
1:B:77:ARG:NH1	1:B:425:SER:O	1.82	1.13
1:B:2:ARG:HB3	1:B:3:ALA:HB3	1.11	1.09
1:D:368:THR:HG23	2:D:697:HOH:O	1.60	1.01

There are no symmetry-related clashes.



## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	399/429 (93%)	373 (94%)	21 (5%)	5 (1%)	15	15
1	B	401/429 (94%)	373 (93%)	22 (6%)	6 (2%)	13	12
1	C	401/429 (94%)	376 (94%)	22 (6%)	3 (1%)	26	31
1	D	401/429 (94%)	382 (95%)	18 (4%)	1 (0%)	52	64
All	All	1602/1716 (93%)	1504 (94%)	83 (5%)	15 (1%)	21	24

5 of 15 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	181	ALA
1	B	3	ALA
1	B	369	ARG
1	C	3	ALA
1	C	184	GLU

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	328/343 (96%)	313 (95%)	15 (5%)	33	44
1	B	329/343 (96%)	312 (95%)	17 (5%)	29	38
1	C	329/343 (96%)	315 (96%)	14 (4%)	35	47
1	D	329/343 (96%)	323 (98%)	6 (2%)	66	82
All	All	1315/1372 (96%)	1263 (96%)	52 (4%)	38	52

5 of 52 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	159	ARG
1	B	340	GLN
1	D	107	GLU
1	B	198	ARG
1	B	307	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 10 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	357	GLN
1	C	156	HIS
1	C	308	ASN
1	B	340	GLN
1	C	237	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

There are no ligands in this entry.

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	407/429 (94%)	-0.12	30 (7%) 17 25	14, 25, 66, 97	0
1	B	407/429 (94%)	-0.18	27 (6%) 22 29	13, 25, 65, 96	0
1	C	407/429 (94%)	-0.12	32 (7%) 15 22	13, 25, 62, 111	0
1	D	407/429 (94%)	-0.16	23 (5%) 27 36	13, 25, 61, 96	0
All	All	1628/1716 (94%)	-0.14	112 (6%) 20 27	13, 25, 64, 111	0

The worst 5 of 112 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	1	MET	8.8
1	B	372	MET	8.5
1	C	372	MET	7.9
1	D	372	MET	7.9
1	C	159	ARG	6.8

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.