



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 11:18 AM GMT

PDB ID : 3OO5
Title : R-state human hemoglobin: nitriheme modified
Authors : Yi, J.; Thomas, L.M.; Richter-Addo, G.B.
Deposited on : 2010-08-30
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	3939 (2.10-2.10)
Clashscore	102246	4460 (2.10-2.10)
Ramachandran outliers	100387	4413 (2.10-2.10)
Sidechain outliers	100360	4414 (2.10-2.10)
RSRZ outliers	91569	3948 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	141	
2	B	146	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	NTE	B	147[A]	-	-	-	X

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 2296 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Hemoglobin subunit alpha.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	140	Total	C	N	O	S	0	1	0
			1060	681	183	193	3			

- Molecule 2 is a protein called Hemoglobin subunit beta.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	143	Total	C	N	O	S	0	0	0
			1091	703	189	196	3			

- Molecule 3 is [3,3'-{(3S,4S)-13-ETHENYL-3,7,12,17-TETRAMETHYL-8-[(E)-2-NITROETHENYL]-3,4-DIHYDROPORPHYRIN-2,18-DIYL-KAPPA 4 N 21 ,N 22 ,N 23 ,N 24 }DIPROPANOATO(4-)]IRON (three-letter code: NTE) (formula: C₃₄H₃₁FeN₅O₆).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	Fe	N	O	0	0
			46	34	1	5	6		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	B	1	Total	C	Fe	N	O	0	1
			47	34	2	5	6		

- Molecule 4 is NITRITE ION (three-letter code: NO2) (formula: NO₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	N	O	0	0
			3	1	2		



- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	29	Total	O	0	0
			29	29		
5	B	20	Total	O	0	0
			20	20		



3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Hemoglobin subunit alpha



- Molecule 2: Hemoglobin subunit beta



4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, α , β , γ	53.50Å 53.50Å 192.22Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	27.49 – 2.10 27.49 – 2.10	Depositor EDS
% Data completeness (in resolution range)	100.0 (27.49-2.10) 100.0 (27.49-2.10)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.15 (at 2.10Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.6 _289)	Depositor
R, R_{free}	0.232 , 0.287 0.228 , 0.279	Depositor DCC
R_{free} test set	870 reflections (5.33%)	DCC
Wilson B-factor (Å ²)	48.2	Xtriage
Anisotropy	0.050	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 48.0	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.53$, $\langle L^2 \rangle = 0.37$	Xtriage
Outliers	1 of 17180 reflections (0.006%)	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	2296	wwPDB-VP
Average B, all atoms (Å ²)	53.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.77% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NTE, NO2

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.46	0/1091	0.54	0/1485
2	B	0.42	0/1119	0.49	0/1522
All	All	0.44	0/2210	0.52	0/3007

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1060	0	1065	21	0
2	B	1091	0	1089	22	0
3	A	46	0	29	1	0
3	B	47	0	29	11	0
4	A	3	0	0	0	0
5	A	29	0	0	0	0
5	B	20	0	0	0	0
All	All	2296	0	2212	45	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (45) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:B:147[A]:NTE:HBB	3:B:147[A]:NTE:HMB	1.55	0.87
1:A:31:ARG:HH11	2:B:127:GLN:HE21	1.30	0.79
3:B:147[A]:NTE:HHA	3:B:147[A]:NTE:HBAA	1.67	0.76
1:A:112:HIS:O	1:A:113:LEU:HD23	1.94	0.68
1:A:31:ARG:HD3	2:B:127:GLN:HE22	1.62	0.65
3:B:147[A]:NTE:HMC	3:B:147[A]:NTE:HBC	1.79	0.64
3:A:142:NTE:HMB	3:A:142:NTE:HBB	1.79	0.63
1:A:14:TRP:CZ3	1:A:67:THR:CG2	2.83	0.62
2:B:1:VAL:HG22	2:B:3:LEU:HD13	1.82	0.61
3:B:147[A]:NTE:O2D	3:B:147[A]:NTE:HBAA	2.01	0.60
2:B:124:PRO:HB2	2:B:125:PRO:HD3	1.83	0.59
3:B:147[A]:NTE:HBD	3:B:147[A]:NTE:HHA	1.84	0.59
1:A:14:TRP:CE3	1:A:67:THR:HG22	2.38	0.58
1:A:31:ARG:HD3	2:B:127:GLN:NE2	2.20	0.56
1:A:14:TRP:CZ3	1:A:67:THR:HG23	2.41	0.55
1:A:31:ARG:HH11	2:B:127:GLN:NE2	2.02	0.53
1:A:14:TRP:CZ3	1:A:67:THR:HG22	2.42	0.53
2:B:3:LEU:HG	2:B:7:GLU:HB3	1.89	0.53
1:A:86:LEU:HG	1:A:91:LEU:HD12	1.90	0.53
3:B:147[A]:NTE:HBD	3:B:147[A]:NTE:CHA	2.38	0.53
2:B:75:LEU:HA	2:B:78:LEU:HD21	1.90	0.52
2:B:41:PHE:CE2	3:B:147[A]:NTE:HBCA	2.48	0.49
1:A:86:LEU:HG	1:A:91:LEU:CD1	2.42	0.49
1:A:84:SER:HB3	1:A:136:LEU:HA	1.93	0.48
1:A:113:LEU:HB3	1:A:116:GLU:HB3	1.96	0.48
2:B:41:PHE:CD2	3:B:147[A]:NTE:HBCA	2.49	0.47
1:A:3:SER:O	1:A:7:LYS:HG3	2.14	0.46
1:A:14:TRP:HZ3	1:A:67:THR:HG23	1.81	0.46
1:A:35:SER:HB3	2:B:131:GLN:HG3	1.98	0.45
2:B:53:ALA:O	2:B:57:ASN:HB2	2.16	0.45
1:A:20:HIS:HB3	1:A:24:TYR:CE2	2.53	0.44
1:A:33:PHE:CE2	1:A:48:LEU:HD22	2.53	0.44
2:B:63:HIS:NE2	3:B:147[A]:NTE:CHA	2.81	0.43
2:B:100:PRO:HA	2:B:103:PHE:CD2	2.53	0.43
2:B:50:THR:HB	2:B:51:PRO:HD2	2.01	0.43
2:B:1:VAL:O	2:B:1:VAL:HG13	2.18	0.43
3:B:147[A]:NTE:HHA	3:B:147[A]:NTE:CBD	2.47	0.42
1:A:93:VAL:O	1:A:140:TYR:OH	2.37	0.42
1:A:14:TRP:HZ3	1:A:67:THR:CG2	2.31	0.42
2:B:45:PHE:HA	2:B:59:LYS:HD3	2.02	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:50:THR:O	2:B:54:VAL:HG23	2.20	0.42
2:B:90:GLU:O	2:B:90:GLU:HG2	2.20	0.41
1:A:119:PRO:HG2	2:B:55:MET:HG3	2.04	0.40
2:B:63:HIS:CE1	3:B:147[A]:NTE:CHA	3.04	0.40
2:B:3:LEU:HD22	2:B:78:LEU:HD13	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	139/141 (99%)	137 (99%)	2 (1%)	0	100	100
2	B	141/146 (97%)	134 (95%)	7 (5%)	0	100	100
All	All	280/287 (98%)	271 (97%)	9 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	113/113 (100%)	111 (98%)	2 (2%)	66	72
2	B	115/118 (98%)	114 (99%)	1 (1%)	84	89
All	All	228/231 (99%)	225 (99%)	3 (1%)	76	82

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	8	THR
1	A	92	ARG
2	B	20	VAL

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
2	B	127	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 4 ligands modelled in this entry, 1 is modelled with single atom - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	NTE	A	142	1,4	25,53,53	1.44	5 (20%)	19,86,86	1.91	4 (21%)
4	NO2	A	143	3	2,2,2	0.04	0	1,1,1	0.58	0
3	NTE	B	147[A]	2	25,53,53	1.42	4 (16%)	19,86,86	1.96	7 (36%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NTE	A	142	1,4	-	0/9/57/57	0/0/8/8
4	NO2	A	143	3	-	0/0/0/0	0/0/0/0
3	NTE	B	147[A]	2	-	0/9/57/57	0/0/8/8

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	142	NTE	C4D-ND	-2.34	1.33	1.36
3	B	147[A]	NTE	C4A-CHB	2.75	1.47	1.39
3	A	142	NTE	C4B-CHC	2.78	1.47	1.39
3	A	142	NTE	C1A-CHA	2.93	1.47	1.39
3	A	142	NTE	C4A-CHB	2.94	1.47	1.39
3	B	147[A]	NTE	C4B-CHC	2.98	1.48	1.39
3	A	142	NTE	C1D-CHD	2.98	1.48	1.39
3	B	147[A]	NTE	C1A-CHA	3.06	1.48	1.39
3	B	147[A]	NTE	C1D-CHD	3.25	1.48	1.39

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	142	NTE	CBD-CAD-C3D	-5.16	103.28	112.53
3	A	142	NTE	C3C-CAC-CBC	-3.18	119.81	126.32
3	B	147[A]	NTE	C3B-CAB-CBB	-2.73	119.87	125.24
3	B	147[A]	NTE	C3C-CAC-CBC	-2.62	120.95	126.32
3	B	147[A]	NTE	CAA-CBA-CGA	-2.61	107.95	112.75
3	A	142	NTE	CAA-CBA-CGA	-2.49	108.18	112.75
3	B	147[A]	NTE	CAA-C2A-C3A	-2.10	123.00	129.00
3	B	147[A]	NTE	CMB-C2B-C3B	2.11	129.22	125.09
3	B	147[A]	NTE	CMC-C2C-C3C	2.52	130.02	125.09
3	A	142	NTE	CMC-C2C-C3C	2.57	130.12	125.09
3	B	147[A]	NTE	CAA-C2A-C1A	4.76	132.18	127.01

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 12 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	142	NTE	1	0
3	B	147[A]	NTE	11	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	140/141 (99%)	0.31	8 (5%)	27 35	29, 44, 65, 104	2 (1%)
2	B	143/146 (97%)	1.03	26 (18%)	2 2	33, 57, 92, 116	4 (2%)
All	All	283/287 (98%)	0.67	34 (12%)	6 8	29, 49, 88, 116	6 (2%)

All (34) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	89	SER	7.1
2	B	93	CYS	6.7
2	B	90	GLU	6.0
2	B	86	ALA	5.9
2	B	1	VAL	5.7
2	B	94	ASP	5.6
2	B	91	LEU	5.3
1	A	140	TYR	4.2
2	B	60	VAL	4.2
2	B	82	LYS	4.0
2	B	45	PHE	3.9
2	B	143	HIS	3.8
2	B	97	HIS	3.5
2	B	58	PRO	3.1
2	B	106	LEU	3.0
1	A	101	LEU	2.8
2	B	56	GLY	2.8
2	B	43	GLU	2.8
1	A	14	TRP	2.8
2	B	111	VAL	2.7
1	A	1	VAL	2.6
2	B	59	LYS	2.6
1	A	105	LEU	2.5
2	B	95	LYS	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	106	LEU	2.5
1	A	75	ASP	2.3
2	B	107	GLY	2.2
2	B	109	VAL	2.2
2	B	44	SER	2.2
1	A	16	LYS	2.2
2	B	61	LYS	2.2
2	B	47	ASP	2.1
2	B	96	LEU	2.1
2	B	112	CYS	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	NTE	B	147[A]	46/46	0.88	0.30	2.23	45,55,59,62	46
3	NTE	A	142	46/46	0.95	0.11	-0.43	29,37,59,67	3
4	NO2	A	143	3/3	0.98	0.07	-1.26	35,35,38,38	0
3	NTE	B	147[B]	1/46	0.88	0.30	-	61,61,61,61	1

6.5 Other polymers [i](#)

There are no such residues in this entry.