



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 07:22 PM GMT

PDB ID : 4OO9  
Title : Structure of the human class C GPCR metabotropic glutamate receptor 5 transmembrane domain in complex with the negative allosteric modulator mavoglurant  
Authors : Dore, A.S.; Okrasa, K.; Patel, J.C.; Serrano-Vega, M.; Bennett, K.; Cooke, R.M.; Errey, J.C.; Jazayeri, A.; Khan, S.; Tehan, B.; Weir, M.; Wiggin, G.R.; Marshall, F.H.  
Deposited on : 2014-01-31  
Resolution : 2.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*


The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	2328 (2.60-2.60)
Clashscore	102246	2679 (2.60-2.60)
Ramachandran outliers	100387	2635 (2.60-2.60)
Sidechain outliers	100360	2635 (2.60-2.60)
RSRZ outliers	91569	2334 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	444	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	OLA	A	4001	-	-	-	X

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	OLA	A	4002	-	-	-	X

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 3355 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Metabotropic glutamate receptor 5, Lysozyme, Metabotropic glutamate receptor 5 chimera.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	409	Total	C	N	O	S	0	0	0
			3212	2096	534	558	24			

There are 24 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	566	ALA	-	EXPRESSION TAG	UNP P41594
A	567	ALA	-	EXPRESSION TAG	UNP P41594
A	568	SER	-	EXPRESSION TAG	UNP P41594
A	579	ALA	GLU	ENGINEERED MUTATION	UNP P41594
A	667	TYR	ASN	ENGINEERED MUTATION	UNP P41594
A	669	ALA	ILE	ENGINEERED MUTATION	UNP P41594
A	675	MET	GLY	ENGINEERED MUTATION	UNP P41594
A	1054	THR	CYS	ENGINEERED MUTATION	UNP P00720
A	1097	ALA	CYS	ENGINEERED MUTATION	UNP P00720
A	742	ALA	THR	ENGINEERED MUTATION	UNP P41594
A	753	ALA	SER	ENGINEERED MUTATION	UNP P41594
A	837	ALA	-	EXPRESSION TAG	UNP P41594
A	838	ALA	-	EXPRESSION TAG	UNP P41594
A	839	ALA	-	EXPRESSION TAG	UNP P41594
A	840	HIS	-	EXPRESSION TAG	UNP P41594
A	841	HIS	-	EXPRESSION TAG	UNP P41594
A	842	HIS	-	EXPRESSION TAG	UNP P41594
A	843	HIS	-	EXPRESSION TAG	UNP P41594
A	844	HIS	-	EXPRESSION TAG	UNP P41594
A	845	HIS	-	EXPRESSION TAG	UNP P41594
A	846	HIS	-	EXPRESSION TAG	UNP P41594
A	847	HIS	-	EXPRESSION TAG	UNP P41594
A	848	HIS	-	EXPRESSION TAG	UNP P41594
A	849	HIS	-	EXPRESSION TAG	UNP P41594

- Molecule 2 is OLEIC ACID (three-letter code: OLA) (formula: C<sub>18</sub>H<sub>34</sub>O<sub>2</sub>).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			20	18	2		
2	A	1	Total	C	O	0	0
			15	13	2		
2	A	1	Total	C	O	0	0
			14	12	2		
2	A	1	Total	C	O	0	0
			13	11	2		

- Molecule 3 is MAVOGLURANT (three-letter code: 2U8) (formula: C<sub>19</sub>H<sub>23</sub>NO<sub>3</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			23	19	1	3		

- Molecule 4 is 2-(N-MORPHOLINO)-ETHANESULFONIC ACID (three-letter code: MES) (formula: C<sub>6</sub>H<sub>13</sub>NO<sub>4</sub>S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O	S	0	0
			12	6	1	4	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	46	Total	O	0	0
			46	46		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Metabotropic glutamate receptor 5, Lysozyme, Metabotropic glutamate receptor 5 chimera



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	143.24Å 43.55Å 82.05Å 90.00° 99.37° 90.00°	Depositor
Resolution (Å)	29.74 – 2.60 34.51 – 2.60	Depositor EDS
% Data completeness (in resolution range)	94.1 (29.74-2.60) 94.1 (34.51-2.60)	Depositor EDS
$R_{merge}$	0.12	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.05 (at 2.61Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.8.1_1168)	Depositor
R, $R_{free}$	0.239 , 0.275 0.244 , 0.280	Depositor DCC
$R_{free}$ test set	691 reflections (4.90%)	DCC
Wilson B-factor (Å <sup>2</sup> )	41.4	Xtriage
Anisotropy	0.910	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 47.1	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 14787 reflections	Xtriage
$F_o, F_c$ correlation	0.92	EDS
Total number of atoms	3355	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	40.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.88% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.



## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: OLA, 2U8, MES, YCM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.22	0/3255	0.38	0/4413

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3212	0	3342	24	0
2	A	62	0	87	3	0
3	A	23	0	23	1	0
4	A	12	0	13	0	0
5	A	46	0	0	3	0
All	All	3355	0	3465	24	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (24) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:607:THR:HB	1:A:832:ASN:HD21	1.51	0.74
1:A:625:ILE:HG23	1:A:810:ALA:HB1	1.77	0.66
1:A:1011:GLU:O	5:A:4142:HOH:O	2.16	0.62
1:A:1053:ASN:OD1	1:A:1055:ASN:ND2	2.32	0.61
1:A:1002:ASN:ND2	5:A:4112:HOH:O	2.35	0.59
1:A:709:VAL:HG11	2:A:4003:OLA:H81	1.86	0.57
1:A:1123:GLN:OE1	1:A:1125:ARG:NH2	2.35	0.57
1:A:653:LEU:HD11	1:A:705:LEU:HD13	1.88	0.55
1:A:633:PHE:HE1	2:A:4004:OLA:H62	1.72	0.54
1:A:613:SER:OG	1:A:665:LYS:NZ	2.42	0.51
1:A:639:PRO:HA	1:A:731:LEU:HD13	1.92	0.50
1:A:741:VAL:HG13	2:A:4001:OLA:H72	1.93	0.50
1:A:635:LEU:HD21	1:A:651:ILE:HD11	1.99	0.45
1:A:638:LYS:NZ	5:A:4109:HOH:O	2.51	0.43
1:A:759:PHE:O	1:A:762:ARG:NH2	2.44	0.42
1:A:747:ASN:ND2	3:A:4005:2U8:O2	2.45	0.42
1:A:1050:ILE:HD12	1:A:1062:GLU:HB3	2.01	0.41
1:A:612:SER:OG	1:A:767:ASN:ND2	2.53	0.41
1:A:654:SER:OG	1:A:655:PRO:HD3	2.20	0.41
1:A:711:LEU:HD13	1:A:739:GLY:HA2	2.02	0.41
1:A:764:VAL:HA	1:A:765:PRO:HD3	1.87	0.41
1:A:573:LEU:HD13	1:A:581:ILE:HG21	2.03	0.41
1:A:635:LEU:HB3	1:A:799:ILE:HG23	2.02	0.40
1:A:718:ASP:OD1	1:A:719:ILE:N	2.54	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	401/444 (90%)	390 (97%)	11 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	342/371 (92%)	340 (99%)	2 (1%)	90	97

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	586	PHE
1	A	1008	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	832	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
1	YCM	A	634	1	8,9,10	1.26	1 (12%)	5,10,12	1.62	1 (20%)
1	YCM	A	691	1	8,9,10	1.22	1 (12%)	5,10,12	1.61	1 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	YCM	A	634	1	-	0/6/8/10	0/0/0/0
1	YCM	A	691	1	-	0/6/8/10	0/0/0/0

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	634	YCM	CD-SG	-2.18	1.76	1.81
1	A	691	YCM	CD-SG	-2.18	1.76	1.81

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	634	YCM	CD-CE-NZ2	3.00	118.78	115.48
1	A	691	YCM	CD-CE-NZ2	3.07	118.86	115.48

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.5 Carbohydrates

There are no carbohydrates in this entry.

## 5.6 Ligand geometry

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	OLA	A	4001	-	16,19,19	0.12	0	16,19,19	0.65	0
2	OLA	A	4002	-	11,14,19	0.13	0	10,14,19	0.57	0
2	OLA	A	4003	-	10,13,19	0.23	0	10,13,19	0.64	0
2	OLA	A	4004	-	9,12,19	0.19	0	9,12,19	0.70	0
3	2U8	A	4005	-	22,25,25	1.36	2 (9%)	33,36,36	1.95	8 (24%)
4	MES	A	4006	-	11,12,12	0.46	0	14,16,16	2.21	2 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	OLA	A	4001	-	-	0/15/17/17	0/0/0/0
2	OLA	A	4002	-	-	0/10/12/17	0/0/0/0
2	OLA	A	4003	-	-	0/9/11/17	0/0/0/0
2	OLA	A	4004	-	-	0/8/10/17	0/0/0/0
3	2U8	A	4005	-	-	0/10/38/38	0/3/3/3
4	MES	A	4006	-	-	0/6/14/14	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	4005	2U8	C4-C9	2.90	1.23	1.19
3	A	4005	2U8	C18-N1	3.50	1.40	1.35

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	4005	2U8	C15-C10-C9	-4.90	105.91	110.90
4	A	4006	MES	O1-C6-C5	-2.06	107.11	111.84
3	A	4005	2U8	C11-C12-C13	2.02	114.75	111.39
3	A	4005	2U8	C13-C14-C15	3.13	117.25	113.40
3	A	4005	2U8	C12-C11-C10	3.19	114.25	111.92
3	A	4005	2U8	C19-O3-C18	3.29	119.50	115.40
3	A	4005	2U8	C12-C13-C14	3.44	118.72	110.38
3	A	4005	2U8	C16-C17-N1	3.62	107.23	103.34
3	A	4005	2U8	C10-C15-C14	4.62	119.93	111.73
4	A	4006	MES	O2S-S-C8	6.84	112.74	106.91

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	4001	OLA	1	0
2	A	4003	OLA	1	0
2	A	4004	OLA	1	0
3	A	4005	2U8	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	407/444 (91%)	0.11	10 (2%) 61 54	19, 38, 66, 76	0

All (10) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	612	SER	5.0
1	A	831	ARG	4.9
1	A	638	LYS	3.8
1	A	1041	ALA	3.0
1	A	1076	ARG	2.6
1	A	1036	SER	2.3
1	A	660	SER	2.3
1	A	1037	PRO	2.2
1	A	640	LYS	2.1
1	A	1035	LYS	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
1	YCM	A	691	10/11	0.87	0.27	-	49,53,80,114	0
1	YCM	A	634	10/11	0.87	0.19	-	33,36,61,73	0

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
2	OLA	A	4002	15/20	0.75	0.21	2.50	55,57,68,68	0
2	OLA	A	4001	20/20	0.88	0.23	2.24	39,49,62,62	0
2	OLA	A	4003	14/20	0.82	0.23	0.75	61,64,68,68	0
2	OLA	A	4004	13/20	0.82	0.27	0.49	68,69,70,70	0
3	2U8	A	4005	23/23	0.95	0.20	0.36	23,31,40,42	0
4	MES	A	4006	12/12	0.98	0.17	-0.06	38,50,51,51	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.