



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 11:18 AM GMT

PDB ID : 3OPE
Title : Structural Basis of Auto-inhibitory mechanism of Histone methyltransferase
Authors : An, S.; Song, J.
Deposited on : 2010-08-31
Resolution : 2.90 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1451 (2.90-2.90)
Clashscore	102246	1668 (2.90-2.90)
Ramachandran outliers	100387	1630 (2.90-2.90)
Sidechain outliers	100360	1632 (2.90-2.90)
RSRZ outliers	91569	1456 (2.90-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	222	<div> <div>5%</div> <div>47%</div> <div>42%</div> <div>7%</div> <div>.</div> </div>
1	B	222	<div> <div>3%</div> <div>33%</div> <div>46%</div> <div>6%</div> <div>.</div> <div>13%</div> </div>

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 3391 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Probable histone-lysine N-methyltransferase ASH1L.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	213	Total	C	N	O	S	0	0	0
			1726	1070	308	326	22			
1	B	193	Total	C	N	O	S	0	0	0
			1578	984	281	293	20			


There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	2067	GLY	-	EXPRESSION TAG	UNP Q9NR48
A	2068	SER	-	EXPRESSION TAG	UNP Q9NR48
B	2067	GLY	-	EXPRESSION TAG	UNP Q9NR48
B	2068	SER	-	EXPRESSION TAG	UNP Q9NR48

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	3	Total	Zn	0	0
			3	3		
2	A	3	Total	Zn	0	0
			3	3		

- Molecule 3 is S-ADENOSYLMETHIONINE (three-letter code: SAM) (formula: C₁₅H₂₂N₆O₅S).



- | Mol | Chain | Residues | Atoms | ZeroOcc | AltConf |
|-----|-------|----------|------------------|---------|---------|
| 4 | A | 12 | Total O
12 12 | 0 | 0 |
| 4 | B | 15 | Total O
15 15 | 0 | 0 |

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Chain A:
-
- 5% 47% 42% 7%
- | Residue | Category |
|---------|----------|
| D2221 | Green |
| P2222 | Green |
| M2226 | Green |
| Q2227 | Green |
| K2228 | Green |
| W2229 | Green |
| Y2235 | Green |
| R2236 | Green |
| L2239 | Green |
| D2244 | Green |
| M2245 | Green |
| P2246 | Green |
| T2249 | Green |
| Y2253 | Green |
| D2254 | Green |
| Y2255 | Green |
| N2256 | Green |
| F2257 | Green |
| H2258 | Green |
| S2259 | Green |
| F2260 | Green |
| N2261 | Green |
| V2262 | Green |
| E2263 | Green |
| K2264 | Green |
| Q2265 | Green |
| Q2266 | Green |
| K2269 | Green |
| I2279 | Green |
| G2280 | Green |
| LVS | Grey |
| SER | Grey |
| GLN | Grey |
| ARG | Grey |
| VAL | Grey |
| ASN | Grey |
| GLY | Grey |
- | Residue | Category |
|---------|----------|
| L2141 | Green |
| L2142 | Green |
| F2145 | Green |
| R2146 | Green |
| A2147 | Green |
| E2148 | Green |
| L2149 | Green |
| K2150 | Green |
| T2154 | Green |
| L2155 | Green |
| T2156 | Green |
| L2160 | Green |
| K2161 | Green |
| F2165 | Green |
| E2168 | Green |
| V2169 | Green |
| L2170 | Green |
| V2173 | Green |
| E2178 | Green |
| F2179 | Green |
| R2180 | Green |
| H2181 | Green |
| R2182 | Green |
| Q2186 | Green |
| V2187 | Green |
| L2188 | Green |
| H2189 | Green |
| H2190 | Green |
| S2191 | Green |
| D2192 | Green |
| H2193 | Green |
| Y2194 | Green |
| C2195 | Green |
| L2196 | Green |
| N2197 | Green |
| L2198 | Green |
| G2201 | Green |
| N2202 | Green |
| T2203 | Green |
| T2204 | Green |
| R2208 | Green |
| E2212 | Green |
| A2213 | Green |
| R2214 | Green |
| H2218 | Green |
| S2219 | Green |
| C2220 | Green |
- | Residue | Category |
|---------|----------|
| S2068 | Green |
| T2072 | Green |
| S2073 | Green |
| S2074 | Green |
| T2075 | Green |
| V2076 | Green |
| T2077 | Green |
| V2078 | Green |
| T2079 | Green |
| V2080 | Green |
| R2081 | Green |
| F2082 | Green |
| T2089 | Red |
| T2090 | Green |
| C2091 | Green |
| R2092 | Red |
| C2093 | Green |
| R2094 | Red |
| K2095 | Green |
| P2096 | Red |
| R2097 | Red |
| T2100 | Red |
| V2102 | Red |
| C2103 | Red |
| C2104 | Red |
| V2105 | Red |
| D2107 | Green |
| C2108 | Orange |
| L2109 | Orange |
| R2110 | Orange |
| R2111 | Orange |
| F2114 | Green |
| A2115 | Green |
| E2116 | Green |
| C2117 | Green |
| N2120 | Red |
| C2121 | Red |
| C2122 | Green |
| F2123 | Green |
| C2124 | Green |
| G2125 | Orange |
| C2126 | Green |
| Q2127 | Green |
| C2128 | Orange |
| C2129 | Orange |
| N2130 | Green |
| Q2131 | Green |
| T2132 | Green |
| Q2133 | Green |
| R2135 | Green |
| H2136 | Green |

- Chain B:
-
- | Amino Acid | Percentage |
|------------|------------|
| GLY | 3% |
| S2068 | 33% |
| Y2069 | 46% |
| K2070 | 46% |
| R2073 | 46% |
| S2074 | 46% |
| N2075 | 46% |
| V2078 | 46% |
| D2079 | 46% |
| V2080 | 46% |
| R2081 | 46% |
| P2082 | 46% |
| L2083 | 46% |
| S2084 | 46% |
| GLY | 46% |
| TTR | 46% |
| GLU | 46% |
| ALA | 46% |
| THR | 46% |
| THR | 46% |
| THR | 46% |
| CYS | 46% |
| ASN | 46% |
| CYS | 46% |
| LYS | 46% |
| LVS | 46% |
| PRO | 46% |
| ASP | 46% |
| ASP | 46% |
| ASP | 46% |
| THR | 46% |
| ARG | 46% |
| LVS | 46% |
| GLY | 46% |
| C2104 | 46% |
| V2105 | 46% |
| D2106 | 46% |
| D2107 | 46% |
| C2108 | 46% |
| L2109 | 46% |
| N2110 | 46% |
| L2113 | 46% |
| F2114 | 46% |
| S2118 | 46% |
| P2119 | 46% |
| N2120 | 46% |
| T2121 | 46% |
| C2122 | 46% |
| P2123 | 46% |
| C2124 | 46% |
| G2125 | 46% |
| C2128 | 46% |
| C2129 | 46% |
| N2130 | 46% |
| Q2131 | 46% |
| B2132 | 46% |
| R2135 | 3% |
| H2136 | 3% |
| E2137 | 3% |
| V2138 | 3% |
| Q2140 | 3% |
| C2141 | 3% |
| L2142 | 3% |
| E2143 | 3% |
| R2144 | 3% |
| F2145 | 3% |
| R2146 | 3% |
| A2147 | 3% |
| K2150 | 3% |
| G2151 | 3% |
| N2152 | 3% |
| R2155 | 3% |
| T2156 | 3% |
| K2157 | 3% |
| L2160 | 3% |
| K2161 | 3% |
| A2162 | 3% |
| G2163 | 3% |
| Q2164 | 3% |
| F2165 | 3% |
| E2168 | 3% |
| Y2169 | 3% |
| L2170 | 3% |
| Q2177 | 3% |
| R2180 | 3% |
| M2183 | 3% |
| Y2187 | 3% |
| H2188 | 3% |
| N2189 | 3% |
| H2193 | 3% |
| Y2194 | 3% |
| C2195 | 3% |
| L2196 | 3% |
| N2197 | 3% |
| L2198 | 3% |
| D2199 | 3% |
| N2202 | 3% |
| V2203 | 3% |
| I2204 | 3% |
| D2205 | 3% |
| S2206 | 3% |
| Y2207 | 3% |
| R2208 | 3% |
| N2209 | 3% |
| G2210 | 3% |
| N2211 | 3% |
| E2212 | 3% |
| A2213 | 3% |
| R2214 | 3% |
| F2215 | 3% |
| I2216 | 3% |
| N2217 | 3% |
| H2218 | 3% |
| S2219 | 3% |
| R2220 | 3% |
| D2221 | 3% |
| P2222 | 3% |
| N2223 | 3% |
| C2224 | 3% |
| E2225 | 3% |
| N2226 | 3% |
| W2229 | 3% |
| V2234 | 3% |
| Y2235 | 3% |
| R2236 | 3% |
| I2237 | 3% |
| G2238 | 3% |
| L2239 | 3% |
| K2243 | 3% |
| D2244 | 3% |
| M2245 | 3% |
| F2246 | 3% |
| A2247 | 3% |
| G2248 | 3% |
| T2249 | 3% |
| E2250 | 3% |
| L2251 | 3% |
| T2252 | 3% |
| Y2253 | 3% |
| D2254 | 3% |
| Y2255 | 3% |
| H2258 | 3% |
| S2259 | 3% |
| F2260 | 3% |
| N2261 | 3% |
| V2262 | 3% |
| E2263 | 3% |
| K2264 | 3% |
| Q2265 | 3% |
| L2266 | 3% |
| T2267 | 3% |
| C2268 | 3% |
| K2269 | 3% |
| G2270 | 3% |
| G2271 | 3% |
| F2272 | 3% |
| E2273 | 3% |
| K2274 | 3% |
| Q2275 | 3% |
| R2276 | 3% |

4 Data and refinement statistics

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, α , β , γ	59.53Å 59.53Å 233.80Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	50.00 – 2.90 38.66 – 2.91	Depositor EDS
% Data completeness (in resolution range)	(Not available) (50.00-2.90) 98.0 (38.66-2.91)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	0.10	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.81 (at 2.90Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.235 , 0.299 0.247 , 0.319	Depositor DCC
R_{free} test set	542 reflections (4.90%)	DCC
Wilson B-factor (Å ²)	67.1	Xtriage
Anisotropy	0.131	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 50.8	EDS
Estimated twinning fraction	0.036 for -h,-k,l	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Outliers	0 of 20348 reflections	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	3391	wwPDB-VP
Average B, all atoms (Å ²)	67.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.28% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, SAM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.44	1/1763 (0.1%)	0.66	0/2371
1	B	0.42	0/1612	0.68	1/2167 (0.0%)
All	All	0.43	1/3375 (0.0%)	0.67	1/4538 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	2195	CYS	CB-SG	-5.00	1.73	1.81

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	2275	CYS	CA-CB-SG	5.84	124.51	114.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1726	0	1635	89	0
1	B	1578	0	1500	119	0
2	A	3	0	0	0	0
2	B	3	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	27	0	22	3	0
3	B	27	0	22	6	0
4	A	12	0	0	0	0
4	B	15	0	0	3	0
All	All	3391	0	3179	207	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 32.

The worst 5 of 207 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:2245:MET:HB3	1:A:2249:THR:HG21	1.30	1.08
1:B:2220:CYS:CB	1:B:2275:CYS:HB2	1.88	1.03
1:B:2218:HIS:HD2	1:B:2255:TYR:H	1.17	0.90
1:B:2267:LEU:HB2	1:B:2269:LYS:HE3	1.55	0.89
1:B:2214:ARG:O	3:B:8:SAM:HE1	1.77	0.84

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	211/222 (95%)	159 (75%)	38 (18%)	14 (7%)	1	4
1	B	189/222 (85%)	139 (74%)	38 (20%)	12 (6%)	2	5
All	All	400/444 (90%)	298 (74%)	76 (19%)	26 (6%)	1	4

5 of 26 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	2096	PRO
1	A	2105	VAL
1	A	2109	LEU
1	A	2148	GLU
1	B	2136	HIS

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	191/197 (97%)	181 (95%)	10 (5%)	29	64
1	B	175/197 (89%)	165 (94%)	10 (6%)	25	59
All	All	366/394 (93%)	346 (94%)	20 (6%)	27	61

5 of 20 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	2257	PHE
1	B	2107	ASP
1	B	2196	LEU
1	A	2190	HIS
1	A	2244	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 13 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	2266	GLN
1	B	2075	ASN
1	B	2197	ASN
1	A	2218	HIS
1	B	2130	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 6 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	SAM	A	7	-	21,29,29	1.20	2 (9%)	17,42,42	3.14	6 (35%)
3	SAM	B	8	-	21,29,29	1.18	1 (4%)	17,42,42	3.15	6 (35%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	SAM	A	7	-	-	0/8/33/33	0/3/3/3
3	SAM	B	8	-	-	0/8/33/33	0/3/3/3

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	7	SAM	O4'-C4'	-2.05	1.40	1.45
3	B	8	SAM	CG-CB	3.24	1.56	1.52
3	A	7	SAM	CG-CB	3.33	1.56	1.52

The worst 5 of 12 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
3	B	8	SAM	N3-C2-N1	-10.77	120.64	128.89
3	A	7	SAM	N3-C2-N1	-10.73	120.68	128.89
3	A	7	SAM	C4-C5-N7	-3.64	106.13	109.48
3	B	8	SAM	C4-C5-N7	-3.62	106.15	109.48
3	A	7	SAM	N6-C6-N1	-2.40	114.05	119.20

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 9 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	7	SAM	3	0
3	B	8	SAM	6	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	213/222 (95%)	0.33	11 (5%) 31 24	36, 63, 121, 133	0
1	B	193/222 (86%)	0.23	7 (3%) 46 38	31, 67, 107, 119	0
All	All	406/444 (91%)	0.28	18 (4%) 38 32	31, 64, 113, 133	0

The worst 5 of 18 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	2101	ARG	4.4
1	A	2103	GLY	3.9
1	B	2271	GLY	3.8
1	A	2100	THR	3.2
1	B	2123	PRO	3.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	SAM	B	8	27/27	0.72	0.32	1.01	94,99,102,103	0
3	SAM	A	7	27/27	0.91	0.18	-0.34	69,72,82,82	0
2	ZN	A	1	1/1	0.99	0.10	-1.12	74,74,74,74	1
2	ZN	A	2	1/1	0.98	0.10	-1.95	86,86,86,86	1
2	ZN	A	4	1/1	0.73	0.08	-2.10	71,71,71,71	1
2	ZN	B	3	1/1	0.89	0.06	-2.86	72,72,72,72	1
2	ZN	B	5	1/1	0.97	0.04	-3.33	81,81,81,81	1
2	ZN	B	6	1/1	0.84	0.06	-	83,83,83,83	1

6.5 Other polymers [i](#)

There are no such residues in this entry.