



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 11:21 AM GMT

PDB ID : 3OSN
Title : Structural Basis for Proficient Incorporation of dTTP Opposite O6-Methylguanine by Human DNA Polymerase Iota
Authors : Pence, M.G.
Deposited on : 2010-09-09
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	4755 (1.90-1.90)
Clashscore	102246	5398 (1.90-1.90)
Ramachandran outliers	100387	5338 (1.90-1.90)
Sidechain outliers	100360	5339 (1.90-1.90)
RSRZ outliers	91569	4766 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	B	18	
1	C	18	
2	A	420	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	TTP	A	421	-	-	X	X
4	MG	A	422	-	-	-	X

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 3611 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a DNA chain called 5'-D(*TP*CP*TP*(6OG)P*GP*GP*GP*TP*CP*CP*TP*AP*GP*GP*AP*CP*CP*(DOC))-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	B	7	Total	C	N	O	P	0	0	0
			142	67	29	39	7			
1	C	8	Total	C	N	O	P	0	0	0
			167	79	30	50	8			

- Molecule 2 is a protein called DNA polymerase iota.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	A	385	Total	C	N	O	S	0	0	0
			2999	1890	525	563	21			

- Molecule 3 is THYMIDINE-5'-TRIPHOSPHATE (three-letter code: TTP) (formula: C₁₀H₁₇N₂O₁₄P₃).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			29	10	2	14	3		

- Molecule 4 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	2	Total	Mg	0	0
			2	2		

- Molecule 5 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total	Na	0	0
			1	1		

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	B	13	Total	O	0	0
			13	13		
6	C	19	Total	O	0	0
			19	19		
6	A	239	Total	O	0	0
			239	239		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: 5'-D(*TP*CP*TP*(6OG)P*GP*GP*GP*TP*CP*CP*TP*AP*GP*GP*AP*CP*CP*(DOC))-3'

Chain B: 




- Molecule 1: 5'-D(*TP*CP*TP*(6OG)P*GP*GP*GP*TP*CP*CP*TP*AP*GP*GP*AP*CP*CP*(DOC))-3'

Chain C: 



- Molecule 2: DNA polymerase iota

Chain A: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 65 2 2	Depositor
Cell constants a, b, c, α , β , γ	97.97Å 97.97Å 202.76Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	47.62 – 1.90 47.62 – 1.90	Depositor EDS
% Data completeness (in resolution range)	99.5 (47.62-1.90) 99.5 (47.62-1.90)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.26 (at 1.90Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.208 , 0.241 0.238 , 0.269	Depositor DCC
R_{free} test set	2323 reflections (5.33%)	DCC
Wilson B-factor (Å ²)	34.2	Xtriage
Anisotropy	0.178	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 37.7	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 45962 reflections	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	3611	wwPDB-VP
Average B, all atoms (Å ²)	52.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.97% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NA, MG, 6OG, TTP, DOC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	B	1.33	0/139	2.54	11/212 (5.2%)
1	C	1.63	0/160	2.47	13/245 (5.3%)
2	A	0.99	5/3040 (0.2%)	1.01	15/4101 (0.4%)
All	All	1.05	5/3339 (0.1%)	1.24	39/4558 (0.9%)

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	267	LYS	C-O	5.76	1.34	1.23
2	A	265	SER	C-O	5.62	1.34	1.23
2	A	203	LYS	CD-CE	-5.24	1.38	1.51
2	A	383	MET	CG-SD	5.12	1.94	1.81
2	A	72	LYS	C-O	5.09	1.33	1.23

All (39) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	9	DG	O4'-C1'-N9	15.79	119.05	108.00
2	A	314	GLU	N-CA-CB	10.01	128.62	110.60
1	C	847	DT	O4'-C1'-N1	-9.57	101.30	108.00
2	A	184	ARG	NE-CZ-NH2	-9.15	115.72	120.30
2	A	184	ARG	NE-CZ-NH1	9.00	124.80	120.30
1	C	847	DT	C6-C5-C7	-8.59	117.75	122.90
1	C	842	DG	O4'-C4'-C3'	-8.22	101.07	106.00
1	B	11	DC	O5'-P-OP2	-7.36	99.08	105.70
1	B	12	DC	O4'-C4'-C3'	-6.90	101.74	104.50
1	C	841	DG	O4'-C1'-N9	6.79	112.75	108.00
1	C	847	DT	C4-C5-C7	6.78	123.07	119.00
2	A	312	SER	N-CA-CB	6.72	120.59	110.50
1	C	843	DG	C8-N9-C4	-6.53	103.79	106.40

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	9	DG	N9-C4-C5	6.52	108.01	105.40
2	A	353	LYS	N-CA-C	6.43	128.36	111.00
1	B	10	DA	N1-C6-N6	-6.40	114.76	118.60
1	B	9	DG	N1-C6-O6	-6.36	116.09	119.90
2	A	312	SER	CA-C-N	-6.33	103.27	117.20
1	C	844	DT	C2-N3-C4	-6.26	123.44	127.20
1	C	843	DG	N9-C4-C5	6.06	107.82	105.40
1	C	846	DC	O4'-C1'-N1	-5.91	103.86	108.00
2	A	350	SER	N-CA-C	-5.89	95.10	111.00
1	C	845	DC	P-O3'-C3'	5.88	126.76	119.70
2	A	313	SER	N-CA-C	-5.85	95.20	111.00
2	A	35	LEU	CB-CG-CD1	5.68	120.65	111.00
1	B	12	DC	C5-C4-N4	-5.63	116.26	120.20
2	A	313	SER	CB-CA-C	5.62	120.77	110.10
2	A	123	LEU	CB-CG-CD2	5.54	120.42	111.00
2	A	356	GLY	N-CA-C	5.39	126.58	113.10
1	B	9	DG	C6-C5-N7	5.38	133.62	130.40
1	C	844	DT	O4'-C4'-C3'	-5.33	102.37	104.50
1	B	11	DC	C4'-C3'-C2'	-5.32	98.31	103.10
2	A	314	GLU	N-CA-C	-5.30	96.69	111.00
1	B	10	DA	C4'-C3'-C2'	-5.23	98.39	103.10
2	A	352	GLU	N-CA-C	-5.23	96.89	111.00
1	C	841	DG	C1'-O4'-C4'	-5.19	104.91	110.10
1	B	11	DC	N1-C2-O2	-5.14	115.82	118.90
1	C	843	DG	O4'-C4'-C3'	-5.13	102.45	104.50
2	A	35	LEU	CA-CB-CG	-5.05	103.69	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	142	0	78	1	1
1	C	167	0	93	4	1
2	A	2999	0	3073	67	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	29	0	13	9	0
4	A	2	0	0	0	0
5	A	1	0	0	0	0
6	A	239	0	0	5	0
6	B	13	0	0	0	0
6	C	19	0	0	0	0
All	All	3611	0	3257	76	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

All (76) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:310:LYS:HE3	2:A:402:HIS:CB	1.67	1.23
2:A:399:MET:N	2:A:400:PRO:HD3	1.36	1.19
2:A:73:LEU:HD23	2:A:73:LEU:N	1.56	1.13
2:A:350:SER:CB	2:A:351:SER:HA	1.66	1.11
2:A:399:MET:N	2:A:400:PRO:CD	2.20	1.05
2:A:267:LYS:H	2:A:267:LYS:HD3	1.15	1.03
2:A:350:SER:HB3	2:A:351:SER:HA	1.00	1.00
2:A:144:GLN:O	2:A:145:SER:OG	1.80	0.99
2:A:310:LYS:CE	2:A:402:HIS:CB	2.41	0.99
2:A:249:CYS:HB2	6:A:644:HOH:O	1.62	0.98
2:A:350:SER:HB3	2:A:351:SER:CA	1.95	0.96
2:A:398:LYS:C	2:A:400:PRO:HD3	1.86	0.96
2:A:73:LEU:CD2	2:A:73:LEU:N	2.30	0.93
2:A:315:VAL:O	2:A:315:VAL:HG13	1.69	0.93
2:A:164:ASN:H	2:A:170:HIS:HD2	1.11	0.91
2:A:290:ASN:HB2	6:A:560:HOH:O	1.71	0.91
2:A:267:LYS:N	2:A:267:LYS:HD3	1.93	0.80
2:A:72:LYS:C	2:A:73:LEU:HD23	2.04	0.77
2:A:164:ASN:H	2:A:170:HIS:CD2	2.01	0.75
2:A:399:MET:H	2:A:400:PRO:HD3	1.46	0.75
2:A:189:ASN:HB3	6:A:626:HOH:O	1.85	0.75
2:A:267:LYS:H	2:A:267:LYS:CD	1.97	0.72
2:A:316:GLU:O	2:A:320:LYS:HG2	1.89	0.72
2:A:73:LEU:H	2:A:73:LEU:HD23	1.57	0.69
2:A:35:LEU:HD12	2:A:195:GLY:HA3	1.74	0.69
2:A:144:GLN:C	2:A:145:SER:HG	1.89	0.68
2:A:315:VAL:O	2:A:315:VAL:CG1	2.41	0.68

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:421:TTP:H5'1	3:A:421:TTP:PB	2.38	0.64
2:A:350:SER:CB	2:A:351:SER:CA	2.58	0.62
3:A:421:TTP:H5'2	3:A:421:TTP:C6	2.38	0.59
2:A:396:ASN:O	2:A:396:ASN:ND2	2.30	0.58
2:A:325:LEU:CD1	2:A:329:LEU:HD22	2.33	0.57
1:C:844:DT:H2''	1:C:845:DC:H5'	1.85	0.57
2:A:202:ASN:ND2	2:A:205:LEU:H	2.02	0.56
2:A:105:MET:HE1	6:A:524:HOH:O	2.05	0.56
3:A:421:TTP:O2B	3:A:421:TTP:H3'	2.06	0.55
3:A:421:TTP:H6	3:A:421:TTP:H5'2	1.70	0.55
3:A:421:TTP:O2B	3:A:421:TTP:H5'1	2.05	0.55
2:A:202:ASN:C	2:A:202:ASN:HD22	2.10	0.54
2:A:112:LEU:HD23	2:A:112:LEU:C	2.28	0.54
1:C:841:DG:OP2	2:A:307:SER:HB2	2.10	0.52
2:A:310:LYS:HE2	2:A:402:HIS:CB	2.35	0.50
2:A:304:GLU:HG3	2:A:328:LEU:HG	1.93	0.49
2:A:235:HIS:CE1	2:A:237:LYS:HE3	2.48	0.49
2:A:316:GLU:OE1	2:A:316:GLU:HA	2.13	0.48
2:A:398:LYS:C	2:A:400:PRO:CD	2.70	0.48
2:A:68:TYR:O	2:A:72:LYS:HG3	2.13	0.48
2:A:371:LYS:N	2:A:371:LYS:HD3	2.29	0.48
2:A:309:LYS:HD3	2:A:309:LYS:HA	1.54	0.47
2:A:300:GLN:NE2	6:A:639:HOH:O	2.46	0.47
1:C:840:6OG:H8	2:A:59:GLN:OE1	2.15	0.46
2:A:266:PRO:O	2:A:270:GLU:HG3	2.16	0.46
2:A:370:GLN:HA	2:A:370:GLN:OE1	2.14	0.46
2:A:61:TYR:N	2:A:61:TYR:CD1	2.84	0.46
2:A:235:HIS:ND1	2:A:237:LYS:HE3	2.31	0.45
2:A:314:GLU:O	2:A:315:VAL:HG12	2.16	0.45
2:A:196:CYS:HA	2:A:217:GLN:O	2.17	0.45
2:A:90:GLN:OE1	2:A:90:GLN:N	2.30	0.44
1:B:13:DOC:H2'	3:A:421:TTP:C6	2.53	0.44
2:A:267:LYS:N	2:A:267:LYS:CD	2.67	0.44
2:A:365:PRO:HB2	2:A:368:VAL:HG23	1.99	0.44
2:A:72:LYS:C	2:A:73:LEU:CD2	2.81	0.43
2:A:396:ASN:ND2	2:A:400:PRO:HA	2.32	0.43
2:A:249:CYS:SG	2:A:273:LEU:HD21	2.57	0.43
2:A:237:LYS:HB3	2:A:237:LYS:HE2	1.88	0.43
2:A:126:ASP:OD1	3:A:421:TTP:O5'	2.36	0.43
2:A:184:ARG:HD2	2:A:195:GLY:O	2.19	0.43
2:A:82:ARG:O	2:A:86:GLU:HB2	2.19	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:421:TTP:H6	3:A:421:TTP:C5'	2.33	0.42
2:A:66:CYS:HB2	2:A:70:ALA:HB3	2.02	0.42
1:C:840:6OG:O6	3:A:421:TTP:O4	2.39	0.41
2:A:303:SER:O	2:A:304:GLU:HG2	2.20	0.41
2:A:338:LYS:HA	2:A:339:PRO:HD3	1.90	0.41
2:A:105:MET:HB2	2:A:105:MET:HE3	1.95	0.40
2:A:248:LYS:HD3	2:A:248:LYS:HA	1.88	0.40
2:A:170:HIS:HE1	2:A:224:GLU:OE2	2.04	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:7:DA:P	1:C:847:DT:O3'[10_665]	1.92	0.28

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	A	381/420 (91%)	369 (97%)	12 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
2	A	342/376 (91%)	306 (90%)	36 (10%)	8 3

All (36) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	A	26	SER
2	A	27	SER
2	A	35	LEU
2	A	51	LYS
2	A	62	LEU
2	A	73	LEU
2	A	80	ASN
2	A	86	GLU
2	A	117	SER
2	A	123	LEU
2	A	142	GLN
2	A	149	SER
2	A	160	ASN
2	A	170	HIS
2	A	202	ASN
2	A	237	LYS
2	A	248	LYS
2	A	265	SER
2	A	267	LYS
2	A	272	GLU
2	A	309	LYS
2	A	313	SER
2	A	314	GLU
2	A	315	VAL
2	A	325	LEU
2	A	329	LEU
2	A	330	ASN
2	A	334	GLN
2	A	344	LEU
2	A	370	GLN
2	A	371	LYS
2	A	378	ASP
2	A	396	ASN
2	A	397	VAL
2	A	404	THR
2	A	405	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such

sidechains are listed below:

Mol	Chain	Res	Type
2	A	58	GLN
2	A	170	HIS
2	A	202	ASN
2	A	262	GLN
2	A	290	ASN
2	A	412	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	DOC	B	13	1	11,19,20	2.35	3 (27%)	14,26,29	2.62	1 (7%)
1	6OG	C	840	-	16,25,26	1.41	4 (25%)	21,36,39	4.73	7 (33%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	DOC	B	13	1	-	0/3/18/19	0/2/2/2
1	6OG	C	840	-	-	0/5/23/24	0/3/3/3

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	840	6OG	C5-C4	-2.24	1.35	1.40

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	840	6OG	C5-N7	-2.12	1.32	1.39
1	C	840	6OG	C5'-C4'	2.54	1.59	1.51
1	B	13	DOC	C4-N3	3.34	1.41	1.35
1	C	840	6OG	O4'-C1'	3.51	1.50	1.42
1	B	13	DOC	O4'-C1'	3.92	1.51	1.42
1	B	13	DOC	C6-N1	5.08	1.42	1.35

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	840	6OG	C-O6-C6	-19.62	96.90	117.26
1	B	13	DOC	O4'-C4'-C5'	-9.17	96.01	109.54
1	C	840	6OG	O4'-C1'-N9	-5.30	98.55	107.72
1	C	840	6OG	N3-C2-N1	-2.65	123.40	127.44
1	C	840	6OG	O4'-C4'-C5'	2.06	116.71	109.32
1	C	840	6OG	C2'-C1'-N9	2.11	119.29	114.16
1	C	840	6OG	O6-C6-N1	3.35	123.82	119.01
1	C	840	6OG	C4-C5-N7	4.17	113.32	109.48

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	B	13	DOC	1	0
1	C	840	6OG	2	0

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 3 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length

(or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	TTP	A	421	4	21,30,30	2.35	7 (33%)	31,47,47	2.69	10 (32%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	TTP	A	421	4	-	0/18/34/34	0/2/2/2

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	421	TTP	C2'-C3'	2.13	1.58	1.52
3	A	421	TTP	O4'-C4'	2.47	1.50	1.45
3	A	421	TTP	PB-O1B	3.33	1.63	1.51
3	A	421	TTP	C4-N3	3.64	1.39	1.33
3	A	421	TTP	C2'-C1'	3.95	1.63	1.52
3	A	421	TTP	C6-N1	4.78	1.42	1.35
3	A	421	TTP	O3'-C3'	5.64	1.56	1.43

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	421	TTP	O4'-C1'-N1	-9.19	91.81	107.72
3	A	421	TTP	C5'-C4'-C3'	-5.81	77.78	114.64
3	A	421	TTP	O4'-C4'-C5'	-3.53	96.70	109.32
3	A	421	TTP	C5-C4-N3	-3.26	121.50	125.14
3	A	421	TTP	O3G-PG-O2G	-3.03	95.82	107.38
3	A	421	TTP	O2B-PB-O3A	-2.03	95.87	105.09
3	A	421	TTP	C5M-C5-C6	2.14	122.92	118.62
3	A	421	TTP	O3'-C3'-C2'	2.77	119.91	110.74
3	A	421	TTP	O4'-C4'-C3'	2.82	112.75	105.67
3	A	421	TTP	C4-N3-C2	5.02	119.58	115.25

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 9 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	421	TTP	9	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	B	6/18 (33%)	-0.01	0	100 100	34, 50, 57, 59	0
1	C	7/18 (38%)	-0.08	0	100 100	29, 33, 45, 59	0
2	A	385/420 (91%)	1.29	68 (17%)	2 2	20, 47, 120, 168	0
All	All	398/456 (87%)	1.25	68 (17%)	2 2	20, 47, 120, 168	0

All (68) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	A	355	TYR	14.8
2	A	401	PHE	12.2
2	A	311	CYS	12.1
2	A	351	SER	10.6
2	A	350	SER	9.9
2	A	399	MET	9.7
2	A	354	HIS	9.6
2	A	377	TYR	8.0
2	A	395	VAL	6.9
2	A	400	PRO	6.5
2	A	397	VAL	6.3
2	A	25	ALA	6.2
2	A	349	TYR	5.7
2	A	313	SER	5.6
2	A	315	VAL	5.3
2	A	398	LYS	5.1
2	A	316	GLU	5.1
2	A	368	VAL	5.0
2	A	391	PHE	4.9
2	A	348	ARG	4.4
2	A	352	GLU	4.2
2	A	148	LEU	4.0
2	A	353	LYS	4.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	A	396	ASN	3.9
2	A	312	SER	3.8
2	A	61	TYR	3.6
2	A	403	LEU	3.5
2	A	390	LEU	3.5
2	A	367	HIS	3.5
2	A	364	ILE	3.4
2	A	381	THR	3.4
2	A	344	LEU	3.3
2	A	402	HIS	3.2
2	A	26	SER	3.2
2	A	78	LEU	3.1
2	A	378	ASP	3.1
2	A	356	GLY	3.1
2	A	309	LYS	3.0
2	A	144	GLN	3.0
2	A	333	CYS	2.9
2	A	82	ARG	2.9
2	A	142	GLN	2.8
2	A	404	THR	2.8
2	A	273	LEU	2.7
2	A	277	VAL	2.7
2	A	244	TYR	2.6
2	A	389	LYS	2.6
2	A	141	GLN	2.6
2	A	145	SER	2.5
2	A	275	ILE	2.4
2	A	308	PHE	2.3
2	A	346	ILE	2.3
2	A	371	LYS	2.3
2	A	62	LEU	2.3
2	A	383	MET	2.3
2	A	69	GLU	2.3
2	A	334	GLN	2.3
2	A	317	ALA	2.2
2	A	358	GLU	2.2
2	A	52	ASP	2.2
2	A	306	ASP	2.2
2	A	233	LEU	2.2
2	A	319	ASN	2.1
2	A	269	LEU	2.1
2	A	84	ALA	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	A	279	GLN	2.1
2	A	138	LYS	2.0
2	A	272	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
1	DOC	B	13	18/19	0.98	0.13	-	30,31,34,34	0
1	6OG	C	840	23/24	0.86	0.14	-	28,37,49,50	0

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	TTP	A	421	29/29	0.82	0.26	3.45	12,33,38,44	25
4	MG	A	422	1/1	0.94	0.23	2.90	31,31,31,31	1
5	NA	A	423	1/1	0.96	0.09	-1.22	34,34,34,34	0
4	MG	A	424	1/1	0.66	0.39	-	30,30,30,30	1

6.5 Other polymers [i](#)

There are no such residues in this entry.