



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 04:56 AM GMT

PDB ID : 2OT3
Title : Crystal structure of rabex-5 VPS9 domain in complex with nucleotide free RAB21
Authors : Delprato, A.; Lambright, D.
Deposited on : 2007-02-07
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	3939 (2.10-2.10)
Clashscore	102246	4460 (2.10-2.10)
Ramachandran outliers	100387	4413 (2.10-2.10)
Sidechain outliers	100360	4414 (2.10-2.10)
RSRZ outliers	91569	3948 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	274	
2	B	170	

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 3642 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Rab5 GDP/GTP exchange factor.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	253	1970	1256	327	370	17	1	0	0

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	124	GLY	-	CLONING ARTIFACT	UNP Q9UJ41
A	125	SER	-	CLONING ARTIFACT	UNP Q9UJ41
A	126	HIS	-	EXPRESSION TAG	UNP Q9UJ41
A	127	HIS	-	EXPRESSION TAG	UNP Q9UJ41
A	128	HIS	-	EXPRESSION TAG	UNP Q9UJ41
A	129	HIS	-	EXPRESSION TAG	UNP Q9UJ41
A	130	HIS	-	EXPRESSION TAG	UNP Q9UJ41
A	131	HIS	-	EXPRESSION TAG	UNP Q9UJ41

- Molecule 2 is a protein called Ras-related protein Rab-21.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	157	1209	772	207	223	7	0	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	14	GLY	-	CLONING ARTIFACT	UNP Q9UL25
B	15	SER	-	CLONING ARTIFACT	UNP Q9UL25

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	295	Total	O	1	0
			295	295		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	168	Total 168	O 168	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($\text{RSRZ} > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Rab5 GDP/GTP exchange factor



- Molecule 2: Ras-related protein Rab-21



4 Data and refinement statistics

Property	Value	Source
Space group	P 2 ₁ 2 ₁ 2	Depositor
Cell constants a, b, c, α , β , γ	71.90 Å 113.40 Å 60.53 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 2.10 19.88 – 2.10	Depositor EDS
% Data completeness (in resolution range)	99.9 (20.00-2.10) 99.9 (19.88-2.10)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	7.50	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.57 (at 2.09 Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, R_{free}	0.188 , 0.243 0.188 , 0.243	Depositor DCC
R_{free} test set	1500 reflections (5.35%)	DCC
Wilson B-factor (Å ²)	24.5	Xtriage
Anisotropy	0.511	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 55.8	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 29521 reflections	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	3642	wwPDB-VP
Average B, all atoms (Å ²)	33.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.71% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.54	0/2007	0.61	1/2719 (0.0%)
2	B	0.55	0/1228	0.64	0/1655
All	All	0.54	0/3235	0.62	1/4374 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	187	HIS	CB-CA-C	7.84	126.07	110.40

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1970	0	1923	35	0
2	B	1209	0	1194	30	0
3	A	295	0	0	9	0
3	B	168	0	0	19	0
All	All	3642	0	3117	65	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (65) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:159:TYR:HD2	3:B:254:HOH:O	1.42	1.01
2:B:94:GLY:HA3	3:B:308:HOH:O	1.60	1.00
2:B:21:LYS:H	2:B:93:ASN:HD22	1.16	0.93
1:A:216:MET:SD	3:A:423:HOH:O	2.27	0.93
1:A:343:ASN:HD22	1:A:344:PRO:HD2	1.38	0.88
1:A:274:ASP:OD2	1:A:297:HIS:HD2	1.55	0.88
1:A:145:PHE:CZ	1:A:207:ILE:HG22	2.12	0.83
2:B:22:VAL:HG22	3:B:308:HOH:O	1.78	0.83
1:A:193:MET:HE1	1:A:207:ILE:HD11	1.61	0.81
1:A:307:ASN:HB3	3:A:510:HOH:O	1.90	0.71
2:B:134:ILE:HG21	2:B:160:HIS:HB3	1.70	0.71
1:A:161:LYS:NZ	1:A:214:TYR:OH	2.22	0.69
1:A:145:PHE:CE2	1:A:207:ILE:HG22	2.29	0.68
2:B:63:ILE:HD12	2:B:68:VAL:HG21	1.75	0.68
1:A:143:ILE:HG22	1:A:147:LYS:HE2	1.77	0.67
2:B:24:LEU:HD22	2:B:36:VAL:HG21	1.76	0.66
2:B:140:ARG:NH2	3:B:293:HOH:O	2.30	0.65
2:B:133:LYS:HB2	3:B:333:HOH:O	1.97	0.64
2:B:142:VAL:CG1	3:B:314:HOH:O	2.45	0.64
1:A:331:LEU:HG	3:A:552:HOH:O	1.99	0.63
1:A:145:PHE:HZ	1:A:207:ILE:HG22	1.62	0.62
1:A:185:PHE:CE1	1:A:189:VAL:HG21	2.34	0.62
1:A:193:MET:CE	1:A:207:ILE:HD11	2.31	0.61
2:B:102:THR:HG23	3:B:333:HOH:O	2.01	0.60
1:A:190:ALA:HA	3:A:676:HOH:O	2.03	0.59
2:B:123:ASN:N	2:B:123:ASN:OD1	2.33	0.59
2:B:146:GLU:HA	3:B:348:HOH:O	2.01	0.59
2:B:59:LYS:HG2	2:B:61:LEU:HD21	1.85	0.58
1:A:151:LYS:HA	3:A:536:HOH:O	2.05	0.57
1:A:270:LYS:HD3	3:A:511:HOH:O	2.04	0.57
2:B:21:LYS:H	2:B:93:ASN:ND2	1.96	0.55
2:B:82:HIS:HE1	3:B:221:HOH:O	1.89	0.55
1:A:216:MET:HE3	1:A:341:PHE:O	2.06	0.55
2:B:142:VAL:HG13	3:B:314:HOH:O	2.06	0.53
1:A:274:ASP:OD2	1:A:297:HIS:CD2	2.48	0.53
1:A:285:ARG:NH1	1:A:346:ARG:NH1	2.56	0.53
1:A:139:SER:N	3:A:496:HOH:O	2.43	0.51
2:B:27:GLU:O	2:B:30:VAL:HG22	2.13	0.49
2:B:123:ASN:ND2	3:B:281:HOH:O	2.45	0.49
1:A:343:ASN:HD22	1:A:344:PRO:CD	2.15	0.48
1:A:285:ARG:CZ	1:A:346:ARG:NH1	2.77	0.48
2:B:68:VAL:HG13	3:B:285:HOH:O	2.13	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:176:LEU:HD21	3:B:308:HOH:O	2.15	0.47
1:A:230:ASP:O	1:A:234:LYS:HG2	2.16	0.46
2:B:78:GLN:HG3	3:B:188:HOH:O	2.15	0.46
1:A:320:ILE:HD13	1:A:365:PHE:CD2	2.51	0.46
1:A:193:MET:HE2	1:A:193:MET:HB3	1.89	0.46
1:A:175:ILE:HB	3:A:665:HOH:O	2.15	0.45
1:A:247:TRP:CD2	1:A:384:TYR:HB3	2.52	0.45
2:B:21:LYS:N	2:B:93:ASN:HD22	1.98	0.45
2:B:78:GLN:NE2	3:B:188:HOH:O	2.36	0.44
1:A:346:ARG:O	1:A:346:ARG:HG3	2.17	0.44
1:A:281:LYS:O	1:A:287:LYS:NZ	2.48	0.44
2:B:159:TYR:CD2	3:B:254:HOH:O	2.32	0.43
2:B:36:VAL:HG12	2:B:57:LEU:HD12	2.00	0.43
1:A:240:LYS:CB	3:A:534:HOH:O	2.66	0.42
2:B:82:HIS:CE1	3:B:221:HOH:O	2.68	0.42
1:A:200:PRO:HA	1:A:201:PRO:HD3	1.93	0.41
2:B:171:GLU:OE1	2:B:171:GLU:N	2.33	0.41
1:A:179:SER:O	1:A:183:GLN:HG2	2.21	0.41
2:B:149:SER:HB3	3:B:348:HOH:O	2.21	0.41
1:A:145:PHE:O	1:A:148:THR:HB	2.21	0.41
1:A:175:ILE:HD12	1:A:175:ILE:HA	1.88	0.41
2:B:85:GLY:N	3:B:246:HOH:O	2.54	0.40
1:A:325:LYS:HA	1:A:325:LYS:HE2	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	251/274 (92%)	244 (97%)	5 (2%)	2 (1%)	24	17
2	B	153/170 (90%)	151 (99%)	2 (1%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
All	All	404/444 (91%)	395 (98%)	7 (2%)	2 (0%)	34 30

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	195	THR
1	A	197	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	212/253 (84%)	207 (98%)	5 (2%)	57 61
2	B	125/147 (85%)	122 (98%)	3 (2%)	57 61
All	All	337/400 (84%)	329 (98%)	8 (2%)	57 61

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	162	LEU
1	A	164	LEU
1	A	195	THR
1	A	198	LYS
1	A	390	SER
2	B	38	ARG
2	B	123	ASN
2	B	157	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	188	ASN
1	A	297	HIS
1	A	343	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	B	93	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	253/274 (92%)	0.19	22 (8%) 13 17	13, 27, 66, 85	1 (0%)
2	B	157/170 (92%)	0.23	10 (6%) 23 30	15, 27, 46, 54	0
All	All	410/444 (92%)	0.21	32 (7%) 16 22	13, 27, 55, 85	1 (0%)

All (32) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	197	GLY	7.8
1	A	199	VAL	6.1
1	A	203	ARG	5.9
1	A	196	ARG	5.9
1	A	198	LYS	5.7
1	A	200	PRO	5.3
2	B	64	GLY	4.5
1	A	195	THR	4.1
1	A	201	PRO	4.0
2	B	65	GLY	3.7
2	B	67	ARG	3.5
1	A	151	LYS	3.5
1	A	204	VAL	3.4
1	A	349	THR	3.3
2	B	17	ALA	3.3
1	A	194	GLN	3.3
1	A	222	TYR	3.3
1	A	148	THR	3.2
1	A	348	MET	3.0
1	A	144	GLU	2.9
2	B	123	ASN	2.9
1	A	211	ILE	2.7
1	A	149	PHE	2.7
1	A	202	GLU	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	96	ILE	2.6
2	B	66	LYS	2.5
1	A	193	MET	2.4
2	B	62	ASN	2.3
2	B	18	TYR	2.2
1	A	346	ARG	2.2
2	B	182	GLU	2.1
1	A	215	ILE	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.