



wwPDB X-ray Structure Validation Summary Report ⓘ

Jan 31, 2016 – 09:25 PM GMT

PDB ID : 1OXP
Title : ASPARTATE AMINOTRANSFERASE, H-ASP COMPLEX, CLOSED CON-
FORMATION
Authors : Hohenester, E.; Schirmer, T.; Jansonius, J.N.
Deposited on : 1995-12-23
Resolution : 2.50 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	102246	4242 (2.50-2.50)
Ramachandran outliers	100387	4156 (2.50-2.50)
Sidechain outliers	100360	4158 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	401	

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 3397 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called ASPARTATE AMINOTRANSFERASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	401	Total	C	N	O	S	5	0	0
			3161	2004	558	581	18			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	47	PRO	SER	CONFLICT	UNP P00508

- Molecule 2 is 4'-DEOXY-4'-ACETYLYAMINO-PYRIDOXAL-5'-PHOSPHATE (three-letter code: IK2) (formula: $C_{10}H_{15}N_2O_8P$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			21	10	2	8	1		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	215	Total 215	O 215	0	0

Note EDS was not executed.

Chain A: 39% 43% 15%

V370	E371	R372	L373	E376	T305	A237	T244	E245	E246	G247	I248	D249	E250	V251	L252	S253	Q254	S255	A256	A257	K258	N259	M260	G261	L262	Y263	R266	T271	V272	L273	C274	R275	D276	A277	E278	E279	R282	V283	E284	L287	K288	L289	L290	L291	R292	M294	Y295	S296	N297	P298	M300	A303	R303
V380	K383	R386	I387	V389	P313	E314	R316	K317	E318	K319	L320	V321	E322	V323	K324	G325	K326	A327	D328	K329	L330	L331	S332	K333	R334	T335	Q336	L337	V338	L341	K343	E344	G345	S346	R348	K349	K350	K352	K353	K354	K355	K356	I357	C361	P362	L365	P367	R368	C369				
S3	H8	M11	D15	P16	I17	L18	R19	G19	V20	T21	E22	K25	R26	D27	K32	M33	N34	L35	G36	V37	G38	A39	Y40	R41	D42	D43	M44	G45	K46	P47	Y48	V49	L50	V53	R54	K55	A56	E57	A58	M59	K68	E69	Y70	L71	P72	I73	D78	F79	R81	A82			
E85	L88	G89	R90	N91	S92	E93	A94	F95	K96	R99	Y100	V101	T102	Y103	Q104	S107	G108	S111	L112	R113	Q120	R121	F122	F123	R124	F125	N126	R129	D130	V133	Y134	L135	P136	K137	P138	S139	W140	G141	N142	L144	P145	I146	F147	R148	D149	A150	G151	L152	Q154	L155	Q156	A157	Y159
R159	Y160	Y161	D162	P163	K164	L168	D169	F170	T171	E175	D176	I177	S178	K179	I180	P181	E182	K183	S184	I185	I186	H193	N194	P195	T196	G197	V198	R201	R202	E203	Q204	W205	K206	E207	S210	V211	K214	R215	N216	L217	L218	A219	V220	F221	D222	M223	Q226	I233	M234	R235	P236		
A237	W238	A239	L240	R241	T244	E245	E246	G247	I248	D249	E250	V251	L252	S253	Q254	S255	A256	A257	K258	N259	M260	G261	L262	Y263	R266	T271	V272	L273	C274	R275	D276	A277	E278	E279	R282	V283	E284	L287	K288	L289	L290	L291	R292	M294	Y295	S296	N297	P298	M300	A303	R303		
T305	A306	S307	L308	L309	T312	P313	E314	R316	K317	E318	K319	L320	V321	E322	V323	K324	G325	K326	A327	D328	K329	L330	L331	S332	K333	R334	T335	Q336	L337	V338	L341	K343	E344	G345	S346	R348	K349	K350	K352	K353	K354	K355	K356	I357	C361	P362	L365	P367	R368	C369			

4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section will therefore be incomplete.

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	69.50Å 91.55Å 128.36Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	8.00 – 2.50	Depositor
% Data completeness (in resolution range)	(Not available) (8.00-2.50)	Depositor
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	PROLSQ	Depositor
R, R_{free}	(Not available) , (Not available)	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	3397	wwPDB-VP
Average B, all atoms (Å ²)	12.0	wwPDB-VP

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: IK2

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.96	1/3231 (0.0%)	2.47	174/4360 (4.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	96	LYS	CA-CB	-7.89	1.36	1.53

The worst 5 of 174 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	121	ARG	NE-CZ-NH1	21.04	130.82	120.30
1	A	41	ARG	NE-CZ-NH2	-20.30	110.15	120.30
1	A	304	ARG	CD-NE-CZ	18.37	149.32	123.60
1	A	148	ARG	NE-CZ-NH1	18.16	129.38	120.30
1	A	292	ARG	NE-CZ-NH1	-16.72	111.94	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3161	0	3154	134	1
2	A	21	0	10	3	1

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	215	0	0	18	1
All	All	3397	0	3164	135	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 21.

The worst 5 of 135 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:177:ILE:HA	1:A:180:ILE:HD12	1.60	0.83
1:A:233:ILE:HD13	1:A:320:LEU:HD21	1.64	0.80
1:A:370:VAL:HG12	1:A:383:LYS:HE2	1.65	0.78
1:A:350:TRP:HB3	1:A:353:ILE:HD12	1.66	0.78
1:A:338:VAL:HG21	1:A:354:THR:HG23	1.67	0.77

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:413:HOH:O	3:A:413:HOH:O[3_655]	2.08	0.12
1:A:70:TYR:OH	2:A:411:IK2:O1P[3_655]	2.14	0.06

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	399/401 (100%)	366 (92%)	27 (7%)	6 (2%)	13 22

5 of 6 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	92	SER
1	A	202	GLN
1	A	263	TYR
1	A	296	SER
1	A	266	ARG

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	335/335 (100%)	295 (88%)	40 (12%)	6 12

5 of 40 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	168	LEU
1	A	220	TYR
1	A	366	LYS
1	A	195	PRO
1	A	252	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 10 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	234	ASN
1	A	242	HIS
1	A	297	ASN
1	A	156	GLN
1	A	286	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	IK2	A	411	-	17,21,21	2.33	5 (29%)	21,29,29	4.26	13 (61%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	IK2	A	411	-	-	0/9/13/13	0/1/1/1

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	411	IK2	OX-N4A	-3.96	1.40	1.45
2	A	411	IK2	P-O3P	-2.81	1.44	1.54
2	A	411	IK2	C3-C4	-2.34	1.36	1.40
2	A	411	IK2	C5-C4	3.98	1.46	1.40
2	A	411	IK2	C3-C2	6.10	1.45	1.40

The worst 5 of 13 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
2	A	411	IK2	O3-C3-C2	-10.08	100.14	117.66
2	A	411	IK2	C4A-C4-C5	-9.64	111.11	119.71
2	A	411	IK2	C3-C2-N1	-4.21	114.80	120.61
2	A	411	IK2	C5A-C5-C4	-4.06	114.07	121.89
2	A	411	IK2	C5-C6-N1	-2.74	119.10	123.86

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	411	IK2	3	1

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

EDS was not executed - this section will therefore be empty.

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

EDS was not executed - this section will therefore be empty.

6.3 Carbohydrates ⓘ

EDS was not executed - this section will therefore be empty.

6.4 Ligands ⓘ

EDS was not executed - this section will therefore be empty.

6.5 Other polymers ⓘ

EDS was not executed - this section will therefore be empty.