



Full wwPDB X-ray Structure Validation Report ⓘ

Jan 31, 2016 – 09:31 PM GMT

PDB ID : 1PCW
Title : Aquifex aeolicus KDO8PS in complex with cadmium and APP, a bisubstrate inhibitor
Authors : Xu, X.; Wang, J.; Grison, C.; Petek, S.; Coutrot, P.; Birck, M.; Woodard, R.W.; Gatti, D.L.
Deposited on : 2003-05-17
Resolution : 1.85 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1745 (1.86-1.86)
Clashscore	102246	1898 (1.86-1.86)
Ramachandran outliers	100387	1875 (1.86-1.86)
Sidechain outliers	100360	1875 (1.86-1.86)
RSRZ outliers	91569	1747 (1.86-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	267	<div> <div>4%</div> <div>79%</div> <div>16%</div> <div>•</div> </div>
1	B	267	<div> <div>7%</div> <div>75%</div> <div>18%</div> <div>• 5%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	H4P	B	2268[A]	-	-	-	X
3	H4P	B	2268[B]	-	-	-	X

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 4440 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called 2-dehydro-3-deoxyphosphooctonate aldolase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	257	Total	C	N	O	S	0	0	0
			2026	1308	338	374	6			
1	B	254	Total	C	N	O	S	0	0	0
			2009	1299	335	369	6			

- Molecule 2 is CADMIUM ION (three-letter code: CD) (formula: Cd).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total	Cd	0	0
			1	1		
2	A	1	Total	Cd	0	0
			1	1		

- Molecule 3 is 1-DEOXY-6-O-PHOSPHONO-1-[(PHOSPHONOMETHYL)AMINO]-L-THR EO-HEXITOL (three-letter code: H4P) (formula: C₇H₁₉NO₁₁P₂).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total 21	C 7	N 1	O 11	P 2	0	0
3	B	1	Total 42	C 14	N 2	O 22	P 4	0	1


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	194	Total 194	O 194	0	0
4	B	146	Total 146	O 146	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: 2-dehydro-3-deoxyphosphooctonate aldolase



- Molecule 1: 2-dehydro-3-deoxyphosphooctonate aldolase



4 Data and refinement statistics

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, α , β , γ	84.48Å 84.48Å 159.07Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	21.47 – 1.85 21.47 – 1.85	Depositor EDS
% Data completeness (in resolution range)	95.9 (21.47-1.85) 96.0 (21.47-1.85)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	0.12	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.77 (at 1.85Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.196 , 0.221 0.197 , 0.223	Depositor DCC
R_{free} test set	5551 reflections (10.18%)	DCC
Wilson B-factor (Å ²)	31.5	Xtriage
Anisotropy	0.240	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 48.1	EDS
Estimated twinning fraction	0.024 for -h,-k,l	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 56235 reflections	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	4440	wwPDB-VP
Average B, all atoms (Å ²)	34.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.99% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CD, H4P

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.36	1/2066 (0.0%)	0.57	0/2786
1	B	0.32	0/2049	0.56	0/2764
All	All	0.34	1/4115 (0.0%)	0.57	0/5550

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	1072	GLU	CD-OE2	7.08	1.33	1.25

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2026	0	2060	30	0
1	B	2009	0	2048	38	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	21	0	12	0	0
3	B	42	0	27	0	0
4	A	194	0	0	2	0
4	B	146	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	4440	0	4147	65	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (65) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1185:HIS:O	1:A:1188:GLN:HG2	1.71	0.89
1:A:1161:ASN:HD21	1:B:2132:TRP:HE1	1.18	0.88
1:A:1132:TRP:HE1	1:B:2161:ASN:HD21	1.24	0.82
1:B:2238:LEU:HD21	1:B:2246:ILE:HD12	1.63	0.80
1:B:2011:CYS:H	1:B:2237:GLN:NE2	1.84	0.76
1:B:2200:MET:HG3	1:B:2202:GLU:HG2	1.66	0.75
1:A:1174:LYS:HG3	1:A:1264:ILE:HG21	1.72	0.71
1:A:1205:PHE:O	1:A:1208:ILE:HG22	1.98	0.63
1:B:2011:CYS:H	1:B:2237:GLN:HE22	1.45	0.63
1:B:2041:LYS:HD3	1:B:2042:SER:N	2.16	0.60
1:B:2041:LYS:HD3	1:B:2041:LYS:C	2.22	0.60
1:B:2025:GLU:HG2	1:B:2240:LEU:HD22	1.86	0.58
1:A:1212:VAL:HG11	1:A:1250:ILE:HG23	1.85	0.57
1:B:2205:PHE:HB3	1:B:2206:PRO:HD3	1.85	0.57
1:A:1208:ILE:HD11	1:A:1250:ILE:HD11	1.89	0.54
1:A:1041:LYS:C	1:A:1041:LYS:HD3	2.28	0.53
1:A:1041:LYS:HD3	1:A:1042:SER:N	2.24	0.52
1:B:2020:LEU:HD21	1:B:2069:LYS:HG3	1.91	0.52
1:A:1048:ASN:HD22	1:A:1048:ASN:H	1.58	0.51
1:A:1208:ILE:CD1	1:A:1250:ILE:HD11	2.40	0.51
1:B:2263:THR:HG22	1:B:2264:ILE:HG22	1.93	0.51
1:B:2048:ASN:HD22	1:B:2048:ASN:H	1.59	0.50
1:A:1044:PHE:CZ	1:A:1060:LEU:HD23	2.46	0.49
1:B:2124:LYS:HD3	1:B:2152:THR:HB	1.93	0.49
1:A:1201:ARG:HG3	1:A:1204:ILE:HD12	1.93	0.49
4:A:23:HOH:O	1:B:2083:HIS:HD2	1.94	0.49
1:B:2147:LYS:HZ2	1:B:2147:LYS:HB3	1.78	0.49
1:A:1228:GLU:HG3	1:A:1229:LYS:HG3	1.96	0.48
1:B:2055:PHE:CZ	1:B:2057:GLY:HA2	2.49	0.48
1:A:1174:LYS:HG3	1:A:1264:ILE:HD13	1.95	0.47
1:B:2200:MET:HB3	1:B:2203:PHE:HD2	1.78	0.47
1:B:2044:PHE:CE2	1:B:2060:LEU:HD13	2.50	0.47
1:A:1233:ASP:O	1:A:1237:GLN:HG3	2.15	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:2044:PHE:CZ	1:B:2060:LEU:HD13	2.49	0.47
1:B:2212:VAL:HG11	1:B:2250:ILE:HG23	1.97	0.47
1:A:1060:LEU:HD13	1:A:1060:LEU:C	2.35	0.46
1:B:2204:ILE:O	1:B:2208:ILE:HG13	2.16	0.46
1:A:1003:LYS:HG3	1:A:1035:GLU:O	2.16	0.46
1:B:2183:ALA:HB2	1:B:2208:ILE:HG12	1.99	0.45
1:B:2005:LEU:HD12	1:B:2037:GLU:HB3	1.99	0.45
1:A:1020:LEU:HD21	1:A:1069:LYS:HG3	1.98	0.44
1:A:1050:SER:O	1:B:2136:ASN:HB3	2.17	0.44
1:A:1189:LEU:HD21	1:A:1200:MET:CE	2.48	0.44
1:A:1049:ARG:HD2	4:A:216:HOH:O	2.17	0.43
1:B:2182:ASP:HA	1:B:2220:PHE:HB3	2.00	0.43
1:B:2147:LYS:HA	1:B:2147:LYS:HZ3	1.83	0.43
1:B:2048:ASN:HD22	1:B:2048:ASN:N	2.17	0.42
1:B:2162:ASN:ND2	1:B:2188:GLN:O	2.49	0.42
1:B:2200:MET:CG	1:B:2202:GLU:HG2	2.45	0.42
1:A:1089:GLU:HB3	1:A:1090:PRO:CD	2.50	0.42
1:A:1026:ILE:HD12	1:A:1040:PHE:CD1	2.55	0.42
1:B:2228:GLU:HG3	1:B:2229:LYS:HG3	2.02	0.42
1:A:1014:GLU:HG2	1:A:1231:LEU:HD12	2.02	0.41
1:B:2187:VAL:O	1:B:2199:GLY:HA3	2.20	0.41
1:A:1182:ASP:HA	1:A:1220:PHE:HB3	2.02	0.41
1:B:2068:ARG:O	1:B:2072:GLU:HG3	2.20	0.41
1:A:1048:ASN:HD22	1:A:1048:ASN:N	2.16	0.41
1:A:1188:GLN:O	1:A:1190:PRO:HD3	2.21	0.41
1:B:2016:GLU:O	1:B:2020:LEU:HG	2.20	0.41
1:A:1003:LYS:HE2	1:A:1036:VAL:C	2.41	0.41
1:B:2041:LYS:HB2	1:B:2079:THR:CG2	2.51	0.41
1:B:2089:GLU:HB3	1:B:2090:PRO:CD	2.52	0.40
1:B:2060:LEU:HD12	1:B:2064:VAL:HG23	2.03	0.40
1:A:1029:LEU:HD21	1:A:1244:GLU:HB2	2.03	0.40
1:B:2040:PHE:HB3	1:B:2078:ILE:HD13	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries

of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	253/267 (95%)	250 (99%)	3 (1%)	0	100	100
1	B	250/267 (94%)	243 (97%)	7 (3%)	0	100	100
All	All	503/534 (94%)	493 (98%)	10 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	216/224 (96%)	212 (98%)	4 (2%)	65	49
1	B	215/224 (96%)	208 (97%)	7 (3%)	45	25
All	All	431/448 (96%)	420 (97%)	11 (3%)	54	36

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1048	ASN
1	A	1112	LEU
1	A	1154	ARG
1	A	1162	ASN
1	B	2048	ASN
1	B	2147	LYS
1	B	2154	ARG
1	B	2162	ASN
1	B	2188	GLN
1	B	2205	PHE
1	B	2240	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (12) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	1048	ASN
1	A	1058	HIS
1	A	1161	ASN
1	A	1162	ASN
1	B	2048	ASN
1	B	2058	HIS
1	B	2083	HIS
1	B	2122	ASN
1	B	2136	ASN
1	B	2161	ASN
1	B	2162	ASN
1	B	2237	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 2 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	H4P	A	1268	2	18,20,20	4.14	6 (33%)	23,29,29	2.07	8 (34%)
3	H4P	B	2268[A]	2	18,20,20	4.14	6 (33%)	23,29,29	2.52	7 (30%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	H4P	B	2268[B]	-	18,20,20	4.11	6 (33%)	23,29,29	2.01	5 (21%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	H4P	A	1268	2	-	0/25/25/25	0/0/0/0
3	H4P	B	2268[A]	2	-	0/25/25/25	0/0/0/0
3	H4P	B	2268[B]	-	-	0/25/25/25	0/0/0/0

All (18) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	1268	H4P	P-C4	-14.06	1.48	1.81
3	B	2268[A]	H4P	P-C4	-14.04	1.48	1.81
3	B	2268[B]	H4P	P-C4	-13.68	1.49	1.81
3	A	1268	H4P	C3-N	-5.28	1.38	1.47
3	B	2268[B]	H4P	C3-N	-5.13	1.38	1.47
3	B	2268[A]	H4P	C3-N	-4.84	1.39	1.47
3	B	2268[A]	H4P	CA2-CA3	3.53	1.60	1.53
3	A	1268	H4P	CA1-CA2	3.55	1.60	1.53
3	A	1268	H4P	CA4-CA3	3.56	1.60	1.53
3	B	2268[B]	H4P	CA4-CA3	3.57	1.60	1.53
3	B	2268[A]	H4P	CA4-CA3	3.59	1.60	1.53
3	B	2268[B]	H4P	CA2-CA3	3.72	1.61	1.53
3	A	1268	H4P	CA2-CA3	3.73	1.61	1.53
3	B	2268[A]	H4P	CA1-CA2	3.77	1.61	1.53
3	B	2268[B]	H4P	CA1-CA2	4.00	1.61	1.53
3	A	1268	H4P	CA5-CA4	5.78	1.60	1.51
3	B	2268[B]	H4P	CA5-CA4	5.98	1.60	1.51
3	B	2268[A]	H4P	CA5-CA4	6.11	1.61	1.51

All (20) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	2268[B]	H4P	OA1-CA1-C3	-3.99	100.87	110.27
3	B	2268[A]	H4P	OA1-CA1-C3	-3.47	102.09	110.27
3	A	1268	H4P	OA1-CA1-C3	-3.41	102.23	110.27
3	A	1268	H4P	CA1-CA2-CA3	-3.14	107.37	112.47
3	A	1268	H4P	O3P-P-C4	-2.58	101.09	106.69

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	2268[A]	H4P	O2P-P-C4	-2.22	101.88	106.69
3	B	2268[A]	H4P	OP3-PA-OP4	-2.21	100.21	106.56
3	B	2268[B]	H4P	CA1-CA2-CA3	-2.15	108.97	112.47
3	A	1268	H4P	O3P-P-O1P	2.13	117.86	112.40
3	A	1268	H4P	OA1-CA1-CA2	2.67	115.72	109.02
3	A	1268	H4P	OP2-PA-OP4	2.88	114.85	106.56
3	B	2268[B]	H4P	OA3-CA3-CA4	2.97	116.24	108.75
3	B	2268[A]	H4P	OA3-CA3-CA4	3.55	117.70	108.75
3	A	1268	H4P	OA2-CA2-CA3	3.62	118.15	109.45
3	B	2268[A]	H4P	OA2-CA2-CA3	3.99	119.03	109.45
3	B	2268[B]	H4P	OP2-PA-OP4	4.24	118.78	106.56
3	B	2268[A]	H4P	OP2-PA-OP4	4.46	119.42	106.56
3	A	1268	H4P	C4-N-C3	5.11	124.42	112.81
3	B	2268[B]	H4P	OA2-CA2-CA3	5.44	122.51	109.45
3	B	2268[A]	H4P	C4-N-C3	7.82	130.58	112.81

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	257/267 (96%)	0.13	12 (4%) 35 33	22, 29, 47, 61	0
1	B	254/267 (95%)	0.40	20 (7%) 15 15	23, 33, 54, 64	0
All	All	511/534 (95%)	0.26	32 (6%) 23 22	22, 31, 53, 64	0

All (32) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	1192	GLY	6.1
1	B	2264	ILE	6.1
1	B	2263	THR	5.1
1	B	2190	PRO	4.8
1	A	1199	GLY	4.6
1	A	1200	MET	4.6
1	B	2199	GLY	4.4
1	A	1263	THR	4.0
1	A	1264	ILE	4.0
1	A	1228	GLU	3.9
1	B	2228	GLU	3.7
1	B	2200	MET	3.5
1	A	1189	LEU	3.3
1	B	2229	LYS	3.2
1	B	2089	GLU	2.9
1	A	1191	GLY	2.9
1	B	2235	SER	2.9
1	B	2188	GLN	2.8
1	A	1188	GLN	2.8
1	A	1235	SER	2.7
1	A	1061	GLU	2.6
1	B	2189	LEU	2.5
1	A	1017	GLU	2.4
1	B	2116	LYS	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	2147	LYS	2.3
1	B	2003	LYS	2.2
1	B	2061	GLU	2.2
1	B	2022	VAL	2.2
1	B	2017	GLU	2.1
1	B	2143	PHE	2.1
1	B	2187	VAL	2.1
1	B	2069	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	H4P	B	2268[B]	21/21	0.92	0.24	3.55	33,48,51,52	21
3	H4P	B	2268[A]	21/21	0.92	0.24	3.49	32,51,57,57	21
3	H4P	A	1268	21/21	0.93	0.17	0.93	31,49,58,61	0
2	CD	B	2270	1/1	0.95	0.13	-0.12	40,40,40,40	1
2	CD	A	1270	1/1	0.98	0.03	-2.82	32,32,32,32	1

6.5 Other polymers [i](#)

There are no such residues in this entry.