



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 07:35 PM GMT

PDB ID : 4PC1  
Title : Elongation Factor Tu:Ts complex with a bound phosphate  
Authors : Thirup, S.S.  
Deposited on : 2014-04-14  
Resolution : 1.95 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	1833 (1.96-1.96)
Clashscore	102246	1953 (1.96-1.96)
Ramachandran outliers	100387	1936 (1.96-1.96)
Sidechain outliers	100360	1936 (1.96-1.96)
RSRZ outliers	91569	1835 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	394	<div> <div>3%</div> <div>86%</div> <div>6% • 8%</div> </div>
1	B	394	<div> <div>6%</div> <div>87%</div> <div>5% 8%</div> </div>
2	C	282	<div> <div>94%</div> <div>5% •</div> </div>
2	D	282	<div> <div>%</div> <div>95%</div> <div>• • •</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	GOL	A	402	-	-	-	X

## 2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 21400 atoms, of which 10075 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Elongation factor Tu.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	363	Total	C	H	N	O	S	0	5	0
			5658	1786	2840	481	536	15			
1	B	363	Total	C	H	N	O	S	0	5	0
			5658	1786	2840	481	536	15			

- Molecule 2 is a protein called Elongation factor Ts.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	C	280	Total	C	H	N	O	S	0	10	0
			4360	1349	2211	368	417	15			
2	D	278	Total	C	H	N	O	S	0	2	0
			4248	1320	2148	358	411	11			

- Molecule 3 is PHOSPHATE ION (three-letter code: PO4) (formula: O<sub>4</sub>P).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total O P 5 4 1	0	0
3	B	1	Total O P 5 4 1	0	0

- Molecule 4 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C H O 14 3 8 3	0	0
4	C	1	Total C H O 14 3 8 3	0	0
4	D	1	Total C H O 14 3 8 3	0	0

- Molecule 5 is LEAD (II) ION (three-letter code: PB) (formula: Pb).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	C	2	Total Pb 2 2	0	0

- Molecule 6 is 2-AMINO-2-HYDROXYMETHYL-PROPANE-1,3-DIOL (three-letter code: TRS) (formula: C<sub>4</sub>H<sub>12</sub>NO<sub>3</sub>).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
6	C	1	Total	C	H	N	O	0	0
			20	4	12	1	3		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	319	Total	O	0	0
			319	319		
7	B	221	Total	O	0	0
			221	221		
7	C	471	Total	O	0	0
			471	471		
7	D	391	Total	O	0	0
			391	391		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


#### • Molecule 1: Elongation factor Tu




#### • Molecule 1: Elongation factor Tu



#### • Molecule 2: Elongation factor Ts



#### • Molecule 2: Elongation factor Ts



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	73.51Å 109.77Å 195.03Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	68.79 – 1.95 68.79 – 1.90	Depositor EDS
% Data completeness (in resolution range)	98.1 (68.79-1.95) 97.8 (68.79-1.90)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.09	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.44 (at 1.90Å)	Xtriage
Refinement program	PHENIX (phenix.refine: dev_1593)	Depositor
R, $R_{free}$	0.174 , 0.208 0.175 , 0.208	Depositor DCC
$R_{free}$ test set	4523 reflections (3.99%)	DCC
Wilson B-factor (Å <sup>2</sup> )	21.3	Xtriage
Anisotropy	0.311	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.37 , 46.8	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 122983 reflections	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	21400	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	35.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.55% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PB, GOL, PO4, TRS

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.58	0/2885	0.67	0/3905
1	B	0.52	0/2885	0.63	0/3905
2	C	0.74	1/2212 (0.0%)	0.75	2/2964 (0.1%)
2	D	0.71	1/2132 (0.0%)	0.73	3/2862 (0.1%)
All	All	0.63	2/10114 (0.0%)	0.69	5/13636 (0.0%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	262	GLU	CB-CG	-6.52	1.39	1.52
2	D	262	GLU	CB-CG	-5.86	1.41	1.52

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	132	ARG	NE-CZ-NH2	-8.89	115.86	120.30
2	D	202	MET	CG-SD-CE	-7.00	89.00	100.20
2	D	161	ASP	CB-CG-OD1	6.13	123.81	118.30
2	C	132	ARG	NE-CZ-NH2	-6.07	117.27	120.30
2	C	161	ASP	CB-CG-OD1	5.88	123.59	118.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within

the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2818	2840	2825	18	0
1	B	2818	2840	2825	8	0
2	C	2149	2211	2161	11	0
2	D	2100	2148	2134	6	0
3	A	5	0	0	0	0
3	B	5	0	0	0	0
4	A	6	8	8	0	0
4	C	6	8	8	0	0
4	D	6	8	8	0	0
5	C	2	0	0	0	0
6	C	8	12	12	0	0
7	A	319	0	0	10	0
7	B	221	0	0	4	0
7	C	471	0	0	6	1
7	D	391	0	0	4	0
All	All	11325	10075	9981	42	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (42) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:248:LYS:O	7:B:501:HOH:O	1.93	0.84
2:C:195:GLN:NE2	7:C:401:HOH:O	2.12	0.82
1:A:151:MET:SD	7:A:676:HOH:O	2.39	0.79
2:D:28:GLU:OE1	7:D:401:HOH:O	2.03	0.74
1:A:142:ASP:OD2	7:A:501:HOH:O	2.08	0.71
2:D:183:GLU:OE1	7:D:402:HOH:O	2.09	0.70
2:D:117:GLU:OE2	7:D:403:HOH:O	2.10	0.67
2:D:66:ASP:OD2	7:D:404:HOH:O	2.12	0.67
2:C:199:ASP:OD2	7:C:402:HOH:O	2.14	0.65
1:A:74:ARG:NE	7:A:503:HOH:O	2.30	0.64
2:C:150[B]:ARG:HG2	7:C:719:HOH:O	2.02	0.59
1:A:348:GLU:OE2	7:A:502:HOH:O	2.16	0.57
1:A:346:GLY:N	7:A:506:HOH:O	2.39	0.54
2:D:198:LEU:O	2:D:202:MET:HG2	2.08	0.54
1:A:139[A]:MET:HE3	7:A:627:HOH:O	2.06	0.53
1:A:74:ARG:HD2	7:A:519:HOH:O	2.08	0.52
2:C:13:GLU:OE2	7:C:403:HOH:O	2.19	0.51

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:C:216:GLU:HG2	7:C:769:HOH:O	2.12	0.50
1:B:116:ARG:HD2	7:B:635:HOH:O	2.11	0.49
1:A:8:THR:O	1:A:204:ARG:NH2	2.46	0.48
2:C:198:LEU:O	2:C:202:MET:HG2	2.13	0.48
1:A:142:ASP:CG	7:A:501:HOH:O	2.52	0.46
1:A:301:HIS:CE1	1:A:391:VAL:HG11	2.50	0.46
1:A:226:VAL:HG22	1:A:277:LEU:HD22	1.98	0.46
1:B:297:THR:N	7:B:502:HOH:O	2.37	0.46
1:A:145:LEU:HD13	2:C:19[A]:MET:SD	2.56	0.45
2:C:64:LYS:HB2	2:C:96:LEU:HD21	1.99	0.45
2:C:11:LEU:HD22	2:C:26:LEU:HG	1.97	0.45
1:B:294:LYS:O	7:B:502:HOH:O	2.21	0.45
1:B:209:PRO:HB2	1:B:297:THR:HG21	1.98	0.44
2:C:150[A]:ARG:HE	2:C:150[A]:ARG:HB3	1.68	0.44
1:B:230:ARG:NH1	1:B:273:ASN:OD1	2.52	0.43
1:A:277:LEU:N	1:A:277:LEU:HD23	2.34	0.43
1:A:74:ARG:NE	7:A:519:HOH:O	2.51	0.42
2:C:150[A]:ARG:HG3	7:C:719:HOH:O	2.20	0.42
1:A:74:ARG:CD	7:A:519:HOH:O	2.66	0.41
2:D:76:ASN:OD1	2:D:132:ARG:HD2	2.19	0.41
1:B:237:LYS:HD2	1:B:238:VAL:HG12	2.02	0.41
1:A:28:THR:O	1:A:32:THR:HG23	2.21	0.41
1:B:128:PRO:HB2	1:B:129:TYR:CD1	2.55	0.41
1:A:221:SER:C	1:A:223:ARG:H	2.24	0.40
1:A:220:ILE:HG13	1:A:221:SER:N	2.35	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
7:C:759:HOH:O	7:C:816:HOH:O[3_545]	1.98	0.22

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	364/394 (92%)	354 (97%)	10 (3%)	0	100	100
1	B	364/394 (92%)	354 (97%)	8 (2%)	2 (0%)	34	21
2	C	288/282 (102%)	282 (98%)	6 (2%)	0	100	100
2	D	278/282 (99%)	276 (99%)	2 (1%)	0	100	100
All	All	1294/1352 (96%)	1266 (98%)	26 (2%)	2 (0%)	52	43

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	333	ARG
1	B	266	ASP

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	306/326 (94%)	298 (97%)	8 (3%)	54	43
1	B	306/326 (94%)	299 (98%)	7 (2%)	58	50
2	C	227/219 (104%)	224 (99%)	3 (1%)	76	72
2	D	218/219 (100%)	217 (100%)	1 (0%)	92	91
All	All	1057/1090 (97%)	1038 (98%)	19 (2%)	68	60

All (19) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	16	THR
1	A	70	ASP
1	A	141	ASP
1	A	223	ARG
1	A	262	ARG
1	A	277	LEU

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	283	ARG
1	A	324	LYS
1	B	207	ASP
1	B	217	VAL
1	B	233	ARG
1	B	237	LYS
1	B	256	THR
1	B	286	ILE
1	B	381	ARG
2	C	150[A]	ARG
2	C	150[B]	ARG
2	C	176	LYS
2	D	234	VAL

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 2 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	PO4	A	401	-	4,4,4	1.22	0	6,6,6	0.34	0
4	GOL	A	402	-	5,5,5	0.35	0	5,5,5	0.58	0
3	PO4	B	401	-	4,4,4	0.97	0	6,6,6	0.36	0
4	GOL	C	303	-	5,5,5	0.52	0	5,5,5	0.71	0
6	TRS	C	304	-	7,7,7	1.03	1 (14%)	9,9,9	1.03	1 (11%)
4	GOL	D	301	-	5,5,5	0.66	0	5,5,5	0.16	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	PO4	A	401	-	-	0/0/0/0	0/0/0/0
4	GOL	A	402	-	-	0/4/4/4	0/0/0/0
3	PO4	B	401	-	-	0/0/0/0	0/0/0/0
4	GOL	C	303	-	-	0/4/4/4	0/0/0/0
6	TRS	C	304	-	-	0/9/9/9	0/0/0/0
4	GOL	D	301	-	-	0/4/4/4	0/0/0/0

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	C	304	TRS	C-N	-2.30	1.47	1.50

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	C	304	TRS	C1-C-N	-2.12	104.24	108.09

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	363/394 (92%)	0.05	11 (3%) 54 64	14, 32, 56, 138	0
1	B	363/394 (92%)	0.40	25 (6%) 20 30	15, 37, 77, 113	0
2	C	280/282 (99%)	-0.20	0 100 100	10, 18, 38, 74	0
2	D	278/282 (98%)	-0.14	2 (0%) 89 93	11, 21, 46, 95	0
All	All	1284/1352 (94%)	0.05	38 (2%) 54 64	10, 27, 66, 138	0

All (38) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	223	ARG	9.7
1	B	224	GLY	6.4
1	B	221	SER	6.1
1	B	223	ARG	6.0
1	A	224	GLY	5.8
1	B	265	LEU	5.2
1	A	220	ILE	5.1
1	B	220	ILE	4.9
1	B	222	GLY	4.9
1	B	264	LEU	4.5
1	A	222	GLY	4.3
1	A	221	SER	4.2
1	B	218	PHE	4.0
1	B	256	THR	4.0
1	B	266	ASP	3.4
1	B	96	ALA	3.3
1	B	97	GLN	3.0
1	B	95	ALA	2.9
1	B	278	LEU	2.9
1	A	41	GLY	2.8
1	B	238	VAL	2.8

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	261	PHE	2.8
1	B	284	GLU	2.8
1	A	277	LEU	2.6
1	B	283	ARG	2.6
1	B	217	VAL	2.5
1	B	258	VAL	2.5
1	A	8	THR	2.4
1	B	237	LYS	2.4
1	B	261	PHE	2.3
1	B	281	ILE	2.3
1	A	283	ARG	2.2
2	D	267	VAL	2.2
2	D	279	SER	2.2
1	B	236	ILE	2.1
1	A	95	ALA	2.1
1	B	279	ARG	2.1
1	B	239	GLY	2.1

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
4	GOL	A	402	6/6	0.33	0.32	4.62	63,76,79,80	0
3	PO4	B	401	5/5	0.97	0.12	1.10	24,24,25,29	0
4	GOL	D	301	6/6	0.98	0.10	0.81	16,19,22,23	0
3	PO4	A	401	5/5	0.98	0.12	0.34	18,19,21,29	0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
6	TRS	C	304	8/8	0.95	0.09	-0.54	23,29,30,31	0
4	GOL	C	303	6/6	0.97	0.09	-0.55	22,26,29,30	0
5	PB	C	302	1/1	0.99	0.10	-	29,29,29,29	1
5	PB	C	301	1/1	0.99	0.11	-	27,27,27,27	1

## 6.5 Other polymers [i](#)

There are no such residues in this entry.