



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 11:39 AM GMT

PDB ID : 3PIQ
Title : Crystal structure of human 2909 Fab, a quaternary structure-specific antibody against HIV-1
Authors : Changela, A.; Gorny, M.K.; Zolla-Pazner, S.; Kwong, P.D.
Deposited on : 2010-11-07
Resolution : 3.33 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.33 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1198 (3.40-3.24)
Clashscore	102246	1280 (3.40-3.24)
Ramachandran outliers	100387	1260 (3.40-3.24)
Sidechain outliers	100360	1259 (3.40-3.24)
RSRZ outliers	91569	1203 (3.40-3.24)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	231	<div> <div>4%</div> <div>65% 27% 7%</div> </div>
1	C	231	<div> <div>3%</div> <div>70% 25% . .</div> </div>
1	E	231	<div> <div>66% 27% 7%</div> </div>
1	G	231	<div> <div>5%</div> <div>59% 35% . .</div> </div>
1	H	231	<div> <div>%</div> <div>67% 29% . .</div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	J	231	<div><div></div><div>8%</div><div>63%</div><div>31%</div><div></div><div></div></div>
2	B	211	<div><div></div><div>%</div><div>82%</div><div>17%</div><div></div><div></div></div>
2	D	211	<div><div></div><div>2%</div><div>76%</div><div>22%</div><div></div><div></div></div>
2	F	211	<div><div></div><div></div><div>74%</div><div>24%</div><div></div><div></div></div>
2	I	211	<div><div></div><div>%</div><div>77%</div><div>20%</div><div></div><div></div></div>
2	K	211	<div><div></div><div>6%</div><div>73%</div><div>25%</div><div></div><div></div></div>
2	L	211	<div><div></div><div>%</div><div>78%</div><div>20%</div><div></div><div></div></div>

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 19449 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Human monoclonal antibody 2909 Fab heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	H	224	Total	C	N	O	S	0	0	0
			1709	1082	283	337	7			
1	A	214	Total	C	N	O	S	0	0	0
			1622	1027	272	316	7			
1	C	222	Total	C	N	O	S	0	0	0
			1691	1071	280	333	7			
1	E	214	Total	C	N	O	S	0	0	0
			1622	1027	272	316	7			
1	G	222	Total	C	N	O	S	0	0	0
			1691	1071	280	333	7			
1	J	223	Total	C	N	O	S	0	0	0
			1700	1077	282	334	7			


- Molecule 2 is a protein called Human monoclonal antibody 2909 Fab light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	L	209	Total	C	N	O	S	0	0	0
			1577	987	268	318	4			
2	B	209	Total	C	N	O	S	0	0	0
			1577	987	268	318	4			
2	D	208	Total	C	N	O	S	0	0	0
			1565	978	267	316	4			
2	F	208	Total	C	N	O	S	0	0	0
			1565	978	267	316	4			
2	I	208	Total	C	N	O	S	0	0	0
			1565	978	267	316	4			
2	K	208	Total	C	N	O	S	0	0	0
			1565	978	267	316	4			


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Human monoclonal antibody 2909 Fab heavy chain



- Molecule 1: Human monoclonal antibody 2909 Fab heavy chain




- Molecule 1: Human monoclonal antibody 2909 Fab heavy chain




- Molecule 1: Human monoclonal antibody 2909 Fab heavy chain







- Molecule 1: Human monoclonal antibody 2909 Fab heavy chain




- Molecule 1: Human monoclonal antibody 2909 Fab heavy chain




- Molecule 2: Human monoclonal antibody 2909 Fab light chain





- Molecule 2: Human monoclonal antibody 2909 Fab light chain








- Molecule 2: Human monoclonal antibody 2909 Fab light chain





- Molecule 2: Human monoclonal antibody 2909 Fab light chain



- Molecule 2: Human monoclonal antibody 2909 Fab light chain



- Molecule 2: Human monoclonal antibody 2909 Fab light chain



4 Data and refinement statistics

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, α , β , γ	180.37Å 180.37Å 222.53Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	22.38 – 3.33 22.37 – 3.33	Depositor EDS
% Data completeness (in resolution range)	77.6 (22.38-3.33) 80.9 (22.37-3.33)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.16	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.06 (at 3.30Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.6.4_486)	Depositor
R, R_{free}	0.236 , 0.299 0.232 , 0.291	Depositor DCC
R_{free} test set	2565 reflections (5.14%)	DCC
Wilson B-factor (Å ²)	67.8	Xtriage
Anisotropy	0.986	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.26 , 88.7	EDS
Estimated twinning fraction	0.058 for -h,-k,l	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.43$, $\langle L^2 \rangle = 0.26$	Xtriage
Outliers	2 of 49967 reflections (0.004%)	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	19449	wwPDB-VP
Average B, all atoms (Å ²)	115.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.10% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.21	0/1662	0.39	0/2258
1	C	0.22	0/1735	0.40	0/2361
1	E	0.22	0/1662	0.41	0/2258
1	G	0.23	0/1735	0.39	0/2361
1	H	0.23	0/1753	0.41	0/2384
1	J	0.22	0/1744	0.40	0/2372
2	B	0.21	0/1618	0.39	0/2216
2	D	0.21	0/1605	0.40	0/2198
2	F	0.22	0/1605	0.41	0/2198
2	I	0.21	0/1605	0.40	0/2198
2	K	0.22	0/1605	0.40	0/2198
2	L	0.22	0/1618	0.41	0/2216
All	All	0.22	0/19947	0.40	0/27218

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1622	0	1572	48	0
1	C	1691	0	1620	40	0
1	E	1622	0	1572	43	0
1	G	1691	0	1620	69	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	H	1709	0	1642	53	0
1	J	1700	0	1633	61	0
2	B	1577	0	1513	23	0
2	D	1565	0	1504	39	0
2	F	1565	0	1504	40	0
2	I	1565	0	1504	36	0
2	K	1565	0	1504	38	0
2	L	1577	0	1513	37	0
All	All	19449	0	18701	493	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 13.

The worst 5 of 493 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:F:27:ASN:H	2:F:69:ASN:HD22	1.23	0.87
1:J:53:ASN:HD22	1:J:53:ASN:H	1.22	0.87
1:G:33:THR:HG22	1:G:52:SER:HA	1.55	0.87
1:E:169:VAL:HG21	2:F:160:GLU:HB3	1.58	0.85
1:H:35:HIS:HB2	1:H:93:ALA:HB3	1.58	0.85

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	208/231 (90%)	183 (88%)	23 (11%)	2 (1%)	19	59
1	C	218/231 (94%)	193 (88%)	24 (11%)	1 (0%)	34	72
1	E	208/231 (90%)	185 (89%)	23 (11%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	G	218/231 (94%)	190 (87%)	25 (12%)	3 (1%)	14	50
1	H	220/231 (95%)	195 (89%)	23 (10%)	2 (1%)	21	61
1	J	219/231 (95%)	193 (88%)	23 (10%)	3 (1%)	14	50
2	B	207/211 (98%)	180 (87%)	27 (13%)	0	100	100
2	D	206/211 (98%)	177 (86%)	28 (14%)	1 (0%)	34	72
2	F	206/211 (98%)	172 (84%)	31 (15%)	3 (2%)	13	49
2	I	206/211 (98%)	176 (85%)	27 (13%)	3 (2%)	13	49
2	K	206/211 (98%)	181 (88%)	24 (12%)	1 (0%)	34	72
2	L	207/211 (98%)	179 (86%)	27 (13%)	1 (0%)	34	72
All	All	2529/2652 (95%)	2204 (87%)	305 (12%)	20 (1%)	24	63

5 of 20 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	H	98	ASP
2	L	167	GLN
2	K	51	ASP
1	A	144	ASP
1	C	88	ALA

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	182/197 (92%)	181 (100%)	1 (0%)	92	96
1	C	189/197 (96%)	187 (99%)	2 (1%)	80	91
1	E	182/197 (92%)	181 (100%)	1 (0%)	92	96
1	G	189/197 (96%)	187 (99%)	2 (1%)	80	91
1	H	191/197 (97%)	188 (98%)	3 (2%)	70	87
1	J	190/197 (96%)	185 (97%)	5 (3%)	54	82
2	B	176/178 (99%)	175 (99%)	1 (1%)	90	95

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	D	175/178 (98%)	174 (99%)	1 (1%)	90	95
2	F	175/178 (98%)	173 (99%)	2 (1%)	80	91
2	I	175/178 (98%)	173 (99%)	2 (1%)	80	91
2	K	175/178 (98%)	173 (99%)	2 (1%)	80	91
2	L	176/178 (99%)	175 (99%)	1 (1%)	90	95
All	All	2175/2250 (97%)	2152 (99%)	23 (1%)	80	91

5 of 23 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	F	53	ASP
1	G	27	PHE
2	K	85	ASP
2	F	95(B)	HIS
1	G	64	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 30 such sidechains are listed below:

Mol	Chain	Res	Type
1	E	13	GLN
2	F	69	ASN
1	J	76	ASN
2	F	34	HIS
2	F	108	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	214/231 (92%)	-0.00	9 (4%) 40 37	73, 119, 176, 274	0
1	C	222/231 (96%)	0.08	8 (3%) 46 45	52, 113, 219, 322	0
1	E	214/231 (92%)	-0.43	0 100 100	33, 76, 123, 155	0
1	G	222/231 (96%)	0.16	12 (5%) 29 28	65, 123, 204, 311	0
1	H	224/231 (96%)	-0.32	2 (0%) 85 86	32, 78, 142, 249	0
1	J	223/231 (96%)	0.39	18 (8%) 15 14	60, 140, 252, 443	0
2	B	209/211 (99%)	-0.08	2 (0%) 84 84	65, 115, 177, 250	0
2	D	208/211 (98%)	-0.03	5 (2%) 62 61	56, 118, 204, 295	0
2	F	208/211 (98%)	-0.35	1 (0%) 91 92	44, 91, 146, 261	0
2	I	208/211 (98%)	-0.15	3 (1%) 78 78	50, 109, 180, 234	0
2	K	208/211 (98%)	0.14	13 (6%) 23 23	77, 135, 208, 274	0
2	L	209/211 (99%)	-0.48	2 (0%) 84 84	34, 87, 136, 201	0
All	All	2569/2652 (96%)	-0.09	75 (2%) 55 54	32, 108, 197, 443	0

The worst 5 of 75 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	K	20	ARG	5.4
1	J	157	GLY	5.3
1	J	100(G)	TYR	5.1
1	H	100(A)	TYR	5.0
1	G	191	THR	5.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.