



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 05:10 AM GMT

PDB ID : 2PPA  
Title : Anaerobically manipulated wild type oxidized AfNiR bound to nitrous oxide  
Authors : Tocheva, E.I.; Murphy, M.E.P.  
Deposited on : 2007-04-28  
Resolution : 1.69 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : **NOT EXECUTED**  
EDS : **NOT EXECUTED**  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.69 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	102246	3585 (1.70-1.70)
Ramachandran outliers	100387	3527 (1.70-1.70)
Sidechain outliers	100360	3527 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	341	 92% 6% ..
1	B	341	 94% 5% .
1	C	341	 91% 8% .

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	ACT	C	1506	-	-	X	-

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	ACT	C	503	-	-	X	-

## 2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 8557 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Copper-containing nitrite reductase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	336	Total	C	N	O	S	0	2	0
			2578	1650	435	482	11			
1	B	337	Total	C	N	O	S	0	1	0
			2573	1646	432	484	11			
1	C	336	Total	C	N	O	S	0	1	0
			2571	1645	434	481	11			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	341	ILE	-	EXPRESSION TAG	UNP P38501
A	342	GLU	-	EXPRESSION TAG	UNP P38501
A	343	GLY	-	EXPRESSION TAG	UNP P38501
A	344	ARG	-	EXPRESSION TAG	UNP P38501
B	341	ILE	-	EXPRESSION TAG	UNP P38501
B	342	GLU	-	EXPRESSION TAG	UNP P38501
B	343	GLY	-	EXPRESSION TAG	UNP P38501
B	344	ARG	-	EXPRESSION TAG	UNP P38501
C	341	ILE	-	EXPRESSION TAG	UNP P38501
C	342	GLU	-	EXPRESSION TAG	UNP P38501
C	343	GLY	-	EXPRESSION TAG	UNP P38501
C	344	ARG	-	EXPRESSION TAG	UNP P38501


- Molecule 2 is COPPER (I) ION (three-letter code: CU1) (formula: Cu).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total	Cu	0	0
			1	1		
2	A	1	Total	Cu	0	0
			1	1		
2	C	1	Total	Cu	0	0
			1	1		

- Molecule 3 is COPPER (II) ION (three-letter code: CU) (formula: Cu).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total	Cu	0	0
			1	1		
3	A	1	Total	Cu	0	0
			1	1		
3	C	1	Total	Cu	0	0
			1	1		

- Molecule 4 is ACETATE ION (three-letter code: ACT) (formula: C<sub>2</sub>H<sub>3</sub>O<sub>2</sub>).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			4	2	2		
4	B	1	Total	C	O	0	0
			4	2	2		
4	B	1	Total	C	O	0	0
			4	2	2		
4	B	1	Total	C	O	0	0
			4	2	2		
4	C	1	Total	C	O	0	0
			4	2	2		
4	C	1	Total	C	O	0	0
			4	2	2		
4	C	1	Total	C	O	0	0
			4	2	2		
4	C	1	Total	C	O	0	0
			4	2	2		

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	C	1	Total	C	O	0	0
			4	2	2		
4	C	1	Total	C	O	0	0
			4	2	2		

- Molecule 5 is NITROUS OXIDE (three-letter code: N2O) (formula: N<sub>2</sub>O).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	N	O	0	0
			3	2	1		
5	B	1	Total	N	O	0	0
			3	2	1		

- Molecule 6 is 2-AMINO-2-HYDROXYMETHYL-PROPANE-1,3-DIOL (three-letter code: TRS) (formula: C<sub>4</sub>H<sub>12</sub>NO<sub>3</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	1	Total C O 7 4 3	0	0

- Molecule 7 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	272	Total O 272 272	0	0
7	B	249	Total O 249 249	0	0
7	C	255	Total O 255 255	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

- Molecule 1: Copper-containing nitrite reductase

Chain A: 




- Molecule 1: Copper-containing nitrite reductase

Chain B: 



- Molecule 1: Copper-containing nitrite reductase

Chain C: 





## 4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section will therefore be incomplete.

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	60.65Å 101.62Å 146.04Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.00 – 1.69	Depositor
% Data completeness (in resolution range)	95.6 (30.00-1.69)	Depositor
$R_{merge}$	0.13	Depositor
$R_{sym}$	(Not available)	Depositor
Refinement program	REFMAC 5.2.0019	Depositor
R, $R_{free}$	0.191 , 0.224	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	8557	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	23.0	wwPDB-VP

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: TRS, N2O, CU, CU1, ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.48	0/2649	0.62	0/3612
1	B	0.47	0/2644	0.62	0/3606
1	C	0.46	0/2642	0.59	0/3602
All	All	0.47	0/7935	0.61	0/10820

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	0	1
1	C	0	1
All	All	0	3

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	306	HIS	Peptide
1	B	306	HIS	Peptide
1	C	306	HIS	Peptide

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2578	0	2505	17	0
1	B	2573	0	2496	12	0
1	C	2571	0	2497	18	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
3	C	1	0	0	0	0
4	A	4	0	3	0	0
4	B	12	0	9	0	0
4	C	24	0	18	5	0
5	A	3	0	0	0	0
5	B	3	0	0	0	0
6	A	7	0	9	0	0
7	A	272	0	0	2	0
7	B	249	0	0	0	0
7	C	255	0	0	2	0
All	All	8557	0	7537	45	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (45) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:26:HIS:HE1	1:A:74:VAL:H	1.23	0.83
1:C:26:HIS:HE1	1:C:74:VAL:H	1.28	0.81
1:B:26:HIS:HE1	1:B:74:VAL:H	1.36	0.72
1:C:26:HIS:CE1	1:C:74:VAL:H	2.12	0.65
1:C:22:PRO:HB2	1:C:221:ASN:HD21	1.60	0.65
1:B:26:HIS:CE1	1:B:74:VAL:H	2.14	0.65
1:A:26:HIS:CE1	1:A:74:VAL:H	2.10	0.64
1:C:60:HIS:HD2	7:C:1547:HOH:O	1.85	0.59
1:A:26:HIS:HD2	1:A:27:ALA:O	1.86	0.58

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:26:HIS:HD2	1:C:27:ALA:O	1.89	0.55
1:A:26:HIS:HE1	1:A:74:VAL:N	2.00	0.53
1:B:282:PHE:HD2	4:C:1506:ACT:H2	1.75	0.52
1:A:300:ILE:HD12	1:A:321:LYS:HG2	1.92	0.51
1:A:232:ALA:HB3	1:A:321:LYS:HE2	1.94	0.50
1:C:98:ASP:OD1	4:C:503:ACT:H2	2.13	0.49
1:A:257:ILE:HG21	4:C:503:ACT:H3	1.94	0.49
1:B:98:ASP:OD2	1:B:109:GLY:HA3	2.12	0.49
1:B:22:PRO:HB2	1:B:221:ASN:HD21	1.77	0.49
1:C:130:GLY:HA2	1:C:270:PHE:CD1	2.49	0.47
1:C:98:ASP:OD2	1:C:109:GLY:N	2.47	0.47
1:C:257:ILE:HD12	1:C:302:ALA:HB3	1.96	0.46
1:C:98:ASP:OD2	1:C:109:GLY:HA3	2.16	0.46
1:A:13:PRO:HG2	1:A:37:LYS:HG2	1.98	0.45
1:A:300:ILE:CD1	1:A:321:LYS:HG2	2.46	0.45
1:B:249:ASN:O	1:C:307:ASN:HA	2.17	0.44
1:C:226:ALA:O	1:C:231:LYS:HB2	2.18	0.44
1:C:282:PHE:HB3	4:C:1506:ACT:H1	1.99	0.44
1:A:22:PRO:HB2	1:A:221:ASN:HD21	1.83	0.43
1:A:98:ASP:OD2	1:A:109:GLY:HA3	2.19	0.42
1:A:301:TYR:HB2	1:A:320:PHE:HB2	2.02	0.42
1:B:26:HIS:HD2	1:B:27:ALA:O	2.03	0.42
1:A:249:ASN:O	1:B:307:ASN:HA	2.20	0.42
1:C:132:PHE:CE1	1:C:269:LYS:HE3	2.54	0.42
1:C:64:PHE:CE2	1:C:150:MET:HG2	2.55	0.42
1:B:282:PHE:CD2	4:C:1506:ACT:H2	2.54	0.41
1:C:37:LYS:HE2	7:C:1566:HOH:O	2.19	0.41
1:A:33:GLU:HG2	7:A:1680:HOH:O	2.20	0.41
1:B:305:ASN:O	1:B:311:ALA:HB2	2.21	0.41
1:B:301:TYR:HB2	1:B:320:PHE:HB2	2.03	0.41
1:C:143:PRO:O	1:C:147:VAL:HG22	2.21	0.41
1:B:26:HIS:HE1	1:B:74:VAL:N	2.11	0.41
1:A:253:ARG:HA	1:A:281:TRP:O	2.20	0.41
1:A:33:GLU:HB2	7:A:1725:HOH:O	2.20	0.40
1:C:233:MET:O	1:C:320:PHE:HA	2.21	0.40
1:A:69:PRO:HG3	1:A:179:GLY:HA3	2.03	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	336/341 (98%)	333 (99%)	3 (1%)	0	100	100
1	B	336/341 (98%)	328 (98%)	8 (2%)	0	100	100
1	C	335/341 (98%)	329 (98%)	6 (2%)	0	100	100
All	All	1007/1023 (98%)	990 (98%)	17 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	266/268 (99%)	263 (99%)	3 (1%)	80	69
1	B	266/268 (99%)	262 (98%)	4 (2%)	72	56
1	C	265/268 (99%)	264 (100%)	1 (0%)	93	90
All	All	797/804 (99%)	789 (99%)	8 (1%)	82	72

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	33	GLU
1	A	192	LYS
1	A	221	ASN
1	B	33	GLU
1	B	37	LYS

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	168	LYS
1	B	221	ASN
1	C	166	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (15) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	26	HIS
1	A	77	GLN
1	A	163	HIS
1	A	221	ASN
1	B	26	HIS
1	B	77	GLN
1	B	115	ASN
1	B	163	HIS
1	B	221	ASN
1	C	26	HIS
1	C	60	HIS
1	C	77	GLN
1	C	115	ASN
1	C	221	ASN
1	C	260	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 5.6 Ligand geometry [i](#)

Of 19 ligands modelled in this entry, 6 are monoatomic - leaving 13 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
6	TRS	A	1501	-	6,6,7	0.27	0	3,6,9	1.31	0
4	ACT	A	1510	-	1,3,3	1.23	0	0,3,3	0.00	-
5	N2O	A	503	3	0,2,2	0.00	-	0,1,1	0.00	-
4	ACT	B	1503	-	1,3,3	1.21	0	0,3,3	0.00	-
4	ACT	B	1504	-	1,3,3	1.07	0	0,3,3	0.00	-
4	ACT	B	1505	-	1,3,3	1.29	0	0,3,3	0.00	-
5	N2O	B	503	3	0,2,2	0.00	-	0,1,1	0.00	-
4	ACT	C	1502	-	1,3,3	1.48	0	0,3,3	0.00	-
4	ACT	C	1506	-	1,3,3	0.91	0	0,3,3	0.00	-
4	ACT	C	1507	-	1,3,3	0.81	0	0,3,3	0.00	-
4	ACT	C	1508	-	1,3,3	0.99	0	0,3,3	0.00	-
4	ACT	C	1509	-	1,3,3	0.89	0	0,3,3	0.00	-
4	ACT	C	503	3	1,3,3	0.57	0	0,3,3	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	TRS	A	1501	-	-	0/6/6/9	0/0/0/0
4	ACT	A	1510	-	-	0/0/0/0	0/0/0/0
5	N2O	A	503	3	-	0/0/0/0	0/0/0/0
4	ACT	B	1503	-	-	0/0/0/0	0/0/0/0
4	ACT	B	1504	-	-	0/0/0/0	0/0/0/0
4	ACT	B	1505	-	-	0/0/0/0	0/0/0/0
5	N2O	B	503	3	-	0/0/0/0	0/0/0/0
4	ACT	C	1502	-	-	0/0/0/0	0/0/0/0
4	ACT	C	1506	-	-	0/0/0/0	0/0/0/0
4	ACT	C	1507	-	-	0/0/0/0	0/0/0/0
4	ACT	C	1508	-	-	0/0/0/0	0/0/0/0
4	ACT	C	1509	-	-	0/0/0/0	0/0/0/0
4	ACT	C	503	3	-	0/0/0/0	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	C	1506	ACT	3	0
4	C	503	ACT	2	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

EDS was not executed - this section will therefore be empty.

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

EDS was not executed - this section will therefore be empty.

### 6.3 Carbohydrates ⓘ

EDS was not executed - this section will therefore be empty.

### 6.4 Ligands ⓘ

EDS was not executed - this section will therefore be empty.

### 6.5 Other polymers ⓘ

EDS was not executed - this section will therefore be empty.