



# wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 05:18 AM GMT

PDB ID : 2Q5D  
Title : Crystal Structure of Human Importin Beta bound to the Snurportin1 IBB-domain second crystal form  
Authors : Mitrousis, G.; Cingolani, G.  
Deposited on : 2007-05-31  
Resolution : 3.20 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 3.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	1124 (3.24-3.16)
Clashscore	102246	1024 (3.22-3.18)
Ramachandran outliers	100387	1004 (3.22-3.18)
Sidechain outliers	100360	1003 (3.22-3.18)
RSRZ outliers	91569	1129 (3.24-3.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	876	<div> <div>3%</div> <div>60%</div> <div>33%</div> <div>5% ..</div> </div>
1	B	876	<div> <div>11%</div> <div>51%</div> <div>33%</div> <div>11% . .</div> </div>
2	C	40	<div> <div>50%</div> <div>30%</div> <div>10%</div> <div>10%</div> </div>
2	D	40	<div> <div>10%</div> <div>48%</div> <div>10%</div> <div>43%</div> </div>

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 13750 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Importin beta-1 subunit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	871	Total	C	N	O	S	1	0	0
			6762	4259	1135	1321	47			
1	B	845	Total	C	N	O	S	0	0	0
			6550	4129	1100	1275	46			

- Molecule 2 is a protein called Snurportin-1.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	C	36	Total	C	N	O	0	0	0
			323	198	70	55			
2	D	23	Total	C	N	O	0	0	0
			115	69	23	23			






• Molecule 2: Snurportin-1

Chain C: 50% 30% 10% 10%



• Molecule 2: Snurportin-1

Chain D: 10% 48% 10% 43%



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	102.98Å 100.70Å 108.68Å 90.00° 109.23° 90.00°	Depositor
Resolution (Å)	40.00 – 3.20 39.61 – 3.20	Depositor EDS
% Data completeness (in resolution range)	(Not available) (40.00-3.20) 91.3 (39.61-3.20)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.98 (at 3.18Å)	Xtriage
Refinement program	CNS	Depositor
R, $R_{free}$	0.303 , 0.328 0.296 , 0.307	Depositor DCC
$R_{free}$ test set	1598 reflections (5.03%)	DCC
Wilson B-factor (Å <sup>2</sup> )	70.4	Xtriage
Anisotropy	0.152	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.28 , 73.8	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Outliers	1 of 31761 reflections (0.003%)	Xtriage
$F_o, F_c$ correlation	0.87	EDS
Total number of atoms	13750	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	83.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.42% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.47	0/6871	0.68	11/9328 (0.1%)
1	B	0.42	0/6651	0.67	19/9023 (0.2%)
2	C	0.50	0/326	0.67	0/428
2	D	0.47	0/114	0.37	0/158
All	All	0.44	0/13962	0.67	30/18937 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	25
1	B	0	48
All	All	0	73

There are no bond length outliers.

The worst 5 of 30 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	396	GLY	C-N-CD	-5.68	108.11	120.60
1	A	779	ASP	CB-CG-OD2	5.28	123.05	118.30
1	A	690	ASP	CB-CG-OD2	5.28	123.05	118.30
1	A	832	ASP	CB-CG-OD2	5.28	123.05	118.30
1	B	337	ASP	CB-CG-OD2	5.27	123.05	118.30

There are no chirality outliers.

5 of 73 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	251	TYR	Peptide
1	A	396	GLY	Peptide

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Group
1	A	498	CYS	Peptide
1	A	622	GLY	Peptide
1	A	681	LEU	Peptide

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6762	0	6764	426	0
1	B	6550	0	6568	655	0
2	C	323	0	337	15	0
2	D	115	0	48	6	0
All	All	13750	0	13717	1047	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 38.

The worst 5 of 1047 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:686:ILE:CG1	1:B:687:PRO:HD3	1.08	1.56
1:B:686:ILE:CG1	1:B:687:PRO:CD	2.02	1.37
1:B:647:MET:CA	1:B:649:ALA:HB2	1.52	1.35
1:B:647:MET:SD	1:B:649:ALA:HB3	1.65	1.35
1:A:130:ILE:CG2	1:A:131:PRO:HD3	1.61	1.30

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.



The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	867/876 (99%)	761 (88%)	97 (11%)	9 (1%)	19	65
1	B	834/876 (95%)	712 (85%)	110 (13%)	12 (1%)	14	57
2	C	32/40 (80%)	27 (84%)	4 (12%)	1 (3%)	5	34
2	D	21/40 (52%)	16 (76%)	5 (24%)	0	100	100
All	All	1754/1832 (96%)	1516 (86%)	216 (12%)	22 (1%)	15	59

5 of 22 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	441	GLU
1	A	560	GLN
1	A	732	LEU
1	B	47	PRO
1	B	72	ILE

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	746/751 (99%)	695 (93%)	51 (7%)	20	59
1	B	722/751 (96%)	648 (90%)	74 (10%)	9	36
2	C	35/39 (90%)	31 (89%)	4 (11%)	7	31
All	All	1503/1541 (98%)	1374 (91%)	129 (9%)	13	46

5 of 129 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	71	ASP
1	B	400	SER
1	B	825	LEU
1	B	79	ARG
1	B	213	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 51 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	759	ASN
1	B	141	ASN
1	B	712	GLN
1	A	856	ASN
1	B	179	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	871/876 (99%)	-0.19	27 (3%) 52 38	11, 51, 141, 200	1 (0%)
1	B	845/876 (96%)	0.61	96 (11%) 7 4	14, 102, 189, 199	0
2	C	36/40 (90%)	-0.13	0 100 100	31, 63, 103, 133	0
2	D	23/40 (57%)	0.70	4 (17%) 2 1	58, 70, 82, 90	0
All	All	1775/1832 (96%)	0.20	127 (7%) 18 10	11, 67, 178, 200	1 (0%)

The worst 5 of 127 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	830	GLY	14.2
1	B	812	ASP	13.9
1	B	623	GLY	10.9
1	B	811	THR	10.9
1	B	800	PHE	10.5

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.