



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 05:24 AM GMT

PDB ID : 2QHO  
Title : Crystal structure of the UBA domain from EDD ubiquitin ligase in complex with ubiquitin  
Authors : Kozlov, G.; Gehring, K.  
Deposited on : 2007-07-02  
Resolution : 1.85 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	1745 (1.86-1.86)
Clashscore	102246	1898 (1.86-1.86)
Ramachandran outliers	100387	1875 (1.86-1.86)
Sidechain outliers	100360	1875 (1.86-1.86)
RSRZ outliers	91569	1747 (1.86-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	76	 76% 16% 5%
1	C	76	 91% 8%
1	E	76	 87% 11%
1	G	76	 84% 11% 5%
2	B	53	 79% 9% 11%

*Continued on next page...*

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	D	53	<div><div>4%</div><div><div></div><div>75%</div><div>9%</div><div>8%</div><div>8%</div></div></div>
2	F	53	<div><div></div><div><div>83%</div><div>9%</div><div>8%</div></div></div>
2	H	53	<div><div></div><div><div>79%</div><div>6%</div><div>6%</div><div>9%</div></div></div>

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 4144 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Ubiquitin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	72	Total	C	N	O	S	0	0	0
			574	362	98	113	1			
1	C	76	Total	C	N	O	S	0	1	0
			607	381	105	120	1			
1	E	75	Total	C	N	O	S	0	0	0
			597	376	104	116	1			
1	G	72	Total	C	N	O	S	0	0	0
			574	362	98	113	1			

- Molecule 2 is a protein called E3 ubiquitin-protein ligase EDD1.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	B	47	Total	C	N	O	0	1	0
			364	226	68	70			
2	D	49	Total	C	N	O	0	0	0
			374	232	70	72			
2	F	49	Total	C	N	O	0	0	0
			380	235	70	75			
2	H	48	Total	C	N	O	0	0	0
			370	230	69	71			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	178	GLY	-	CLONING ARTIFACT	UNP O95071
B	179	SER	-	CLONING ARTIFACT	UNP O95071
D	178	GLY	-	CLONING ARTIFACT	UNP O95071
D	179	SER	-	CLONING ARTIFACT	UNP O95071
F	178	GLY	-	CLONING ARTIFACT	UNP O95071
F	179	SER	-	CLONING ARTIFACT	UNP O95071
H	178	GLY	-	CLONING ARTIFACT	UNP O95071
H	179	SER	-	CLONING ARTIFACT	UNP O95071


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	45	Total 45	O 45	0	0
3	B	37	Total 37	O 37	0	0
3	C	63	Total 63	O 63	0	0
3	D	33	Total 33	O 33	0	0
3	E	50	Total 50	O 50	0	0
3	F	24	Total 24	O 24	0	0
3	G	26	Total 26	O 26	0	0
3	H	26	Total 26	O 26	0	0

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Ubiquitin

Chain A: 





- Molecule 1: Ubiquitin

Chain C: 




- Molecule 1: Ubiquitin

Chain E: 



- Molecule 1: Ubiquitin

Chain G: 




- Molecule 2: E3 ubiquitin-protein ligase EDD1

Chain B: 




- Molecule 2: E3 ubiquitin-protein ligase EDD1

Chain D: 




- Molecule 2: E3 ubiquitin-protein ligase EDD1

Chain F: 83% 9% 8%



- Molecule 2: E3 ubiquitin-protein ligase EDD1

Chain H: 79% 6% 6% 9%



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	33.85Å 59.33Å 246.67Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	33.53 – 1.85 33.53 – 1.85	Depositor EDS
% Data completeness (in resolution range)	92.9 (33.53-1.85) 92.9 (33.53-1.85)	Depositor EDS
$R_{merge}$	0.05	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	6.24 (at 1.85Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, $R_{free}$	0.207 , 0.258 0.205 , 0.256	Depositor DCC
$R_{free}$ test set	2034 reflections (5.26%)	DCC
Wilson B-factor (Å <sup>2</sup> )	15.5	Xtriage
Anisotropy	0.144	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.39 , 47.4	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 40669 reflections	Xtriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	4144	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	15.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.15% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.



## 5 Model quality

### 5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.82	0/580	0.99	3/781 (0.4%)
1	C	0.85	0/617	0.79	0/828
1	E	0.80	0/603	0.96	2/811 (0.2%)
1	G	0.71	0/580	0.77	0/781
2	B	0.91	0/371	0.86	0/503
2	D	0.86	0/375	0.91	0/508
2	F	0.81	0/381	0.73	0/517
2	H	0.79	0/371	0.76	0/503
All	All	0.82	0/3878	0.86	5/5232 (0.1%)

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	54	ARG	NE-CZ-NH1	-10.09	115.26	120.30
1	E	42	ARG	NE-CZ-NH1	-9.74	115.43	120.30
1	A	54	ARG	NE-CZ-NH2	8.59	124.59	120.30
1	E	42	ARG	NE-CZ-NH2	8.45	124.52	120.30
1	A	39	ASP	CB-CG-OD1	5.25	123.03	118.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	574	0	599	11	0
1	C	607	0	631	5	0
1	E	597	0	626	3	0
1	G	574	0	599	3	0
2	B	364	0	389	3	0
2	D	374	0	400	18	0
2	F	380	0	400	6	0
2	H	370	0	397	5	0
3	A	45	0	0	1	0
3	B	37	0	0	1	0
3	C	63	0	0	1	0
3	D	33	0	0	1	0
3	E	50	0	0	0	0
3	F	24	0	0	0	0
3	G	26	0	0	0	0
3	H	26	0	0	2	0
All	All	4144	0	4041	48	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (48) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:226:ARG:HH21	2:D:226:ARG:HG3	1.30	0.96
2:D:207:ARG:HH11	2:D:210:GLN:HE22	1.04	0.93
2:D:207:ARG:HD3	2:D:210:GLN:NE2	1.85	0.91
2:D:207:ARG:NH1	2:D:210:GLN:HE22	1.79	0.78
1:A:51:GLU:OE1	1:A:54:ARG:HD2	1.84	0.76
2:D:212:THR:HG21	2:D:219:ALA:HB2	1.71	0.72
2:D:207:ARG:HH11	2:D:210:GLN:NE2	1.83	0.70
2:D:226:ARG:HH21	2:D:226:ARG:CG	2.03	0.70
1:C:16:GLU:O	1:C:29:LYS:HE2	1.91	0.70
2:H:212:THR:HG21	2:H:219:ALA:HB2	1.73	0.70
2:D:207:ARG:HD3	2:D:210:GLN:HE21	1.61	0.65
2:D:200:LYS:HB2	2:D:200:LYS:NZ	2.13	0.64
1:A:11:LYS:HE3	1:A:13:ILE:HD11	1.81	0.62
2:D:207:ARG:HD3	2:D:210:GLN:HE22	1.64	0.62
2:B:191:SER:HA	2:B:194:GLN:HE21	1.66	0.61
1:A:51:GLU:HB2	1:A:54:ARG:HG3	1.83	0.60
2:D:208:GLU:O	2:D:212:THR:HB	2.01	0.60
1:A:72:ARG:HD2	3:A:104:HOH:O	2.02	0.58

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:G:24:GLU:HG2	1:G:52:ASP:HB3	1.84	0.58
2:D:207:ARG:O	2:D:211:ARG:HG2	2.05	0.56
1:C:13:ILE:CD1	1:C:34:GLU:HG3	2.36	0.55
2:D:208:GLU:HA	2:D:211:ARG:HG3	1.90	0.54
2:H:208:GLU:O	2:H:212:THR:HB	2.10	0.52
1:A:62:GLN:HE22	2:F:199:GLY:H	1.59	0.50
1:A:63:LYS:HG2	1:A:64:GLU:OE2	2.13	0.49
2:D:207:ARG:CD	2:D:210:GLN:NE2	2.70	0.48
2:H:217:ASN:ND2	3:H:233:HOH:O	2.45	0.48
2:D:207:ARG:CD	2:D:210:GLN:HE22	2.26	0.48
1:A:62:GLN:NE2	2:F:199:GLY:H	2.13	0.47
1:E:22:THR:H	1:E:25:ASN:ND2	2.13	0.47
1:E:61:ILE:HD13	1:E:67:LEU:HD11	1.96	0.47
1:A:57:SER:HA	2:F:198:GLN:HE22	1.81	0.46
1:A:19:PRO:HB3	2:F:198:GLN:CD	2.36	0.46
2:H:207:ARG:HD3	3:H:243:HOH:O	2.17	0.45
1:G:22:THR:O	1:G:25:ASN:HB2	2.17	0.45
1:C:13:ILE:HD13	1:C:34:GLU:HG3	2.00	0.44
1:C:13:ILE:HD11	1:C:34:GLU:HG3	1.99	0.44
1:G:15:LEU:HD11	1:G:30:ILE:HG13	2.00	0.44
1:A:8:LEU:HD12	2:B:217:ASN:HD22	1.83	0.44
2:D:207:ARG:NH1	2:D:210:GLN:NE2	2.55	0.43
2:D:207:ARG:HG2	3:D:257:HOH:O	2.18	0.43
1:E:46:ALA:HA	2:F:198:GLN:NE2	2.34	0.43
2:D:226:ARG:HG3	2:D:226:ARG:NH2	2.12	0.42
1:A:42:ARG:HD2	1:A:72:ARG:HB3	2.02	0.42
2:H:211:ARG:HB3	2:H:211:ARG:CZ	2.50	0.41
2:F:207:ARG:O	2:F:211:ARG:HG3	2.20	0.41
1:C:62:GLN:HB2	3:C:113:HOH:O	2.19	0.41
2:B:213:ASN:ND2	3:B:264:HOH:O	2.47	0.41

There are no symmetry-related clashes.

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	70/76 (92%)	70 (100%)	0	0	100	100
1	C	75/76 (99%)	74 (99%)	1 (1%)	0	100	100
1	E	73/76 (96%)	73 (100%)	0	0	100	100
1	G	70/76 (92%)	70 (100%)	0	0	100	100
2	B	46/53 (87%)	46 (100%)	0	0	100	100
2	D	47/53 (89%)	47 (100%)	0	0	100	100
2	F	47/53 (89%)	47 (100%)	0	0	100	100
2	H	46/53 (87%)	46 (100%)	0	0	100	100
All	All	474/516 (92%)	473 (100%)	1 (0%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	66/68 (97%)	64 (97%)	2 (3%)	48	29
1	C	69/68 (102%)	66 (96%)	3 (4%)	35	16
1	E	68/68 (100%)	64 (94%)	4 (6%)	24	8
1	G	66/68 (97%)	64 (97%)	2 (3%)	48	29
2	B	44/48 (92%)	43 (98%)	1 (2%)	58	41
2	D	44/48 (92%)	39 (89%)	5 (11%)	7	1
2	F	45/48 (94%)	44 (98%)	1 (2%)	60	43
2	H	44/48 (92%)	41 (93%)	3 (7%)	20	5
All	All	446/464 (96%)	425 (95%)	21 (5%)	32	13

All (21) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	8	LEU
1	A	56	LEU
2	B	184	VAL
1	C	8	LEU
1	C	62	GLN
1	C	71	LEU
2	D	200	LYS
2	D	201	SER
2	D	211	ARG
2	D	212	THR
2	D	226	ARG
1	E	6	LYS
1	E	9	THR
1	E	67	LEU
1	E	71	LEU
2	F	228	ASP
1	G	8	LEU
1	G	16	GLU
2	H	207	ARG
2	H	211	ARG
2	H	212	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (26) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	2	GLN
1	A	25	ASN
1	A	60	ASN
1	A	62	GLN
2	B	194	GLN
2	B	210	GLN
2	B	213	ASN
2	B	217	ASN
1	C	2	GLN
1	C	60	ASN
1	C	62	GLN
2	D	210	GLN
2	D	213	ASN
1	E	2	GLN
1	E	25	ASN
1	E	31	GLN
1	E	62	GLN
2	F	194	GLN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
2	F	198	GLN
2	F	210	GLN
2	F	213	ASN
2	F	217	ASN
1	G	31	GLN
1	G	62	GLN
2	H	210	GLN
2	H	217	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

### 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	72/76 (94%)	-0.21	0 100 100	8, 13, 19, 24	0
1	C	76/76 (100%)	-0.35	0 100 100	8, 13, 22, 28	0
1	E	75/76 (98%)	-0.21	1 (1%) 79 79	10, 15, 23, 31	0
1	G	72/76 (94%)	0.16	0 100 100	12, 21, 30, 35	0
2	B	47/53 (88%)	-0.41	0 100 100	7, 11, 20, 26	0
2	D	49/53 (92%)	-0.21	2 (4%) 41 38	6, 11, 25, 29	0
2	F	49/53 (92%)	-0.27	0 100 100	7, 12, 24, 35	0
2	H	48/53 (90%)	-0.11	0 100 100	9, 16, 26, 35	0
All	All	488/516 (94%)	-0.19	3 (0%) 90 90	6, 14, 27, 35	0

All (3) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	D	178	GLY	3.3
2	D	199	GLY	2.3
1	E	13	ILE	2.2

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.