



# wwPDB X-ray Structure Validation Summary Report ⓘ

Jan 31, 2016 – 09:51 PM GMT

PDB ID : 1QWM  
Title : Structure of Helicobacter pylori catalase with formic acid bound  
Authors : Loewen, P.C.; Carpena, X.; Perez-Luque, R.; Rovira, C.; Haas, R.; Odenbreit, S.; Nicholls, P.; Fita, I.  
Deposited on : 2003-09-02  
Resolution : 1.60 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*



The reported resolution of this entry is 1.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	2475 (1.60-1.60)
Clashscore	102246	2732 (1.60-1.60)
Ramachandran outliers	100387	2654 (1.60-1.60)
Sidechain outliers	100360	2653 (1.60-1.60)
RSRZ outliers	91569	2479 (1.60-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	505	 13% 85% 11% ..
1	B	505	 10% 85% 11% ..

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	AZI	A	2600	-	-	-	X
4	FMT	A	1701	-	-	-	X
4	FMT	A	1713	-	-	-	X
4	FMT	A	1714	-	-	X	-
4	FMT	A	1732	-	-	X	-
4	FMT	A	1733	-	-	-	X
4	FMT	A	1734	-	-	X	X
4	FMT	A	1735	-	-	X	-
4	FMT	B	1702	-	-	-	X
4	FMT	B	1703	-	-	-	X
4	FMT	B	1705	-	-	-	X
4	FMT	B	1712	-	-	X	-
4	FMT	B	1721	-	-	X	-
4	FMT	B	1729	-	-	X	-
4	FMT	B	1731	-	-	X	X
4	FMT	B	1736	-	-	-	X

## 2 Entry composition [i](#)


There are 5 unique types of molecules in this entry. The entry contains 9209 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called KatA catalase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	490	Total	C	N	O	S	0	9	0
			4041	2580	704	743	14			
1	B	490	Total	C	N	O	S	0	9	0
			4038	2578	701	745	14			

- Molecule 2 is AZIDE ION (three-letter code: AZI) (formula: N<sub>3</sub>).




Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	N	0	0
			3	3		

- Molecule 3 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula: C<sub>34</sub>H<sub>32</sub>FeN<sub>4</sub>O<sub>4</sub>).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		
3	B	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		

- Molecule 4 is FORMIC ACID (three-letter code: FMT) (formula: CH<sub>2</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			3	1	2		
4	B	1	Total	C	O	0	0
			3	1	2		

Continued on next page...

*Continued from previous page...*

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total 3	C 1	O 2	0	0
4	A	1	Total 3	C 1	O 2	0	0
4	B	1	Total 3	C 1	O 2	0	0
4	A	1	Total 3	C 1	O 2	0	0
4	B	1	Total 3	C 1	O 2	0	0
4	B	1	Total 3	C 1	O 2	0	0
4	A	1	Total 3	C 1	O 2	0	0
4	A	1	Total 3	C 1	O 2	0	0
4	A	1	Total 3	C 1	O 2	0	0
4	B	1	Total 3	C 1	O 2	0	0
4	A	1	Total 3	C 1	O 2	0	0
4	A	1	Total 3	C 1	O 2	0	0
4	A	1	Total 3	C 1	O 2	0	0
4	A	1	Total 3	C 1	O 2	0	0
4	B	1	Total 3	C 1	O 2	0	0
4	A	1	Total 3	C 1	O 2	0	0
4	A	1	Total 3	C 1	O 2	0	0
4	B	1	Total 3	C 1	O 2	0	0
4	B	1	Total 3	C 1	O 2	0	0
4	A	1	Total 3	C 1	O 2	0	0
4	A	1	Total 3	C 1	O 2	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			3	1	2		
4	A	1	Total	C	O	0	0
			3	1	2		
4	B	1	Total	C	O	0	0
			3	1	2		
4	B	1	Total	C	O	0	0
			3	1	2		
4	B	1	Total	C	O	0	0
			3	1	2		
4	B	1	Total	C	O	0	0
			3	1	2		
4	A	1	Total	C	O	0	0
			3	1	2		
4	B	1	Total	C	O	0	0
			3	1	2		
4	A	1	Total	C	O	0	0
			3	1	2		
4	A	1	Total	C	O	0	0
			3	1	2		
4	A	1	Total	C	O	0	0
			3	1	2		
4	A	1	Total	C	O	0	0
			3	1	2		
4	B	1	Total	C	O	0	0
			3	1	2		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	442	Total	O	0	0
			442	442		
5	B	491	Total	O	0	0
			491	491		





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 2 <sub>1</sub> 2 <sub>1</sub> 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	64.76Å 154.96Å 96.16Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.88 – 1.60 29.87 – 1.60	Depositor EDS
% Data completeness (in resolution range)	100.0 (29.88-1.60) 97.4 (29.87-1.60)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.05	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.41 (at 1.60Å)	Xtriage
Refinement program	REFMAC 5.1.24	Depositor
R, $R_{free}$	0.194 , 0.227 0.204 , 0.237	Depositor DCC
$R_{free}$ test set	6282 reflections (5.30%)	DCC
Wilson B-factor (Å <sup>2</sup> )	12.6	Xtriage
Anisotropy	0.076	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.52 , 60.8	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	1 of 124794 reflections (0.001%)	Xtriage
$F_o, F_c$ correlation	0.92	EDS
Total number of atoms	9209	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	15.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.59% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality ⓘ

### 5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: AZI, FMT, HEM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.79	0/4215	0.89	15/5705 (0.3%)
1	B	0.82	0/4214	0.93	16/5704 (0.3%)
All	All	0.80	0/8429	0.91	31/11409 (0.3%)

There are no bond length outliers.

The worst 5 of 31 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	109	ASP	CB-CG-OD2	7.19	124.77	118.30
1	A	32	LEU	CA-CB-CG	-7.04	99.11	115.30
1	B	184	ARG	NE-CZ-NH1	6.38	123.49	120.30
1	B	18	ASP	CB-CG-OD2	6.35	124.01	118.30
1	B	399	ASP	CB-CG-OD2	6.30	123.97	118.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4041	0	3865	35	1
1	B	4038	0	3852	44	0
2	A	3	0	0	1	0
3	A	43	0	30	1	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	43	0	30	5	0
4	A	63	0	21	18	0
4	B	45	0	15	16	1
5	A	442	0	0	9	2
5	B	491	0	0	8	3
All	All	9209	0	7813	85	4

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 85 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:B:1705:FMT:H	5:B:2015:HOH:O	1.54	1.07
4:A:1714:FMT:O1	5:A:2694:HOH:O	1.83	0.94
1:B:353:GLN:HE22	1:B:374:ASN:H	1.30	0.79
1:A:130:ASN:O	4:A:1714:FMT:H	1.84	0.78
1:B:184:ARG:HG2	4:B:1729:FMT:C	2.15	0.77

All (4) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:154:GLN:NE2	5:A:2849:HOH:O[2_665]	1.45	0.75
5:A:2930:HOH:O	5:B:2217:HOH:O[1_554]	1.91	0.29
4:B:1721:FMT:O2	5:B:2177:HOH:O[2_665]	1.94	0.26
5:B:2009:HOH:O	5:B:2128:HOH:O[2_665]	2.17	0.03

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	497/505 (98%)	478 (96%)	19 (4%)	0	100	100
1	B	497/505 (98%)	482 (97%)	14 (3%)	1 (0%)	52	28
All	All	994/1010 (98%)	960 (97%)	33 (3%)	1 (0%)	56	31

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	369	ASP

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	436/442 (99%)	419 (96%)	17 (4%)	39	13
1	B	436/442 (99%)	425 (98%)	11 (2%)	55	26
All	All	872/884 (99%)	844 (97%)	28 (3%)	50	18

5 of 28 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	377	TYR
1	A	475	TYR
1	B	414	TRP
1	A	414	TRP
1	A	418	TYR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	353	GLN
1	A	364	HIS
1	B	287	GLN
1	B	353	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

39 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
4	FMT	A	1701	3	0,2,2	0.00	-	0,1,1	0.00	-
4	FMT	A	1704	-	0,2,2	0.00	-	0,1,1	0.00	-
4	FMT	A	1706	-	0,2,2	0.00	-	0,1,1	0.00	-
4	FMT	A	1709	-	0,2,2	0.00	-	0,1,1	0.00	-
4	FMT	A	1710	1	0,2,2	0.00	-	0,1,1	0.00	-
4	FMT	A	1711	-	0,2,2	0.00	-	0,1,1	0.00	-
4	FMT	A	1713	-	0,2,2	0.00	-	0,1,1	0.00	-
4	FMT	A	1714	-	0,2,2	0.00	-	0,1,1	0.00	-
4	FMT	A	1715	-	0,2,2	0.00	-	0,1,1	0.00	-
4	FMT	A	1716	-	0,2,2	0.00	-	0,1,1	0.00	-
4	FMT	A	1718	-	0,2,2	0.00	-	0,1,1	0.00	-
4	FMT	A	1719	-	0,2,2	0.00	-	0,1,1	0.00	-
4	FMT	A	1722	-	0,2,2	0.00	-	0,1,1	0.00	-
4	FMT	A	1723	-	0,2,2	0.00	-	0,1,1	0.00	-
4	FMT	A	1724	-	0,2,2	0.00	-	0,1,1	0.00	-
4	FMT	A	1725	-	0,2,2	0.00	-	0,1,1	0.00	-
4	FMT	A	1730	-	0,2,2	0.00	-	0,1,1	0.00	-
4	FMT	A	1732	-	0,2,2	0.00	-	0,1,1	0.00	-

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	FMT	A	1733	-	0,2,2	0.00	-	0,1,1	0.00	-
4	FMT	A	1734	-	0,2,2	0.00	-	0,1,1	0.00	-
4	FMT	A	1735	2	0,2,2	0.00	-	0,1,1	0.00	-
2	AZI	A	2600	4	0,2,2	0.00	-	0,1,1	0.00	-
3	HEM	A	550	1,4	30,50,50	2.54	9 (30%)	24,82,82	2.52	13 (54%)
4	FMT	B	1702	-	0,2,2	0.00	-	0,1,1	0.00	-
4	FMT	B	1703	-	0,2,2	0.00	-	0,1,1	0.00	-
4	FMT	B	1705	-	0,2,2	0.00	-	0,1,1	0.00	-
4	FMT	B	1707	-	0,2,2	0.00	-	0,1,1	0.00	-
4	FMT	B	1708	-	0,2,2	0.00	-	0,1,1	0.00	-
4	FMT	B	1712	-	0,2,2	0.00	-	0,1,1	0.00	-
4	FMT	B	1717	-	0,2,2	0.00	-	0,1,1	0.00	-
4	FMT	B	1720	-	0,2,2	0.00	-	0,1,1	0.00	-
4	FMT	B	1721	-	0,2,2	0.00	-	0,1,1	0.00	-
4	FMT	B	1726	-	0,2,2	0.00	-	0,1,1	0.00	-
4	FMT	B	1727	-	0,2,2	0.00	-	0,1,1	0.00	-
4	FMT	B	1728	-	0,2,2	0.00	-	0,1,1	0.00	-
4	FMT	B	1729	-	0,2,2	0.00	-	0,1,1	0.00	-
4	FMT	B	1731	-	0,2,2	0.00	-	0,1,1	0.00	-
4	FMT	B	1736	-	0,2,2	0.00	-	0,1,1	0.00	-
3	HEM	B	550	1	30,50,50	2.15	7 (23%)	24,82,82	2.73	13 (54%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	FMT	A	1701	3	-	0/0/0/0	0/0/0/0
4	FMT	A	1704	-	-	0/0/0/0	0/0/0/0
4	FMT	A	1706	-	-	0/0/0/0	0/0/0/0
4	FMT	A	1709	-	-	0/0/0/0	0/0/0/0
4	FMT	A	1710	1	-	0/0/0/0	0/0/0/0
4	FMT	A	1711	-	-	0/0/0/0	0/0/0/0
4	FMT	A	1713	-	-	0/0/0/0	0/0/0/0
4	FMT	A	1714	-	-	0/0/0/0	0/0/0/0
4	FMT	A	1715	-	-	0/0/0/0	0/0/0/0
4	FMT	A	1716	-	-	0/0/0/0	0/0/0/0
4	FMT	A	1718	-	-	0/0/0/0	0/0/0/0
4	FMT	A	1719	-	-	0/0/0/0	0/0/0/0
4	FMT	A	1722	-	-	0/0/0/0	0/0/0/0
4	FMT	A	1723	-	-	0/0/0/0	0/0/0/0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	FMT	A	1724	-	-	0/0/0/0	0/0/0/0
4	FMT	A	1725	-	-	0/0/0/0	0/0/0/0
4	FMT	A	1730	-	-	0/0/0/0	0/0/0/0
4	FMT	A	1732	-	-	0/0/0/0	0/0/0/0
4	FMT	A	1733	-	-	0/0/0/0	0/0/0/0
4	FMT	A	1734	-	-	0/0/0/0	0/0/0/0
4	FMT	A	1735	2	-	0/0/0/0	0/0/0/0
2	AZI	A	2600	4	-	0/0/0/0	0/0/0/0
3	HEM	A	550	1,4	-	0/10/54/54	0/0/8/8
4	FMT	B	1702	-	-	0/0/0/0	0/0/0/0
4	FMT	B	1703	-	-	0/0/0/0	0/0/0/0
4	FMT	B	1705	-	-	0/0/0/0	0/0/0/0
4	FMT	B	1707	-	-	0/0/0/0	0/0/0/0
4	FMT	B	1708	-	-	0/0/0/0	0/0/0/0
4	FMT	B	1712	-	-	0/0/0/0	0/0/0/0
4	FMT	B	1717	-	-	0/0/0/0	0/0/0/0
4	FMT	B	1720	-	-	0/0/0/0	0/0/0/0
4	FMT	B	1721	-	-	0/0/0/0	0/0/0/0
4	FMT	B	1726	-	-	0/0/0/0	0/0/0/0
4	FMT	B	1727	-	-	0/0/0/0	0/0/0/0
4	FMT	B	1728	-	-	0/0/0/0	0/0/0/0
4	FMT	B	1729	-	-	0/0/0/0	0/0/0/0
4	FMT	B	1731	-	-	0/0/0/0	0/0/0/0
4	FMT	B	1736	-	-	0/0/0/0	0/0/0/0
3	HEM	B	550	1	-	0/10/54/54	0/0/8/8

The worst 5 of 16 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	550	HEM	C3B-C4B	-8.42	1.44	1.51
3	B	550	HEM	C3B-C4B	-6.90	1.45	1.51
3	A	550	HEM	C3D-C4D	-6.43	1.43	1.51
3	B	550	HEM	C3D-C4D	-4.98	1.45	1.51
3	B	550	HEM	C2C-C1C	-3.50	1.45	1.52

The worst 5 of 26 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	550	HEM	CAA-C2A-C1A	-4.33	122.31	127.01
3	B	550	HEM	C3B-CAB-CBB	-3.65	118.86	124.46
3	B	550	HEM	CBA-CAA-C2A	-3.49	106.28	112.53
3	A	550	HEM	C3B-CAB-CBB	-2.81	120.14	124.46

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	550	HEM	CAA-C2A-C1A	-2.66	124.11	127.01

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

19 monomers are involved in 41 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	1701	FMT	1	0
4	A	1710	FMT	1	0
4	A	1714	FMT	4	0
4	A	1718	FMT	1	0
4	A	1730	FMT	1	0
4	A	1732	FMT	3	0
4	A	1734	FMT	4	0
4	A	1735	FMT	4	0
2	A	2600	AZI	1	0
3	A	550	HEM	1	0
4	B	1702	FMT	1	0
4	B	1705	FMT	1	0
4	B	1712	FMT	5	0
4	B	1720	FMT	1	0
4	B	1721	FMT	1	1
4	B	1728	FMT	1	0
4	B	1729	FMT	3	0
4	B	1731	FMT	3	0
3	B	550	HEM	5	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	490/505 (97%)	1.00	65 (13%) 4 4	5, 14, 25, 48	2 (0%)
1	B	490/505 (97%)	0.87	52 (10%) 8 7	5, 12, 23, 47	1 (0%)
All	All	980/1010 (97%)	0.93	117 (11%) 6 5	5, 13, 24, 48	3 (0%)

The worst 5 of 117 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	489	MET	9.1
1	B	489	MET	8.8
1	A	488	LYS	6.8
1	A	266	PRO	5.8
1	B	268	GLU	5.6

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
4	FMT	B	1736	3/3	0.62	0.41	14.47	29,29,31,31	0
4	FMT	B	1702	3/3	0.71	0.26	8.20	38,38,39,41	0
4	FMT	A	1701	3/3	0.70	0.23	6.28	26,26,28,29	0
4	FMT	B	1703	3/3	0.65	0.22	4.79	27,27,30,32	0
4	FMT	B	1731	3/3	0.85	0.21	4.66	13,13,21,27	0
4	FMT	B	1705	3/3	0.88	0.18	4.35	32,32,34,36	0
4	FMT	A	1734	3/3	0.72	0.29	4.08	26,26,28,29	0
2	AZI	A	2600	3/3	0.87	0.16	3.18	11,11,18,22	0
4	FMT	A	1713	3/3	0.87	0.16	2.54	24,24,26,29	0
4	FMT	A	1733	3/3	0.87	0.15	2.24	27,27,27,28	0
4	FMT	B	1721	3/3	0.76	0.20	1.94	38,38,39,40	0
4	FMT	B	1720	3/3	0.82	0.18	1.56	30,30,31,31	0
4	FMT	A	1732	3/3	0.93	0.15	1.35	15,15,23,29	0
4	FMT	B	1729	3/3	0.79	0.18	1.26	45,45,45,45	0
4	FMT	A	1719	3/3	0.82	0.19	0.97	27,27,28,28	0
4	FMT	B	1727	3/3	0.79	0.17	0.76	36,36,37,38	0
4	FMT	B	1712	3/3	0.88	0.16	0.72	24,24,24,26	0
3	HEM	A	550	43/43	0.93	0.15	0.71	7,10,14,20	0
3	HEM	B	550	43/43	0.94	0.13	0.47	5,9,15,21	0
4	FMT	A	1716	3/3	0.88	0.12	0.40	18,18,20,23	0
4	FMT	A	1714	3/3	0.95	0.12	0.24	17,17,22,27	0
4	FMT	B	1717	3/3	0.70	0.18	-0.00	34,34,36,36	0
4	FMT	A	1735	3/3	0.82	0.11	-0.06	22,22,25,28	0
4	FMT	A	1711	3/3	0.91	0.13	-0.07	32,32,33,33	0
4	FMT	A	1710	3/3	0.92	0.09	-0.82	32,32,32,32	0
4	FMT	A	1709	3/3	0.86	0.10	-1.43	39,39,39,39	0
4	FMT	A	1715	3/3	0.71	0.25	-	49,49,49,49	0
4	FMT	B	1707	3/3	0.68	0.18	-	38,38,39,39	0
4	FMT	A	1718	3/3	0.91	0.09	-	28,28,30,30	0
4	FMT	A	1722	3/3	0.84	0.14	-	39,39,39,40	0
4	FMT	B	1728	3/3	0.63	0.16	-	49,49,50,50	0
4	FMT	A	1730	3/3	0.63	0.16	-	38,38,39,39	0
4	FMT	A	1706	3/3	0.84	0.13	-	42,42,42,42	0
4	FMT	A	1704	3/3	0.88	0.20	-	42,42,43,44	0
4	FMT	A	1725	3/3	0.61	0.12	-	41,41,41,42	0
4	FMT	A	1724	3/3	0.53	0.21	-	48,48,48,48	0
4	FMT	B	1726	3/3	0.88	0.11	-	54,54,55,55	0
4	FMT	A	1723	3/3	0.80	0.17	-	54,54,54,54	0
4	FMT	B	1708	3/3	0.96	0.13	-	32,32,33,33	0

## 6.5 Other polymers ⓘ

There are no such residues in this entry.