



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 05:32 AM GMT

PDB ID : 2R6W
Title : Estrogen receptor alpha ligand-binding domain complexed to a SERM
Authors : Wang, Y.
Deposited on : 2007-09-06
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	6249 (2.00-2.00)
Clashscore	102246	7340 (2.00-2.00)
Ramachandran outliers	100387	7248 (2.00-2.00)
Sidechain outliers	100360	7247 (2.00-2.00)
RSRZ outliers	91569	6262 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	248	<div> <div>12%</div> <div>72%</div> <div>18%</div> <div>8%</div> </div>
1	B	248	<div> <div>8%</div> <div>70%</div> <div>19%</div> <div>10%</div> </div>

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 3799 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Estrogen receptor.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	229	Total	C	N	O	S	0	2	0
			1842	1185	313	329	15			
1	B	223	Total	C	N	O	S	0	1	0
			1783	1149	301	318	15			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	381	SER	CYS	ENGINEERED	UNP P03372
A	417	SER	CYS	ENGINEERED	UNP P03372
A	530	SER	CYS	ENGINEERED	UNP P03372
B	381	SER	CYS	ENGINEERED	UNP P03372
B	417	SER	CYS	ENGINEERED	UNP P03372
B	530	SER	CYS	ENGINEERED	UNP P03372

- Molecule 2 is [6-HYDROXY-2-(4-HYDROXYPHENYL)-1-BENZOTHIEN-3-YL]{4-[2-(4-METHYLPIPERIDIN-1-YL)ETHOXY]PHENYL}METHANONE (three-letter code: LLB) (formula: C₂₉H₂₉NO₄S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	S	0	0
			35	29	1	4	1		
2	B	1	Total	C	N	O	S	0	0
			35	29	1	4	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	46	Total	O	0	0
			46	46		
3	B	58	Total	O	0	0
			58	58		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Estrogen receptor



• Molecule 1: Estrogen receptor



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	56.47Å 102.76Å 172.95Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	22.27 – 2.00 22.27 – 1.93	Depositor EDS
% Data completeness (in resolution range)	92.2 (22.27-2.00) 87.0 (22.27-1.93)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.81 (at 1.93Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.208 , 0.257 0.209 , 0.258	Depositor DCC
R_{free} test set	2544 reflections (8.71%)	DCC
Wilson B-factor (Å ²)	29.7	Xtriage
Anisotropy	0.673	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.40 , 57.8	EDS
Estimated twinning fraction	0.021 for 1/2*h-1/2*k,-3/2*h-1/2*k,-l 0.025 for 1/2*h+1/2*k,3/2*h-1/2*k,-l	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Outliers	0 of 33053 reflections	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	3799	wwPDB-VP
Average B, all atoms (Å ²)	42.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.86% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: LLB

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.74	0/1876	0.77	0/2536
1	B	0.76	0/1817	0.83	1/2452 (0.0%)
All	All	0.75	0/3693	0.80	1/4988 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	469	LEU	CA-CB-CG	5.05	126.92	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1842	0	1879	44	0
1	B	1783	0	1829	62	0
2	A	35	0	28	3	0
2	B	35	0	28	6	0
3	A	46	0	0	2	0
3	B	58	0	0	12	0
All	All	3799	0	3764	89	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

All (89) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:418:VAL:HG12	1:B:419:GLU:H	1.24	1.01
1:B:401:LYS:HD3	1:B:409:LEU:HD21	1.46	0.96
1:A:519:ASN:HD22	1:B:519:ASN:HD22	1.08	0.91
1:A:513[B]:HIS:CD2	1:B:459:TYR:CD2	2.60	0.89
1:B:392:VAL:HB	3:B:608:HOH:O	1.74	0.86
1:A:421:MET:HB3	1:A:524:HIS:CE1	2.10	0.86
1:B:392:VAL:CB	3:B:608:HOH:O	2.24	0.85
1:B:392:VAL:CG1	3:B:608:HOH:O	2.24	0.85
1:A:513[B]:HIS:CD2	1:B:459:TYR:HD2	1.94	0.84
1:B:392:VAL:HG12	3:B:608:HOH:O	1.77	0.83
1:A:506:GLN:HG2	3:B:560:HOH:O	1.84	0.75
1:B:418:VAL:HG12	1:B:419:GLU:N	2.02	0.75
1:A:513[B]:HIS:CD2	1:B:459:TYR:CE2	2.79	0.71
1:A:519:ASN:ND2	1:B:519:ASN:HD22	1.88	0.70
1:B:379:LEU:CD1	1:B:544:LEU:HD11	2.23	0.69
1:B:418:VAL:CG1	1:B:419:GLU:H	2.03	0.68
1:A:310:LEU:O	1:A:481:LYS:NZ	2.26	0.68
1:A:519:ASN:HD22	1:B:519:ASN:ND2	1.89	0.67
1:A:459:TYR:CD2	1:B:513[A]:HIS:CD2	2.83	0.65
1:B:377:HIS:CE1	1:B:460:THR:HG23	2.31	0.65
1:B:434:ARG:NH1	3:B:602:HOH:O	2.29	0.64
1:B:377:HIS:NE2	1:B:460:THR:HG23	2.12	0.63
1:A:513[B]:HIS:CG	1:B:459:TYR:HD2	2.17	0.63
1:B:310:LEU:O	1:B:481:LYS:HE3	2.00	0.62
1:B:519:ASN:O	1:B:523:GLU:HG3	2.02	0.60
1:A:501:HIS:HB3	3:A:559:HOH:O	2.02	0.59
1:B:496:THR:O	1:B:500:GLN:HG3	2.02	0.59
1:A:434:ARG:HD3	1:B:459:TYR:HE1	1.68	0.59
1:B:316:VAL:HG21	1:B:489:LEU:HD21	1.85	0.59
1:A:380:GLU:HG3	1:A:536:LEU:HD23	1.86	0.57
1:A:434:ARG:HD3	1:B:459:TYR:CE1	2.39	0.57
1:B:308:LEU:O	1:B:481:LYS:HE2	2.03	0.57
1:A:459:TYR:OH	1:B:434:ARG:HD3	2.03	0.56
1:B:377:HIS:HE1	1:B:457:GLY:O	1.88	0.56
1:A:459:TYR:HD2	1:B:513[A]:HIS:CD2	2.23	0.55
1:A:368:VAL:HG22	3:A:570:HOH:O	2.07	0.55
1:A:526:TYR:CD1	1:A:526:TYR:C	2.79	0.55

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:347:THR:HG23	2:A:1:LLB:H281	1.89	0.54
1:B:501:HIS:HB2	3:B:572:HOH:O	2.07	0.54
1:B:347:THR:HG23	2:B:1:LLB:H282	1.90	0.54
1:B:377:HIS:CE1	1:B:457:GLY:O	2.61	0.53
1:B:316:VAL:HG23	1:B:485:THR:CG2	2.37	0.53
1:A:424:ILE:HG13	2:A:1:LLB:O59	2.08	0.53
1:B:524:HIS:HE1	2:B:1:LLB:H53	1.74	0.53
1:B:341:SER:O	1:B:345:LEU:HG	2.09	0.53
1:A:539:LEU:HB3	1:A:543:MET:CE	2.40	0.52
1:A:513[B]:HIS:NE2	1:B:459:TYR:CD2	2.79	0.51
1:B:434:ARG:HG2	1:B:510:ILE:HD11	1.93	0.51
1:A:412:ARG:HA	1:A:425:PHE:CD1	2.47	0.49
1:B:403:LEU:HD13	1:B:409:LEU:HD12	1.94	0.49
1:B:354:LEU:CD2	2:B:1:LLB:H652	2.42	0.49
1:A:455:ASN:O	1:A:458:VAL:HG12	2.13	0.49
1:B:401:LYS:HD3	1:B:409:LEU:CD2	2.33	0.49
1:A:434:ARG:HA	1:A:437:MET:HE2	1.95	0.47
1:A:539:LEU:O	1:A:543:MET:HG3	2.15	0.47
1:B:392:VAL:HG13	1:B:432:SER:HA	1.97	0.47
1:B:535:PRO:N	3:B:603:HOH:O	2.48	0.47
1:B:524:HIS:HE1	2:B:1:LLB:C53	2.27	0.46
1:A:513[B]:HIS:CD2	1:B:459:TYR:HE2	2.33	0.45
1:A:371:THR:HG21	1:A:467:LYS:HE2	1.98	0.45
1:A:513[B]:HIS:NE2	1:B:459:TYR:HD2	2.15	0.45
1:B:310:LEU:O	1:B:481:LYS:CE	2.65	0.45
1:B:374:ASP:OD2	1:B:471:GLU:OE1	2.35	0.45
1:B:376:VAL:HG21	1:B:541:LEU:HD13	1.99	0.44
1:B:477:ARG:HD3	3:B:558:HOH:O	2.18	0.44
1:A:459:TYR:HD2	1:B:513[A]:HIS:CG	2.36	0.44
1:A:392:VAL:HG13	1:A:432:SER:HA	2.00	0.43
1:B:392:VAL:HG13	1:B:432:SER:CA	2.49	0.43
1:B:535:PRO:CA	3:B:603:HOH:O	2.67	0.43
1:A:392:VAL:HG12	1:A:435:PHE:CD2	2.54	0.43
1:B:413:ASN:HA	1:B:416:LYS:HE3	2.01	0.43
1:A:539:LEU:HD13	2:A:1:LLB:H71	2.01	0.43
1:A:522:MET:HE3	1:A:522:MET:HA	2.00	0.42
1:A:379:LEU:HD12	1:A:544:LEU:HD11	2.01	0.42
1:B:379:LEU:HD12	1:B:544:LEU:HD11	2.00	0.42
1:A:539:LEU:HB3	1:A:543:MET:HE2	2.01	0.42
1:B:437:MET:HE3	3:B:596:HOH:O	2.18	0.42
1:A:424:ILE:H	1:A:424:ILE:HG12	1.64	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:535:PRO:HA	3:B:603:HOH:O	2.19	0.41
1:A:342:MET:HE2	1:A:418:VAL:HG21	2.02	0.41
1:A:315:MET:HE2	1:A:319:LEU:HD11	2.01	0.41
1:A:518:SER:O	1:A:522:MET:HG2	2.20	0.41
1:A:403:LEU:HD11	1:A:406:PRO:HA	2.03	0.41
1:A:459:TYR:OH	1:B:434:ARG:NH1	2.34	0.41
1:A:377[A]:HIS:HE1	1:A:457:GLY:O	2.04	0.41
1:B:524:HIS:CE1	2:B:1:LLB:O59	2.74	0.40
1:B:524:HIS:CE1	2:B:1:LLB:H53	2.54	0.40
1:B:520:LYS:HA	1:B:520:LYS:HD2	1.93	0.40
1:B:377:HIS:CE1	1:B:460:THR:CG2	3.01	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	225/248 (91%)	222 (99%)	2 (1%)	1 (0%)	39	33
1	B	216/248 (87%)	211 (98%)	5 (2%)	0	100	100
All	All	441/496 (89%)	433 (98%)	7 (2%)	1 (0%)	52	48

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	419	GLU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	205/223 (92%)	191 (93%)	14 (7%)	20	13
1	B	198/223 (89%)	187 (94%)	11 (6%)	26	20
All	All	403/446 (90%)	378 (94%)	25 (6%)	22	16

All (25) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	363	ARG
1	A	417	SER
1	A	422	VAL
1	A	424	ILE
1	A	425	PHE
1	A	439	ASN
1	A	460	THR
1	A	464	SER
1	A	466	LEU
1	A	468	SER
1	A	481	LYS
1	A	501	HIS
1	A	522	MET
1	A	547	HIS
1	B	359	ASN
1	B	363	ARG
1	B	372	LEU
1	B	381	SER
1	B	413	ASN
1	B	422	VAL
1	B	425	PHE
1	B	469	LEU
1	B	481	LYS
1	B	526	TYR
1	B	544	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	502	GLN
1	A	506	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	377	HIS
1	B	476	HIS
1	B	488	HIS
1	B	502	GLN
1	B	506	GLN
1	B	519	ASN
1	B	524	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	LLB	A	1	-	34,39,39	1.62	7 (20%)	42,55,55	1.46	5 (11%)
2	LLB	B	1	-	34,39,39	1.54	9 (26%)	42,55,55	1.65	9 (21%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means

no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	LLB	A	1	-	-	0/15/28/28	0/5/5/5
2	LLB	B	1	-	-	0/15/28/28	0/5/5/5

All (16) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	1	LLB	C51-C52	2.01	1.42	1.38
2	A	1	LLB	C50-C51	2.15	1.42	1.38
2	B	1	LLB	C6-C1	2.16	1.43	1.38
2	B	1	LLB	C21-C20	2.36	1.43	1.38
2	B	1	LLB	C2-C1	2.36	1.41	1.37
2	B	1	LLB	C50-C51	2.38	1.43	1.38
2	B	1	LLB	C5-C6	2.42	1.41	1.36
2	B	1	LLB	C22-C16	2.47	1.43	1.39
2	A	1	LLB	C22-C16	2.47	1.43	1.39
2	A	1	LLB	C2-C3	2.59	1.41	1.37
2	A	1	LLB	C21-C22	2.66	1.43	1.38
2	A	1	LLB	C18-C16	2.69	1.43	1.39
2	B	1	LLB	C21-C22	3.00	1.44	1.38
2	B	1	LLB	C18-C16	3.00	1.44	1.39
2	A	1	LLB	C5-C6	3.35	1.43	1.36
2	A	1	LLB	C2-C1	4.23	1.45	1.37

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1	LLB	C5-C4-C12	-4.62	125.84	135.57
2	B	1	LLB	C5-C4-C12	-4.47	126.14	135.57
2	B	1	LLB	C62-C63-C64	-3.90	105.00	112.23
2	B	1	LLB	C6-C5-C4	-3.42	116.64	121.13
2	B	1	LLB	O27-C28-C29	-3.07	100.58	107.67
2	B	1	LLB	C1-C2-C3	-2.74	117.47	120.36
2	B	1	LLB	O17-C15-C12	-2.55	116.69	119.56
2	B	1	LLB	O13-C1-C2	-2.35	113.88	121.04
2	A	1	LLB	C5-C6-C1	-2.35	117.30	120.18
2	B	1	LLB	O27-C20-C19	-2.05	109.77	119.95
2	A	1	LLB	C65-C66-N61	2.52	115.08	110.96
2	B	1	LLB	C12-C15-C16	2.53	123.26	119.55
2	A	1	LLB	C63-C62-N61	3.12	116.06	110.96
2	A	1	LLB	C28-C29-N61	3.26	122.40	113.23

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 9 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1	LLB	3	0
2	B	1	LLB	6	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	229/248 (92%)	0.74	30 (13%) 5 5	24, 40, 72, 82	0
1	B	223/248 (89%)	0.57	21 (9%) 11 11	26, 39, 66, 80	0
All	All	452/496 (91%)	0.66	51 (11%) 7 7	24, 39, 70, 82	0

All (51) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	462	LEU	10.5
1	A	534	VAL	9.6
1	A	526	TYR	9.2
1	A	417	SER	8.6
1	B	418	VAL	7.9
1	A	463	SER	7.6
1	B	340	ALA	6.9
1	B	464	SER	6.9
1	A	535	PRO	6.9
1	B	417	SER	6.1
1	A	331	TYR	6.1
1	B	526	TYR	6.1
1	A	419	GLU	5.7
1	A	538	ASP	4.2
1	B	546	ALA	3.7
1	B	459	TYR	3.7
1	A	418	VAL	3.5
1	A	459	TYR	3.4
1	A	539	LEU	3.3
1	A	309	SER	3.2
1	B	416	LYS	3.2
1	B	535	PRO	3.1
1	A	461	PHE	3.1
1	B	331	TYR	3.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	391	LEU	3.0
1	A	548	ARG	3.0
1	A	549	LEU	2.9
1	A	537	TYR	2.9
1	B	545	ASP	2.8
1	A	330	GLU	2.8
1	A	464	SER	2.7
1	A	425	PHE	2.7
1	B	419	GLU	2.7
1	A	387	LEU	2.6
1	B	342	MET	2.6
1	B	306	LEU	2.6
1	A	536	LEU	2.5
1	A	340	ALA	2.5
1	B	420	GLY	2.4
1	A	306	LEU	2.3
1	A	428	LEU	2.3
1	A	465	THR	2.3
1	B	413	ASN	2.3
1	A	482	ILE	2.3
1	A	546	ALA	2.2
1	B	341	SER	2.2
1	A	308	LEU	2.2
1	B	309	SER	2.1
1	B	448	LEU	2.1
1	B	373	HIS	2.1
1	A	391	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains.

The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	LLB	A	1	35/35	0.90	0.18	0.25	29,34,46,50	0
2	LLB	B	1	35/35	0.91	0.14	-0.16	30,36,47,49	0

6.5 Other polymers [i](#)

There are no such residues in this entry.