



wwPDB X-ray Structure Validation Summary Report ⓘ

Jan 31, 2016 – 10:00 PM GMT

PDB ID : 1RMG
Title : RHAMNOGALACTURONASE A FROM ASPERGILLUS ACULEATUS
Authors : Petersen, T.N.; Kauppinen, S.; Larsen, S.
Deposited on : 1997-02-26
Resolution : 2.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	6249 (2.00-2.00)
Clashscore	102246	7340 (2.00-2.00)
Ramachandran outliers	100387	7248 (2.00-2.00)
Sidechain outliers	100360	7247 (2.00-2.00)
RSRZ outliers	91569	6262 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	422	<div> <div></div> <div>90%</div> <div>9%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	MAN	A	431	-	-	-	X

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	MAN	A	448	-	-	-	X
5	BMA	A	427	-	-	-	X

2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 3576 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called RHAMNOGALACTURONASE A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	422	3101	1939	506	638	18	0	0	0


- Molecule 2 is a polymer of unknown type called SUGAR (5-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
2	A	5	61	34	2	25	0	0

- Molecule 3 is a polymer of unknown type called SUGAR (3-MER).


Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
3	A	3	39	22	2	15	0	0

- Molecule 4 is SUGAR (ALPHA-D-MANNOSE) (three-letter code: MAN) (formula: C₆H₁₂O₆).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			11	6	5		
4	A	1	Total	C	O	0	0
			11	6	5		
4	A	1	Total	C	O	0	0
			11	6	5		
4	A	1	Total	C	O	0	0
			11	6	5		
4	A	1	Total	C	O	0	0
			11	6	5		
4	A	1	Total	C	O	0	0
			11	6	5		
4	A	1	Total	C	O	0	0
			11	6	5		
4	A	1	Total	C	O	0	0
			11	6	5		
4	A	1	Total	C	O	0	0
			11	6	5		
4	A	1	Total	C	O	0	0
			11	6	5		
4	A	1	Total	C	O	0	0
			11	6	5		

- Molecule 5 is SUGAR (BETA-D-MANNOSE) (three-letter code: BMA) (formula: C₆H₁₂O₆).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			11	6	5		
5	A	1	Total	C	O	0	0
			11	6	5		
5	A	1	Total	C	O	0	0
			11	6	5		

- Molecule 6 is SUGAR (GLUCOSE) (three-letter code: GLC) (formula: $C_6H_{12}O_6$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			11	6	5		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	177	Total 177	O 177	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: RHAMNOGALACTURONASE A



4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	62.90Å 125.40Å 137.00Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.00 – 2.00 29.25 – 2.00	Depositor EDS
% Data completeness (in resolution range)	92.7 (30.00-2.00) 92.2 (29.25-2.00)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	5.21 (at 2.00Å)	Xtriage
Refinement program	X-PLOR 3.1	Depositor
R, R_{free}	0.174 , 0.214 0.163 , 0.201	Depositor DCC
R_{free} test set	3399 reflections (11.09%)	DCC
Wilson B-factor (Å ²)	19.9	Xtriage
Anisotropy	0.361	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 54.1	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	1 of 34050 reflections (0.003%)	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	3576	wwPDB-VP
Average B, all atoms (Å ²)	23.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.69% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GLC, BMA, NAG, MAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.47	0/3169	0.74	0/4335

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3101	0	2933	18	0
2	A	61	0	52	0	0
3	A	39	0	34	0	0
4	A	154	0	140	2	0
5	A	33	0	30	1	0
6	A	11	0	10	0	0
7	A	177	0	0	1	0
All	All	3576	0	3199	18	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 18 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:113:GLN:HE21	1:A:115:PHE:H	1.37	0.70
1:A:184:SER:HB3	1:A:207:ASN:OD1	1.92	0.69
1:A:335:ASP:HB3	7:A:552:HOH:O	1.97	0.64
1:A:374:THR:HG21	4:A:432:MAN:O2	2.05	0.56
1:A:370:THR:HG21	5:A:427:BMA:O2	2.08	0.54

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	420/422 (100%)	391 (93%)	27 (6%)	2 (0%)	34 26

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	195	ASN
1	A	366	HIS

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	336/336 (100%)	325 (97%)	11 (3%)	45 43

5 of 11 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	184	SER
1	A	215	TYR
1	A	349	LEU
1	A	134	ASP
1	A	330	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	77	GLN
1	A	113	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

8 carbohydrates are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	NAG	A	423	1,2	14,14,15	0.74	0	15,19,21	0.75	0
2	NAG	A	424	2	14,14,15	0.50	0	15,19,21	0.78	1 (6%)
3	NAG	A	425	1,3	14,14,15	0.55	0	15,19,21	0.60	0
2	BMA	A	441	2	11,11,12	0.69	0	14,15,17	0.78	0
2	BMA	A	442	2	11,11,12	0.56	0	14,15,17	0.88	1 (7%)
3	NAG	A	443	3	14,14,15	0.78	1 (7%)	15,19,21	1.31	2 (13%)
2	MAN	A	445	2	11,11,12	0.58	0	14,15,17	0.71	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	BMA	A	446	3	11,11,12	0.56	0	14,15,17	0.55	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	A	423	1,2	-	0/6/23/26	0/1/1/1
2	NAG	A	424	2	-	0/6/23/26	0/1/1/1
3	NAG	A	425	1,3	-	0/6/23/26	0/1/1/1
2	BMA	A	441	2	-	0/2/19/22	0/1/1/1
2	BMA	A	442	2	-	0/2/19/22	0/1/1/1
3	NAG	A	443	3	-	0/6/23/26	0/1/1/1
2	MAN	A	445	2	-	0/2/19/22	1/1/1/1
3	BMA	A	446	3	-	0/2/19/22	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	443	NAG	C1-C2	2.26	1.55	1.52

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	443	NAG	C4-C3-C2	-3.31	106.09	111.23
2	A	424	NAG	C2-N2-C7	-2.38	119.98	123.04
3	A	443	NAG	C1-O5-C5	2.22	115.06	112.25
2	A	442	BMA	C1-C2-C3	2.41	112.39	109.54

There are no chirality outliers.

There are no torsion outliers.

All (1) ring outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	445	MAN	C1-C2-C3-C4-C5-O5

No monomer is involved in short contacts.

5.6 Ligand geometry

18 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
4	MAN	A	426	1	11,11,12	0.59	0	14,15,17	0.77	0
5	BMA	A	427	1	11,11,12	0.47	0	14,15,17	1.12	2 (14%)
4	MAN	A	428	1	11,11,12	0.46	0	14,15,17	0.66	0
4	MAN	A	429	1	11,11,12	0.56	0	14,15,17	0.68	0
4	MAN	A	430	1	11,11,12	0.66	0	14,15,17	0.66	0
4	MAN	A	431	1	11,11,12	0.47	0	14,15,17	0.74	0
4	MAN	A	432	1	11,11,12	0.33	0	14,15,17	0.77	0
6	GLC	A	433	1	11,11,12	0.61	0	14,15,17	1.15	0
4	MAN	A	434	1	11,11,12	0.53	0	14,15,17	0.81	1 (7%)
5	BMA	A	435	1	11,11,12	0.50	0	14,15,17	0.79	1 (7%)
4	MAN	A	436	1	11,11,12	0.62	0	14,15,17	1.07	2 (14%)
4	MAN	A	437	1	11,11,12	0.55	0	14,15,17	0.87	0
4	MAN	A	438	1	11,11,12	0.54	0	14,15,17	1.05	1 (7%)
5	BMA	A	439	1	11,11,12	0.61	0	14,15,17	0.92	1 (7%)
4	MAN	A	440	1	11,11,12	0.61	0	14,15,17	0.91	1 (7%)
4	MAN	A	444	1	11,11,12	0.65	0	14,15,17	0.75	0
4	MAN	A	447	1	11,11,12	0.52	0	14,15,17	0.94	0
4	MAN	A	448	1	11,11,12	1.15	1 (9%)	14,15,17	1.15	2 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	MAN	A	426	1	-	0/2/19/22	0/1/1/1
5	BMA	A	427	1	-	0/2/19/22	0/1/1/1
4	MAN	A	428	1	-	0/2/19/22	0/1/1/1
4	MAN	A	429	1	-	0/2/19/22	1/1/1/1
4	MAN	A	430	1	-	0/2/19/22	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	MAN	A	431	1	-	0/2/19/22	0/1/1/1
4	MAN	A	432	1	-	0/2/19/22	1/1/1/1
6	GLC	A	433	1	-	0/2/19/22	0/1/1/1
4	MAN	A	434	1	-	0/2/19/22	0/1/1/1
5	BMA	A	435	1	-	0/2/19/22	0/1/1/1
4	MAN	A	436	1	-	0/2/19/22	0/1/1/1
4	MAN	A	437	1	-	0/2/19/22	0/1/1/1
4	MAN	A	438	1	-	0/2/19/22	0/1/1/1
5	BMA	A	439	1	-	0/2/19/22	1/1/1/1
4	MAN	A	440	1	-	0/2/19/22	0/1/1/1
4	MAN	A	444	1	-	0/2/19/22	0/1/1/1
4	MAN	A	447	1	-	0/2/19/22	0/1/1/1
4	MAN	A	448	1	-	0/2/19/22	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	448	MAN	O5-C5	2.32	1.48	1.43

The worst 5 of 11 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	435	BMA	C1-C2-C3	-2.15	107.00	109.54
5	A	427	BMA	C1-O5-C5	2.23	115.08	112.25
4	A	436	MAN	O5-C1-C2	2.25	114.51	110.86
4	A	440	MAN	C1-O5-C5	2.27	115.13	112.25
4	A	434	MAN	C1-C2-C3	2.31	112.28	109.54

There are no chirality outliers.

There are no torsion outliers.

All (3) ring outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	439	BMA	C1-C2-C3-C4-C5-O5
4	A	429	MAN	C1-C2-C3-C4-C5-O5
4	A	432	MAN	C1-C2-C3-C4-C5-O5

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	427	BMA	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	432	MAN	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	422/422 (100%)	-0.53	1 (0%) 95 95	9, 19, 38, 57	0

All (1) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	288	THR	2.7

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	NAG	A	423	14/15	0.97	0.08	1.02	18,26,34,34	0
2	BMA	A	441	11/12	0.71	0.26	-	62,64,67,67	0
3	NAG	A	425	14/15	0.89	0.18	-	40,48,57,61	0
3	BMA	A	446	11/12	0.74	0.52	-	78,79,80,82	0
2	MAN	A	445	11/12	0.70	0.50	-	71,75,78,80	0
2	BMA	A	442	11/12	0.83	0.25	-	56,60,64,68	0
2	NAG	A	424	14/15	0.91	0.14	-	29,38,51,53	0
3	NAG	A	443	14/15	0.73	0.34	-	66,69,72,75	0

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
4	MAN	A	448	11/12	0.68	0.39	10.74	65,69,70,72	0
5	BMA	A	427	11/12	0.91	0.23	7.15	51,55,60,66	0
4	MAN	A	431	11/12	0.94	0.14	2.92	28,36,40,45	0
4	MAN	A	434	11/12	0.89	0.11	1.29	45,46,49,51	0
4	MAN	A	447	11/12	0.88	0.16	0.24	53,58,61,63	0
4	MAN	A	430	11/12	0.97	0.08	0.10	11,18,23,30	0
4	MAN	A	437	11/12	0.95	0.07	-0.82	29,34,39,43	0
4	MAN	A	438	11/12	0.80	0.21	-	42,47,53,57	0
6	GLC	A	433	11/12	0.91	0.17	-	42,50,54,58	0
4	MAN	A	429	11/12	0.76	0.27	-	48,58,65,68	0
4	MAN	A	426	11/12	0.85	0.23	-	47,57,61,62	0
5	BMA	A	439	11/12	0.81	0.26	-	57,61,65,70	0
4	MAN	A	428	11/12	0.88	0.14	-	38,43,47,47	0
4	MAN	A	436	11/12	0.90	0.19	-	47,51,54,59	0
4	MAN	A	432	11/12	0.84	0.18	-	47,53,57,63	0
4	MAN	A	440	11/12	0.74	0.28	-	54,58,61,63	0
5	BMA	A	435	11/12	0.85	0.20	-	48,53,59,61	0
4	MAN	A	444	11/12	0.85	0.21	-	42,50,57,62	0

6.5 Other polymers [i](#)

There are no such residues in this entry.