



# wwPDB X-ray Structure Validation Summary Report ⓘ

Jan 31, 2016 – 10:04 PM GMT

PDB ID : 1RXX  
Title : Structure of arginine deiminase  
Authors : Galkin, A.; Kulakova, L.; Sarikaya, E.; Lim, K.; Howard, A.; Herzberg, O.;  
Structure 2 Function Project (S2F)  
Deposited on : 2003-12-18  
Resolution : 2.45 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.45 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	4776 (2.50-2.42)
Clashscore	102246	1030 (2.48-2.44)
Ramachandran outliers	100387	1024 (2.48-2.44)
Sidechain outliers	100360	1024 (2.48-2.44)
RSRZ outliers	91569	4787 (2.50-2.42)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	421	<div> <div>2%</div> <div>67%</div> <div>25%</div> <div>5%</div> <div>.</div> </div>
1	B	421	<div> <div>%</div> <div>68%</div> <div>23%</div> <div>5%</div> <div>.</div> </div>
1	C	421	<div> <div>3%</div> <div>70%</div> <div>24%</div> <div>.</div> <div>.</div> <div>.</div> </div>
1	D	421	<div> <div>%</div> <div>65%</div> <div>27%</div> <div>.</div> <div>5%</div> </div>

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 13187 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Arginine deiminase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	408	Total	C	N	O	S	Se	0	0	0
			3185	2015	553	600	7	10			
1	B	404	Total	C	N	O	S	Se	0	0	0
			3160	2001	549	593	7	10			
1	C	412	Total	C	N	O	S	Se	0	0	0
			3214	2033	557	607	7	10			
1	D	402	Total	C	N	O	S	Se	0	0	0
			3136	1986	543	590	7	10			

There are 56 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	GLY	-	CLONING ARTIFACT	UNP P13981
A	-1	SER	-	CLONING ARTIFACT	UNP P13981
A	0	HIS	-	CLONING ARTIFACT	UNP P13981
A	1	MSE	-	CLONING ARTIFACT	UNP P13981
A	21	MSE	MET	MODIFIED RESIDUE	UNP P13981
A	62	MSE	MET	MODIFIED RESIDUE	UNP P13981
A	72	MSE	MET	MODIFIED RESIDUE	UNP P13981
A	143	MSE	MET	MODIFIED RESIDUE	UNP P13981
A	180	MSE	MET	MODIFIED RESIDUE	UNP P13981
A	229	MSE	MET	MODIFIED RESIDUE	UNP P13981
A	240	MSE	MET	MODIFIED RESIDUE	UNP P13981
A	277	MSE	MET	MODIFIED RESIDUE	UNP P13981
A	316	MSE	MET	MODIFIED RESIDUE	UNP P13981
A	407	MSE	MET	MODIFIED RESIDUE	UNP P13981
B	-2	GLY	-	CLONING ARTIFACT	UNP P13981
B	-1	SER	-	CLONING ARTIFACT	UNP P13981
B	0	HIS	-	CLONING ARTIFACT	UNP P13981
B	1	MSE	-	CLONING ARTIFACT	UNP P13981
B	21	MSE	MET	MODIFIED RESIDUE	UNP P13981
B	62	MSE	MET	MODIFIED RESIDUE	UNP P13981
B	72	MSE	MET	MODIFIED RESIDUE	UNP P13981

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
B	143	MSE	MET	MODIFIED RESIDUE	UNP P13981
B	180	MSE	MET	MODIFIED RESIDUE	UNP P13981
B	229	MSE	MET	MODIFIED RESIDUE	UNP P13981
B	240	MSE	MET	MODIFIED RESIDUE	UNP P13981
B	277	MSE	MET	MODIFIED RESIDUE	UNP P13981
B	316	MSE	MET	MODIFIED RESIDUE	UNP P13981
B	407	MSE	MET	MODIFIED RESIDUE	UNP P13981
C	-2	GLY	-	CLONING ARTIFACT	UNP P13981
C	-1	SER	-	CLONING ARTIFACT	UNP P13981
C	0	HIS	-	CLONING ARTIFACT	UNP P13981
C	1	MSE	-	CLONING ARTIFACT	UNP P13981
C	21	MSE	MET	MODIFIED RESIDUE	UNP P13981
C	62	MSE	MET	MODIFIED RESIDUE	UNP P13981
C	72	MSE	MET	MODIFIED RESIDUE	UNP P13981
C	143	MSE	MET	MODIFIED RESIDUE	UNP P13981
C	180	MSE	MET	MODIFIED RESIDUE	UNP P13981
C	229	MSE	MET	MODIFIED RESIDUE	UNP P13981
C	240	MSE	MET	MODIFIED RESIDUE	UNP P13981
C	277	MSE	MET	MODIFIED RESIDUE	UNP P13981
C	316	MSE	MET	MODIFIED RESIDUE	UNP P13981
C	407	MSE	MET	MODIFIED RESIDUE	UNP P13981
D	-2	GLY	-	CLONING ARTIFACT	UNP P13981
D	-1	SER	-	CLONING ARTIFACT	UNP P13981
D	0	HIS	-	CLONING ARTIFACT	UNP P13981
D	1	MSE	-	CLONING ARTIFACT	UNP P13981
D	21	MSE	MET	MODIFIED RESIDUE	UNP P13981
D	62	MSE	MET	MODIFIED RESIDUE	UNP P13981
D	72	MSE	MET	MODIFIED RESIDUE	UNP P13981
D	143	MSE	MET	MODIFIED RESIDUE	UNP P13981
D	180	MSE	MET	MODIFIED RESIDUE	UNP P13981
D	229	MSE	MET	MODIFIED RESIDUE	UNP P13981
D	240	MSE	MET	MODIFIED RESIDUE	UNP P13981
D	277	MSE	MET	MODIFIED RESIDUE	UNP P13981
D	316	MSE	MET	MODIFIED RESIDUE	UNP P13981
D	407	MSE	MET	MODIFIED RESIDUE	UNP P13981

- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	113	Total O 113 113	0	0
2	B	147	Total O 147 147	0	0

*Continued on next page...*


*Continued from previous page...*

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	C	144	Total 144	O 144	0	0
2	D	88	Total 88	O 88	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: Arginine deiminase




#### • Molecule 1: Arginine deiminase




#### • Molecule 1: Arginine deiminase





• Molecule 1: Arginine deiminase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	114.90 Å   114.90 Å   300.70 Å 90.00°   90.00°   90.00°	Depositor
Resolution (Å)	20.00 – 2.45 19.95 – 2.45	Depositor EDS
% Data completeness (in resolution range)	99.9 (20.00-2.45) 99.9 (19.95-2.45)	Depositor EDS
$R_{merge}$	0.10	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	5.34 (at 2.44 Å)	Xtriage
Refinement program	CNS 1.0	Depositor
R, $R_{free}$	0.198   ,   0.264 0.200   ,   0.264	Depositor DCC
$R_{free}$ test set	3802 reflections (5.09%)	DCC
Wilson B-factor (Å <sup>2</sup> )	42.8	Xtriage
Anisotropy	0.316	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 59.7	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.51$ , $\langle L^2 \rangle = 0.34$	Xtriage
Outliers	2 of 74675 reflections (0.003%)	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	13187	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	44.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 16.68% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.



## 5 Model quality

### 5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.70	2/3242 (0.1%)	0.91	6/4379 (0.1%)
1	B	0.70	1/3216 (0.0%)	0.89	4/4342 (0.1%)
1	C	0.70	1/3273 (0.0%)	0.93	7/4422 (0.2%)
1	D	0.65	2/3192 (0.1%)	0.85	1/4313 (0.0%)
All	All	0.69	6/12923 (0.0%)	0.90	18/17456 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	C	0	2

The worst 5 of 6 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	180	MSE	SE-CE	-7.41	1.51	1.95
1	D	363	CYS	CB-SG	-6.15	1.71	1.82
1	A	363	CYS	CB-SG	-6.08	1.72	1.82
1	A	169	CYS	CB-SG	-5.85	1.72	1.81
1	C	37	CYS	CB-SG	-5.61	1.72	1.81

The worst 5 of 18 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	347	SER	N-CA-C	9.03	135.38	111.00
1	C	400	GLY	N-CA-C	-8.59	91.63	113.10
1	C	401	ARG	NE-CZ-NH2	-7.93	116.34	120.30
1	A	373	ASP	CB-CG-OD2	6.32	123.99	118.30
1	C	346	ASN	N-CA-C	6.18	127.68	111.00

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	C	144	TYR	Sidechain
1	C	348	PHE	Sidechain

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3185	0	3161	121	0
1	B	3160	0	3139	114	0
1	C	3214	0	3186	99	0
1	D	3136	0	3119	106	0
2	A	113	0	0	15	0
2	B	147	0	0	16	0
2	C	144	0	0	20	0
2	D	88	0	0	13	0
All	All	13187	0	12605	432	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 17.

The worst 5 of 432 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:352:ARG:HG3	1:B:352:ARG:HH11	1.13	1.11
1:A:343:THR:HG21	1:A:358:GLY:H	1.15	1.08
1:B:343:THR:HG21	1:B:358:GLY:H	1.21	1.04
1:D:343:THR:HG21	1:D:358:GLY:H	1.20	1.02
1:B:363:CYS:HB2	2:B:517:HOH:O	1.61	1.01

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	402/421 (96%)	381 (95%)	18 (4%)	3 (1%)	26	32
1	B	396/421 (94%)	373 (94%)	21 (5%)	2 (0%)	34	41
1	C	408/421 (97%)	385 (94%)	18 (4%)	5 (1%)	16	17
1	D	396/421 (94%)	371 (94%)	23 (6%)	2 (0%)	34	41
All	All	1602/1684 (95%)	1510 (94%)	80 (5%)	12 (1%)	26	32

5 of 12 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	402	GLY
1	C	347	SER
1	C	402	GLY
1	D	152	SER
1	A	402	GLY

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	347/346 (100%)	318 (92%)	29 (8%)	14	17
1	B	343/346 (99%)	316 (92%)	27 (8%)	15	20
1	C	349/346 (101%)	328 (94%)	21 (6%)	24	33
1	D	342/346 (99%)	318 (93%)	24 (7%)	19	25
All	All	1381/1384 (100%)	1280 (93%)	101 (7%)	17	23

5 of 101 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	305	SER
1	C	20	VAL
1	D	305	SER
1	B	331	GLU
1	B	382	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 18 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	317	ASN
1	C	29	HIS
1	D	81	GLN
1	B	55	HIS
1	B	199	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	398/421 (94%)	-0.28	8 (2%) 68 71	26, 41, 70, 91	0
1	B	394/421 (93%)	-0.35	5 (1%) 79 81	22, 38, 66, 97	0
1	C	402/421 (95%)	-0.25	12 (2%) 54 57	23, 39, 69, 99	0
1	D	392/421 (93%)	-0.23	3 (0%) 87 89	31, 46, 73, 89	0
All	All	1586/1684 (94%)	-0.28	28 (1%) 71 74	22, 42, 71, 99	0

The worst 5 of 28 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	351	GLU	8.2
1	B	418	TYR	5.9
1	C	347	SER	5.7
1	C	350	ALA	5.5
1	C	399	ARG	5.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.