



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 01:03 PM GMT

PDB ID : 3SQ0  
Title : DNA Polymerase(L561A/S565G/Y567A) Ternary Complex with dUpNpp  
Opposite dA (Mn2+)  
Authors : Xia, S.; Konigsberg, W.H.; Wang, J.  
Deposited on : 2011-07-04  
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	6249 (2.00-2.00)
Clashscore	102246	7340 (2.00-2.00)
Ramachandran outliers	100387	7248 (2.00-2.00)
Sidechain outliers	100360	7247 (2.00-2.00)
RSRZ outliers	91569	6262 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	903	<div> <div>3%</div> <div>91%</div> <div>8%</div> </div>
2	T	18	<div> <div>11%</div> <div>83%</div> <div>17%</div> </div>
3	P	13	<div> <div>23%</div> <div>85%</div> <div>8%</div> <div>8%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	MN	A	905	-	-	-	X

## 2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 8343 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA polymerase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	903	Total	C	N	O	S	0	2	0
			7377	4737	1231	1376	33			

There are 5 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	222	ALA	ASP	ENGINEERED MUTATION	UNP Q38087
A	327	ALA	ASP	ENGINEERED MUTATION	UNP Q38087
A	561	ALA	LEU	ENGINEERED MUTATION	UNP Q38087
A	565	GLY	SER	ENGINEERED MUTATION	UNP Q38087
A	567	ALA	TYR	ENGINEERED MUTATION	UNP Q38087


- Molecule 2 is a DNA chain called 5'-D(\*TP\*CP\*AP\*AP\*GP\*TP\*AP\*AP\*GP\*CP\*AP\*GP\*TP\*CP\*CP\*GP\*CP\*G)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	T	18	Total	C	N	O	P	0	0	0
			367	175	71	104	17			

- Molecule 3 is a DNA chain called 5'-D(\*GP\*CP\*GP\*GP\*AP\*CP\*TP\*GP\*CP\*TP\*TP\*AP\*(DOC))-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	P	13	Total	C	N	O	P	0	0	0
			262	126	48	76	12			

- Molecule 4 is 2'-DEOXYURIDINE 5'-ALPHA,BETA-IMIDO-TRIPHOSPHATE (three-letter code: DUP) (formula: C<sub>9</sub>H<sub>16</sub>N<sub>3</sub>O<sub>13</sub>P<sub>3</sub>).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O	P	0	0
			28	9	3	13	3		

- Molecule 5 is MANGANESE (II) ION (three-letter code: MN) (formula: Mn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	5	Total	Mn	0	0
			5	5		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	280	Total	O	0	2
			282	282		
6	T	16	Total	O	0	0
			16	16		
6	P	6	Total	O	0	0
			6	6		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $\text{RSRZ} > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: DNA polymerase



- Molecule 2: 5'-D(\*TP\*CP\*AP\*AP\*GP\*TP\*AP\*AP\*GP\*CP\*AP\*GP\*TP\*CP\*CP\*GP\*CP\*G)-3'



- Molecule 3: 5'-D(\*GP\*CP\*GP\*GP\*AP\*CP\*TP\*GP\*CP\*TP\*TP\*AP\*(DOC))-3'



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	78.15Å 119.67Å 130.64Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	88.24 – 2.00 47.51 – 2.00	Depositor EDS
% Data completeness (in resolution range)	98.7 (88.24-2.00) 98.6 (47.51-2.00)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.30 (at 2.00Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, $R_{free}$	0.209 , 0.245 0.213 , 0.250	Depositor DCC
$R_{free}$ test set	4113 reflections (5.26%)	DCC
Wilson B-factor (Å <sup>2</sup> )	36.4	Xtriage
Anisotropy	0.167	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 39.3	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 82448 reflections	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	8343	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	45.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.10% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: DUP, DOC, MN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.42	0/7564	0.53	0/10220
2	T	0.72	0/412	1.27	3/634 (0.5%)
3	P	0.73	0/273	1.20	1/420 (0.2%)
All	All	0.45	0/8249	0.63	4/11274 (0.0%)

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	T	7	DA	O4'-C1'-N9	-5.97	103.82	108.00
3	P	105	DG	P-O3'-C3'	5.87	126.74	119.70
2	T	8	DA	O4'-C1'-N9	5.79	112.06	108.00
2	T	12	DG	O4'-C1'-N9	5.29	111.70	108.00

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	7377	0	7275	53	0
2	T	367	0	203	0	0
3	P	262	0	148	1	0
4	A	28	0	12	0	0

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	A	5	0	0	0	0
6	A	282	0	0	5	0
6	P	6	0	0	0	0
6	T	16	0	0	0	0
All	All	8343	0	7638	54	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (54) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:893:LYS:HA	1:A:894:LYS:HB2	1.01	1.01
1:A:893:LYS:CA	1:A:894:LYS:HB2	1.91	1.00
1:A:893:LYS:HA	1:A:894:LYS:CB	1.93	0.97
1:A:506:PRO:HA	1:A:507:ASN:CB	2.03	0.88
1:A:506:PRO:HA	1:A:507:ASN:HB2	1.62	0.81
1:A:330:ARG:HH11	1:A:333:GLN:HE22	1.32	0.75
1:A:467:ARG:HG3	1:A:467:ARG:HH11	1.55	0.72
1:A:203:ASN:ND2	1:A:241:ARG:HH22	1.90	0.69
1:A:203:ASN:HD22	1:A:241:ARG:HH22	1.40	0.69
1:A:467:ARG:HH11	1:A:467:ARG:CG	2.08	0.67
1:A:825:VAL:HB	1:A:828:GLU:HG3	1.79	0.65
1:A:871:LEU:CD2	6:A:930:HOH:O	2.43	0.65
1:A:871:LEU:HD21	6:A:930:HOH:O	1.97	0.62
1:A:854:ILE:HD13	1:A:862:VAL:HG21	1.82	0.61
1:A:170:LEU:HA	1:A:177:GLU:HG3	1.84	0.59
1:A:506:PRO:HA	1:A:507:ASN:HB3	1.86	0.57
1:A:468:ASP:HB2	6:A:979:HOH:O	2.04	0.56
1:A:467:ARG:NH1	1:A:467:ARG:CG	2.72	0.53
1:A:655:ALA:HA	1:A:659:MET:HE2	1.91	0.51
1:A:415:LEU:HD22	1:A:623:ASP:HB3	1.92	0.51
1:A:408:MET:HE1	1:A:659:MET:HE1	1.93	0.50
1:A:408:MET:CE	1:A:659:MET:HE1	2.41	0.49
1:A:862:VAL:O	1:A:866:MET:HG3	2.12	0.49
1:A:203:ASN:HD22	1:A:241:ARG:NH2	2.10	0.48
1:A:330:ARG:NH1	1:A:333:GLN:HE22	2.08	0.47
1:A:638:GLU:HA	1:A:639:SER:HA	1.52	0.46
1:A:204:PHE:CE1	1:A:208:LYS:HD2	2.50	0.46
1:A:894:LYS:HD3	1:A:894:LYS:HA	1.68	0.45
1:A:412:LEU:HB2	1:A:623:ASP:HB2	1.99	0.44

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:664:ASP:OD1	1:A:679:HIS:HE1	2.00	0.44
1:A:40:HIS:HD2	6:A:1117:HOH:O	2.01	0.44
1:A:279:LYS:HE2	1:A:358:VAL:O	2.18	0.43
3:P:105:DG:H2"	3:P:106:DG:OP2	2.18	0.43
1:A:441:ASP:HB3	1:A:447:ALA:HB2	2.00	0.43
1:A:660:GLU:HB3	1:A:661:PRO:HD3	2.00	0.43
1:A:760:LEU:HD13	1:A:891:TYR:HA	2.00	0.43
1:A:514:LEU:HD21	1:A:529:LYS:HB3	1.99	0.43
1:A:436:VAL:HG12	1:A:437:ALA:O	2.19	0.43
1:A:793:VAL:HG23	1:A:793:VAL:O	2.20	0.42
1:A:440[B]:HIS:HB3	6:A:962:HOH:O	2.19	0.42
1:A:785:ALA:HB1	1:A:788:ILE:HD11	2.01	0.42
1:A:330:ARG:HH11	1:A:333:GLN:NE2	2.07	0.42
1:A:408:MET:CE	1:A:659:MET:CE	2.97	0.42
1:A:752:MET:HG3	1:A:760:LEU:HG	2.01	0.42
1:A:152:LEU:HD11	1:A:161:GLU:HG3	2.02	0.42
1:A:636:VAL:HG12	1:A:637:GLY:O	2.20	0.41
1:A:202:LEU:O	1:A:206:GLN:HG2	2.20	0.41
1:A:279:LYS:HE3	1:A:280:PHE:CE2	2.55	0.41
1:A:40:HIS:HE1	1:A:51:ASP:OD2	2.03	0.41
1:A:506:PRO:CA	1:A:507:ASN:CB	2.85	0.41
1:A:901:PHE:HD1	1:A:903:PHE:H	1.68	0.41
1:A:465:LYS:O	1:A:677:LYS:HE2	2.21	0.41
1:A:410:PHE:HB3	1:A:683:MET:HG2	2.01	0.40
1:A:41:CYS:HB2	1:A:42:PRO:HD2	2.04	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	903/903 (100%)	878 (97%)	21 (2%)	4 (0%)	39	33

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	894	LYS
1	A	507	ASN
1	A	638	GLU
1	A	898	PHE

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	799/797 (100%)	776 (97%)	23 (3%)	50	49

All (23) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	34	LYS
1	A	44	SER
1	A	66	ARG
1	A	76	GLU
1	A	177	GLU
1	A	309	ILE
1	A	332	LEU
1	A	467	ARG
1	A	468	ASP
1	A	505	ASN
1	A	507	ASN
1	A	514	LEU
1	A	580	LEU
1	A	658	ARG
1	A	739	LYS
1	A	760	LEU
1	A	819	ILE
1	A	835	LEU
1	A	878	LYS
1	A	893	LYS
1	A	894	LYS

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	900	MET
1	A	901	PHE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (14) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	40	HIS
1	A	112	ASN
1	A	131	HIS
1	A	203	ASN
1	A	206	GLN
1	A	333	GLN
1	A	339	GLN
1	A	376	GLN
1	A	546	GLN
1	A	564	ASN
1	A	679	HIS
1	A	761	GLN
1	A	786	ASN
1	A	823	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	DOC	P	115	3,2	11,19,20	0.54	0	14,26,29	1.64	2 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	DOC	P	115	3,2	-	0/3/18/19	0/2/2/2

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	P	115	DOC	C2-N3-C4	3.52	120.57	115.61
3	P	115	DOC	O4'-C4'-C5'	3.81	115.16	109.54

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 5 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	DUP	A	904	5	23,29,29	2.64	5 (21%)	32,45,45	2.18	9 (28%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical

component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	DUP	A	904	5	-	0/13/34/34	0/2/2/2

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	904	DUP	PA-O2A	-2.12	1.50	1.56
4	A	904	DUP	PA-O1A	4.38	1.51	1.46
4	A	904	DUP	O4-C4	4.50	1.35	1.24
4	A	904	DUP	PB-O1B	4.91	1.51	1.46
4	A	904	DUP	PG-O3G	8.73	1.86	1.54

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	904	DUP	O1A-PA-N3A	-4.04	105.70	111.90
4	A	904	DUP	C2'-C1'-N1	-3.85	104.80	114.16
4	A	904	DUP	O1B-PB-N3A	-3.77	106.11	111.90
4	A	904	DUP	PG-O3B-PB	-2.41	124.58	132.67
4	A	904	DUP	O1G-PG-O2G	2.58	118.88	110.58
4	A	904	DUP	O4'-C1'-N1	3.24	113.32	107.72
4	A	904	DUP	O2B-PB-O1B	3.57	117.45	110.00
4	A	904	DUP	O2A-PA-O1A	4.07	118.49	110.00
4	A	904	DUP	C4-N3-C2	6.11	120.19	114.14

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å <sup>2</sup> )	Q<0.9
1	A	903/903 (100%)	0.20	29 (3%)	51	52	27, 41, 66, 105	1 (0%)
2	T	18/18 (100%)	0.49	2 (11%)	7	8	32, 53, 110, 134	0
3	P	12/13 (92%)	1.11	3 (25%)	1	1	32, 78, 120, 125	0
All	All	933/934 (99%)	0.22	34 (3%)	46	48	27, 41, 70, 134	1 (0%)

All (34) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	256	MET	8.1
2	T	1	DT	6.9
1	A	253	ILE	6.4
1	A	508	LEU	6.3
1	A	257	TYR	4.6
1	A	510	VAL	4.5
1	A	901	PHE	4.3
1	A	254	GLU	4.2
1	A	903	PHE	3.5
1	A	507	ASN	3.5
1	A	637	GLY	3.4
3	P	104	DC	3.3
1	A	851	GLY	3.1
1	A	252	VAL	3.0
1	A	45	GLN	2.9
1	A	44	SER	2.8
1	A	255	ASN	2.8
1	A	260	ARG	2.8
1	A	48	LYS	2.7
1	A	532	LYS	2.7
1	A	466	ASP	2.6
3	P	106	DG	2.6
1	A	893	LYS	2.6

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	899	ASP	2.6
1	A	49	TYR	2.4
3	P	103	DG	2.4
1	A	819	ILE	2.3
1	A	63	ALA	2.3
1	A	854	ILE	2.3
1	A	46	ALA	2.3
1	A	820	ASP	2.2
1	A	375	GLU	2.2
2	T	15	DC	2.1
1	A	192	ASP	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
3	DOC	P	115	18/19	0.97	0.15	-	27,33,37,40	0

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
5	MN	A	905	1/1	0.99	0.17	2.03	30,30,30,30	0
4	DUP	A	904	28/28	0.98	0.18	1.91	25,29,32,33	0
5	MN	A	907	1/1	0.98	0.06	-4.50	53,53,53,53	0

*Continued on next page...*



*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
5	MN	A	906	1/1	1.00	0.17	-	32,32,32,32	0
5	MN	A	908	1/1	0.98	0.06	-	52,52,52,52	0
5	MN	A	909	1/1	0.85	0.08	-	109,109,109,109	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.