



Full wwPDB X-ray Structure Validation Report ⓘ

Dec 20, 2016 – 04:46 PM EST

PDB ID : 5T1A
Title : Structure of CC Chemokine Receptor 2 with Orthosteric and Allosteric Antagonists
Authors : Zheng, Y.; Qin, L.; Ortiz Zacarias, N.V.; de Vries, H.; Han, G.W.; Gustavsson, M.; Dabros, M.; Zhao, C.; Cherney, R.J.; Carter, P.; Stamos, D.; Abagyan, R.; Cherezov, V.; Stevens, R.C.; IJzerman, A.P.; Heitman, L.H.; Tebben, A.; Kufareva, I.; Handel, T.M.
Deposited on : 2016-08-18
Resolution : 2.81 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7.1 (RC1), CSD as537be (2016)
Xtriage (Phenix)	:	1.9-1692
EDS	:	rb-20028442
Percentile statistics	:	20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	rb-20028442


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.81 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	2393 (2.80-2.80)
Clashscore	102246	2827 (2.80-2.80)
Ramachandran outliers	100387	2782 (2.80-2.80)
Sidechain outliers	100360	2784 (2.80-2.80)
RSRZ outliers	91569	2404 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	508	<div> <div>3%</div> <div>77%</div> <div>10%</div> <div>12%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	OLC	A	1207	-	-	-	X

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 3580 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Chimera protein of CC chemokine receptor type 2 isoform B and T4-lysozyme, Lysozyme.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	445	Total	C	N	O	S	0	0	0
			3458	2270	575	594	19			

There are 38 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-9	ASP	-	expression tag	UNP P41597
A	-8	TYR	-	expression tag	UNP P41597
A	-7	LYS	-	expression tag	UNP P41597
A	-6	ASP	-	expression tag	UNP P41597
A	-5	ASP	-	expression tag	UNP P41597
A	-4	ASP	-	expression tag	UNP P41597
A	-3	ASP	-	expression tag	UNP P41597
A	-2	LYS	-	expression tag	UNP P41597
A	-1	PRO	-	expression tag	UNP P41597
A	0	GLY	-	expression tag	UNP P41597
A	1	THR	-	expression tag	UNP P41597
A	226	SER	LEU	engineered mutation	UNP P41597
A	227	ARG	LYS	engineered mutation	UNP P41597
A	228	ALA	THR	engineered mutation	UNP P41597
A	229	SER	LEU	engineered mutation	UNP P41597
A	230	LYS	LEU	engineered mutation	UNP P41597
A	231	SER	ARG	engineered mutation	UNP P41597
A	232	ARG	CYS	engineered mutation	UNP P41597
A	233	ILE	ARG	engineered mutation	UNP P41597
A	1054	THR	CYS	engineered mutation	UNP D9IEF7
A	1097	ALA	CYS	engineered mutation	UNP D9IEF7
A	1162	PRO	LYS	engineered mutation	UNP D9IEF7
A	234	PRO	ASN	engineered mutation	UNP P41597
A	235	PRO	GLU	engineered mutation	UNP P41597
A	236	SER	LYS	engineered mutation	UNP P41597
A	237	ARG	LYS	engineered mutation	UNP P41597

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
A	238	GLU	ARG	engineered mutation	UNP P41597
A	239	LYS	HIS	engineered mutation	UNP P41597
A	240	LYS	ARG	engineered mutation	UNP P41597
A	329	GLY	-	engineered mutation	UNP P41597
A	330	ARG	-	engineered mutation	UNP P41597
A	331	PRO	-	engineered mutation	UNP P41597
A	332	LEU	-	engineered mutation	UNP P41597
A	333	GLU	-	engineered mutation	UNP P41597
A	334	VAL	-	expression tag	UNP P41597
A	335	LEU	-	expression tag	UNP P41597
A	336	PHE	-	expression tag	UNP P41597
A	337	GLN	-	expression tag	UNP P41597

- Molecule 2 is (3S)-1-{(1S,2R,4R)-4-[methyl(propan-2-yl)amino]-2-propylcyclohexyl}-3-{[6-(trifluoromethyl)quinazolin-4-yl]amino}pyrrolidin-2-one (three-letter code: 73R) (formula: $C_{26}H_{36}F_3N_5O$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	F	N	O	0	0
			35	26	3	5	1		

- Molecule 3 is (2 {R})-1-(4-chloranyl-2-fluoranyl-phenyl)-2-cyclohexyl-3-ethanoyl-4-oxidanyl-2 {H}-pyrrol-5-one (three-letter code: VT5) (formula: $C_{18}H_{19}ClFNO_3$).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
3	A	1	Total	C	Cl	F	N	O	0	0
			24	18	1	1	1	3		

- Molecule 4 is SULFATE ION (three-letter code: SO4) (formula: O₄S).




Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	O S	0	0
			5	4 1		
4	A	1	Total	O S	0	0
			5	4 1		
4	A	1	Total	O S	0	0
			5	4 1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	O	S	0	0
			5	4	1		

- Molecule 5 is (2R)-2,3-dihydroxypropyl (9Z)-octadec-9-enoate (three-letter code: OLC) (formula: C₂₁H₄₀O₄).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			25	21	4		

- Molecule 6 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	1	Total	Zn	0	0
			1	1		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	17	Total	O	0	0
			17	17		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Chimera protein of CC chemokine receptor type 2 isoform B and T4-lysozyme, Lysozyme



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	59.23 Å 64.69 Å 169.98 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	24.32 – 2.81 48.59 – 2.81	Depositor EDS
% Data completeness (in resolution range)	87.2 (24.32-2.81) 87.2 (48.59-2.81)	Depositor EDS
R_{merge}	0.24	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.50 (at 2.81 Å)	Xtriage
Refinement program	PHENIX	Depositor
R, R_{free}	0.234 , 0.274 0.235 , 0.277	Depositor DCC
R_{free} test set	746 reflections (5.14%)	DCC
Wilson B-factor (Å ²)	40.3	Xtriage
Anisotropy	0.120	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 51.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.85	EDS
Total number of atoms	3580	wwPDB-VP
Average B, all atoms (Å ²)	41.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.57% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, OLC, 73R, VT5, SO4, YCM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.22	0/3527	0.38	0/4802

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3458	0	3478	29	0
2	A	35	0	0	0	0
3	A	24	0	0	1	0
4	A	20	0	0	2	0
5	A	25	0	40	2	0
6	A	1	0	0	0	0
7	A	17	0	0	1	0
All	All	3580	0	3518	29	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (29) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1006:MET:HG3	1:A:1161:TYR:CZ	2.36	0.61
1:A:314:ARG:NH2	4:A:1203:SO4:O3	2.34	0.60
1:A:1015:LEU:HA	1:A:1058:ILE:HG22	1.84	0.60
1:A:1146:ALA:O	1:A:1150:ILE:HG12	2.06	0.56
1:A:1148:ARG:HB2	1:A:1160:ALA:HB1	1.89	0.55
1:A:1077:GLY:HA3	1:A:1108:GLU:OE2	2.08	0.54
1:A:232:ARG:HG3	1:A:239:LYS:HD3	1.90	0.52
1:A:71:LYS:HD2	1:A:78:ASP:OD2	2.10	0.52
1:A:1052:ARG:HH21	1:A:1054:THR:HA	1.76	0.50
1:A:1078:ILE:HG23	1:A:1084:LEU:HB3	1.96	0.48
1:A:1083:LYS:HE3	1:A:1115:THR:HG22	1.95	0.47
1:A:203:THR:O	1:A:207:ASN:ND2	2.45	0.47
1:A:311:LYS:HB2	3:A:1202:VT5:C4	2.44	0.47
1:A:208:ILE:HA	1:A:212:VAL:HB	1.96	0.46
1:A:135:THR:HG23	1:A:246:PHE:HE1	1.81	0.45
1:A:242:VAL:HA	1:A:245:ILE:HG22	1.98	0.45
1:A:132:ILE:HG23	1:A:217:ILE:HG21	1.99	0.44
1:A:1083:LYS:NZ	7:A:1302:HOH:O	2.50	0.44
1:A:213:LEU:HB3	1:A:214:PRO:HD3	2.01	0.43
1:A:127:GLY:O	1:A:131:ILE:HG13	2.19	0.43
1:A:1117:SER:OG	1:A:1132:ASN:ND2	2.51	0.43
1:A:243:ARG:HG2	5:A:1207:OLC:H2A	2.02	0.42
1:A:243:ARG:HH21	5:A:1207:OLC:H24A	1.85	0.42
1:A:1003:ILE:HG13	1:A:1006:MET:HE2	2.02	0.42
1:A:41:GLY:O	1:A:45:LEU:HB2	2.20	0.42
1:A:76:LEU:HD22	1:A:152:ARG:HA	2.01	0.41
1:A:1005:GLU:O	1:A:1009:ILE:HG12	2.21	0.41
1:A:257:THR:N	1:A:258:PRO:HD2	2.35	0.41
1:A:1144:ASN:HB2	4:A:1206:SO4:O4	2.21	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	442/508 (87%)	429 (97%)	13 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	357/442 (81%)	354 (99%)	3 (1%)	86	97

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	37	VAL
1	A	149	LEU
1	A	1153	PHE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	1069	GLN
1	A	1132	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length

(or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	YCM	A	181	1	7,9,10	1.23	1 (14%)	5,10,12	1.57	1 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	YCM	A	181	1	-	0/6/8/10	0/0/0/0

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	181	YCM	CD-SG	-2.08	1.77	1.81

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	181	YCM	CD-CE-NZ2	2.92	118.70	115.48

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 1 is monoatomic - leaving 7 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link

column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	73R	A	1201	-	36,38,38	1.98	5 (13%)	44,56,56	2.94	13 (29%)
3	VT5	A	1202	-	24,26,26	2.08	5 (20%)	30,38,38	1.54	3 (10%)
4	SO4	A	1203	-	4,4,4	0.25	0	6,6,6	0.06	0
4	SO4	A	1204	-	4,4,4	0.25	0	6,6,6	0.10	0
4	SO4	A	1205	-	4,4,4	0.23	0	6,6,6	0.07	0
4	SO4	A	1206	-	4,4,4	0.29	0	6,6,6	0.05	0
5	OLC	A	1207	-	24,24,24	0.69	1 (4%)	25,25,25	0.94	1 (4%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	73R	A	1201	-	-	0/25/51/51	0/4/4/4
3	VT5	A	1202	-	-	0/12/40/40	0/3/3/3
4	SO4	A	1203	-	-	0/0/0/0	0/0/0/0
4	SO4	A	1204	-	-	0/0/0/0	0/0/0/0
4	SO4	A	1205	-	-	0/0/0/0	0/0/0/0
4	SO4	A	1206	-	-	0/0/0/0	0/0/0/0
5	OLC	A	1207	-	-	0/24/24/24	0/0/0/0

All (11) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	1202	VT5	C12-N	-7.76	1.33	1.43
2	A	1201	73R	C14-C15	-3.89	1.46	1.52
3	A	1202	VT5	C4-N	-2.48	1.34	1.37
2	A	1201	73R	C8-C1	-2.31	1.37	1.41
2	A	1201	73R	C18-C14	-2.09	1.47	1.54
3	A	1202	VT5	C15-CL	2.28	1.79	1.74
5	A	1207	OLC	O20-C1	2.64	1.41	1.33
3	A	1202	VT5	O1-C4	2.65	1.28	1.23
2	A	1201	73R	C17-N16	2.95	1.53	1.47
3	A	1202	VT5	O2-C3	4.33	1.48	1.33
2	A	1201	73R	C15-N16	8.55	1.46	1.35

All (17) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1201	73R	N6-C5-N4	-9.30	121.56	128.87
2	A	1201	73R	C18-C17-N16	-7.80	95.24	103.30
2	A	1201	73R	C36-C35-C31	-4.03	107.45	114.61
2	A	1201	73R	C2-C1-N6	-3.93	118.57	122.83
2	A	1201	73R	C17-N16-C15	-3.56	111.06	113.56
2	A	1201	73R	C12-C13-C2	-2.22	120.17	122.24
2	A	1201	73R	F32-CAP-C12	-2.04	108.62	112.92
2	A	1201	73R	C31-C30-C29	2.02	113.56	109.20
2	A	1201	73R	C13-C2-C3	2.32	126.18	124.87
5	A	1207	OLC	O20-C1-C2	2.71	120.20	111.85
2	A	1201	73R	C15-C14-N7	3.03	115.70	111.37
3	A	1202	VT5	C15-C16-C17	3.67	120.21	117.31
2	A	1201	73R	C27-C28-C29	3.74	117.56	109.61
3	A	1202	VT5	C17-C12-N	3.78	123.91	120.43
3	A	1202	VT5	C7-C6-C5	3.82	115.04	111.29
2	A	1201	73R	C5-N6-C1	6.30	120.88	115.19
2	A	1201	73R	C5-N4-C3	8.70	122.73	116.47

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	1202	VT5	1	0
4	A	1203	SO4	1	0
4	A	1206	SO4	1	0
5	A	1207	OLC	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	444/508 (87%)	-0.11	17 (3%) 44 32	21, 37, 74, 118	0

All (17) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	277	CYS	8.4
1	A	273	GLY	5.1
1	A	187	VAL	4.8
1	A	188	TYR	3.7
1	A	318	VAL	3.7
1	A	276	ASN	3.5
1	A	275	SER	3.4
1	A	186	SER	3.1
1	A	37	VAL	3.0
1	A	184	GLU	3.0
1	A	281	SER	2.8
1	A	278	GLU	2.6
1	A	279	SER	2.4
1	A	320	PHE	2.4
1	A	269	GLN	2.4
1	A	185	ASP	2.3
1	A	103	ALA	2.2

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
1	YCM	A	181	10/11	0.92	0.19	-	51,56,65,65	0

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
5	OLC	A	1207	25/25	0.82	0.27	2.34	38,44,52,54	0
4	SO4	A	1206	5/5	0.86	0.25	0.57	66,66,67,67	0
2	73R	A	1201	35/35	0.92	0.20	0.13	26,37,42,42	0
3	VT5	A	1202	24/24	0.94	0.17	-0.19	16,25,34,57	0
4	SO4	A	1205	5/5	0.97	0.15	-0.70	53,53,53,53	0
4	SO4	A	1204	5/5	0.98	0.15	-0.87	29,30,31,33	0
6	ZN	A	1208	1/1	0.99	0.11	-1.63	40,40,40,40	0
4	SO4	A	1203	5/5	0.88	0.23	-	80,80,80,80	0

6.5 Other polymers [i](#)

There are no such residues in this entry.