



Full wwPDB X-ray Structure Validation Report ⓘ

Dec 19, 2016 – 10:00 PM EST

PDB ID : 5T55
Title : LECTIN FROM BAUHINIA FORFICATA IN COMPLEX WITH GLOBOTETRAOSE
Authors : Lubkowski, J.; Wlodawer, A.
Deposited on : 2016-08-30
Resolution : 1.43 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.1 (RC1), CSD as537be (2016)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20028442
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20028442

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	EDO	A	305	-	-	-	X
5	EDO	A	306	-	-	-	X
5	EDO	A	307[A]	-	-	-	X
5	EDO	A	307[B]	-	-	-	X
5	EDO	A	308	-	-	-	X
5	EDO	A	310[A]	-	-	-	X
5	EDO	A	310[B]	-	-	-	X
5	EDO	A	311	-	-	X	X
5	EDO	B	308	-	-	-	X
5	EDO	B	309	-	-	-	X

2 Entry composition

There are 9 unique types of molecules in this entry. The entry contains 4213 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Lectin.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
1	A	229	Total	C	N	O	0	7	0
			1830	1168	307	355			
1	B	229	Total	C	N	O	0	13	0
			1851	1180	313	358			

There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	234	GLY	-	expression tag	UNP P86993
A	235	ALA	-	expression tag	UNP P86993
A	236	ARG	-	expression tag	UNP P86993
A	237	HIS	-	expression tag	UNP P86993
A	238	HIS	-	expression tag	UNP P86993
A	239	HIS	-	expression tag	UNP P86993
A	240	HIS	-	expression tag	UNP P86993
A	241	HIS	-	expression tag	UNP P86993
A	242	HIS	-	expression tag	UNP P86993
B	234	GLY	-	expression tag	UNP P86993
B	235	ALA	-	expression tag	UNP P86993
B	236	ARG	-	expression tag	UNP P86993
B	237	HIS	-	expression tag	UNP P86993
B	238	HIS	-	expression tag	UNP P86993
B	239	HIS	-	expression tag	UNP P86993
B	240	HIS	-	expression tag	UNP P86993
B	241	HIS	-	expression tag	UNP P86993
B	242	HIS	-	expression tag	UNP P86993

- Molecule 2 is CALCIUM ION (three-letter code: CA) (formula: Ca).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	2	Total	Ca	0	0
			2	2		

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	2	Total	Ca		0	0
			2	2			

- Molecule 3 is N-ACETYL-D-GALACTOSAMINE (three-letter code: NGA) (formula: $C_8H_{15}NO_6$).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			15	8	1	6		
3	B	1	Total	C	N	O	0	0
			15	8	1	6		

- Molecule 4 is BETA-D-GALACTOSE (three-letter code: GAL) (formula: $C_6H_{12}O_6$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			10	6	4		
4	B	1	Total	C	O	0	0
			12	6	6		

- Molecule 5 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: $C_2H_6O_2$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			4	2	2		
5	A	1	Total	C	O	0	0
			4	2	2		

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	1
			8	4	4		
5	A	1	Total	C	O	0	0
			4	2	2		
5	A	1	Total	C	O	0	0
			4	2	2		
5	A	1	Total	C	O	0	1
			8	4	4		
5	A	1	Total	C	O	0	0
			4	2	2		
5	B	1	Total	C	O	0	0
			4	2	2		
5	B	1	Total	C	O	0	0
			4	2	2		
5	B	1	Total	C	O	0	0
			4	2	2		

- Molecule 6 is ALPHA D-GALACTOSE (three-letter code: GLA) (formula: $C_6H_{12}O_6$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	B	1	Total	C	O	0	0
			10	6	4		

- Molecule 7 is BETA-D-GLUCOSE (three-letter code: BGC) (formula: $C_6H_{12}O_6$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	B	1	Total	C	O	0	0
			11	6	5		

- Molecule 8 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	B	1	Total	Cl	0	0
			1	1		

- Molecule 9 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	A	209	Total	O	0	0
			209	209		
9	B	196	Total	O	0	1
			197	197		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Lectin



- Molecule 1: Lectin



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	44.68 Å 88.23 Å 110.78 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	39.86 – 1.43 39.86 – 1.43	Depositor EDS
% Data completeness (in resolution range)	93.3 (39.86-1.43) 93.3 (39.86-1.43)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.13 (at 1.43 Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.105 , 0.147 0.104 , 0.147	Depositor DCC
R_{free} test set	3713 reflections (5.17%)	DCC
Wilson B-factor (Å ²)	11.8	Xtriage
Anisotropy	0.717	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 47.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.98	EDS
Total number of atoms	4213	wwPDB-VP
Average B, all atoms (Å ²)	18.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.21% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: BGC, NGA, CL, GLA, CA, EDO, GAL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.03	5/1916 (0.3%)	1.03	5/2617 (0.2%)
1	B	1.00	3/1966 (0.2%)	1.18	11/2682 (0.4%)
All	All	1.01	8/3882 (0.2%)	1.11	16/5299 (0.3%)

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	14	GLU	CD-OE2	-6.54	1.18	1.25
1	A	14	GLU	CB-CG	-6.53	1.39	1.52
1	B	129	GLU	CB-CG	-5.75	1.41	1.52
1	A	129	GLU	CB-CG	-5.54	1.41	1.52
1	A	7	TYR	CZ-OH	5.27	1.46	1.37
1	B	4[A]	SER	CB-OG	-5.22	1.35	1.42
1	B	4[B]	SER	CB-OG	-5.22	1.35	1.42
1	A	158	ARG	CD-NE	-5.02	1.38	1.46

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	42[A]	ARG	NE-CZ-NH2	17.45	129.03	120.30
1	B	42[B]	ARG	NE-CZ-NH2	17.45	129.03	120.30
1	B	203[A]	ARG	NE-CZ-NH1	11.26	125.93	120.30
1	B	203[B]	ARG	NE-CZ-NH1	11.26	125.93	120.30
1	B	42[A]	ARG	NE-CZ-NH1	-10.55	115.03	120.30
1	B	42[B]	ARG	NE-CZ-NH1	-10.55	115.03	120.30
1	A	146	ARG	NE-CZ-NH1	9.00	124.80	120.30
1	A	158	ARG	NE-CZ-NH2	-7.43	116.58	120.30
1	A	146	ARG	NE-CZ-NH2	-5.93	117.33	120.30
1	B	203[A]	ARG	NE-CZ-NH2	-5.90	117.35	120.30
1	B	203[B]	ARG	NE-CZ-NH2	-5.90	117.35	120.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	42[A]	ARG	CD-NE-CZ	5.64	131.49	123.60
1	B	42[B]	ARG	CD-NE-CZ	5.64	131.49	123.60
1	A	35	ASP	CB-CG-OD1	5.31	123.08	118.30
1	A	25	ARG	NE-CZ-NH2	-5.18	117.71	120.30
1	B	129	GLU	CB-CA-C	-5.17	100.05	110.40

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1830	0	1737	23	0
1	B	1851	0	1771	16	1
2	A	2	0	0	0	0
2	B	2	0	0	0	0
3	A	15	0	14	0	0
3	B	15	0	14	0	0
4	A	10	0	8	0	0
4	B	12	0	10	0	0
5	A	36	0	53	16	0
5	B	12	0	16	2	0
6	B	10	0	8	0	0
7	B	11	0	10	0	0
8	B	1	0	0	0	0
9	A	209	0	0	8	0
9	B	197	0	0	11	0
All	All	4213	0	3641	41	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (41) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:113[A]:LYS:HG2	9:A:491:HOH:O	1.36	1.23
1:B:104[B]:ASN:OD1	9:B:401:HOH:O	1.60	1.17
1:A:104[B]:ASN:OD1	5:A:311:EDO:C2	1.99	1.11
1:A:26:ASN:HB3	9:A:497:HOH:O	1.58	1.03
1:A:104[B]:ASN:HD21	5:A:311:EDO:C2	1.75	1.00
1:A:104[B]:ASN:HD21	5:A:311:EDO:C1	1.73	0.99
1:A:104[B]:ASN:OD1	5:A:311:EDO:H21	1.60	0.99
1:A:104[B]:ASN:ND2	5:A:311:EDO:C2	2.26	0.99
1:A:104[B]:ASN:HD21	5:A:311:EDO:H11	1.34	0.92
1:A:104[B]:ASN:OD1	5:A:311:EDO:H22	1.70	0.91
1:A:104[B]:ASN:CG	5:A:311:EDO:C2	2.39	0.90
1:A:104[B]:ASN:ND2	5:A:311:EDO:H21	1.86	0.90
1:B:113[B]:LYS:HE3	9:B:426:HOH:O	1.71	0.89
1:A:104[B]:ASN:CG	5:A:311:EDO:H21	1.94	0.87
1:B:25:ARG:NE	9:B:402:HOH:O	2.09	0.82
1:A:104[B]:ASN:ND2	5:A:311:EDO:H11	1.95	0.81
1:A:104[B]:ASN:HD21	5:A:311:EDO:H21	1.43	0.77
1:A:104[B]:ASN:ND2	5:A:311:EDO:C1	2.46	0.77
1:A:104[B]:ASN:CG	5:A:311:EDO:H22	2.04	0.76
1:B:113[B]:LYS:CE	9:B:426:HOH:O	2.30	0.74
1:B:134:HIS:CE1	9:B:406:HOH:O	2.42	0.72
1:B:70:ARG:HD3	9:B:405:HOH:O	1.92	0.70
1:B:113[B]:LYS:HD3	1:B:197:ILE:HG23	1.75	0.69
1:B:134:HIS:HE1	9:B:406:HOH:O	1.76	0.66
1:B:25:ARG:HD3	1:B:30:GLN:OE1	1.97	0.64
1:B:146:ARG:HG3	1:B:188[A]:LEU:HD22	1.81	0.62
1:A:113[A]:LYS:CG	9:A:491:HOH:O	2.15	0.62
1:A:26:ASN:CB	9:A:497:HOH:O	2.30	0.60
5:B:309:EDO:H22	9:B:410:HOH:O	2.07	0.54
1:B:133:ARG:HA	9:B:470:HOH:O	2.09	0.53
1:A:113[A]:LYS:CE	9:A:491:HOH:O	2.58	0.52
1:A:113[A]:LYS:HE2	9:A:491:HOH:O	2.10	0.51
1:B:129:GLU:HB3	9:B:406:HOH:O	2.12	0.50
1:B:70:ARG:NH1	9:B:405:HOH:O	2.47	0.47
1:A:25:ARG:NH1	9:A:405:HOH:O	2.46	0.44
1:B:113[B]:LYS:HD3	1:B:197:ILE:CG2	2.45	0.44
1:B:69[A]:ILE:HG12	5:B:309:EDO:H11	1.99	0.44
1:A:25:ARG:HD3	5:A:307[B]:EDO:H21	2.01	0.43
5:A:307[B]:EDO:H12	9:A:422:HOH:O	2.19	0.42
1:A:113[B]:LYS:HE3	1:A:198:PHE:CD1	2.56	0.41
1:B:84[B]:ILE:HD13	1:B:84[B]:ILE:HG21	1.82	0.41

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the sym-

metry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:25:ARG:NH1	1:B:193:ASP:OD2[1_455]	2.08	0.12

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	234/242 (97%)	225 (96%)	9 (4%)	0	100	100
1	B	240/242 (99%)	232 (97%)	8 (3%)	0	100	100
All	All	474/484 (98%)	457 (96%)	17 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	205/209 (98%)	200 (98%)	5 (2%)	57	18
1	B	211/209 (101%)	211 (100%)	0	100	100
All	All	416/418 (100%)	411 (99%)	5 (1%)	82	48

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	15[A]	ASP

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	15[B]	ASP
1	A	73	SER
1	A	74	ASN
1	A	182	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	74	ASN
1	B	19	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 23 ligands modelled in this entry, 5 are monoatomic - leaving 18 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	NGA	A	303	4	15,15,15	0.61	0	17,21,21	0.95	1 (5%)
4	GAL	A	304	3	10,10,12	2.34	3 (30%)	10,13,17	5.24	6 (60%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	EDO	A	305	-	3,3,3	0.18	0	2,2,2	0.75	0
5	EDO	A	306	-	3,3,3	0.80	0	2,2,2	1.03	0
5	EDO	A	307[A]	-	3,3,3	0.56	0	2,2,2	0.66	0
5	EDO	A	307[B]	-	3,3,3	0.52	0	2,2,2	1.17	0
5	EDO	A	308	-	3,3,3	0.50	0	2,2,2	1.13	0
5	EDO	A	309	-	3,3,3	0.30	0	2,2,2	0.60	0
5	EDO	A	310[A]	-	3,3,3	0.25	0	2,2,2	0.38	0
5	EDO	A	310[B]	-	3,3,3	0.94	0	2,2,2	1.54	1 (50%)
5	EDO	A	311	-	3,3,3	0.75	0	2,2,2	1.14	0
3	NGA	B	303	6	15,15,15	1.30	2 (13%)	17,21,21	0.94	1 (5%)
6	GLA	B	304	3,4	10,10,12	1.32	2 (20%)	10,13,17	2.58	5 (50%)
4	GAL	B	305	7,6	12,12,12	0.71	0	17,17,17	1.29	2 (11%)
7	BGC	B	306	4	11,11,12	0.68	0	15,15,17	1.34	1 (6%)
5	EDO	B	307	-	3,3,3	0.68	0	2,2,2	0.34	0
5	EDO	B	308	-	3,3,3	0.08	0	2,2,2	0.87	0
5	EDO	B	309	-	3,3,3	2.49	1 (33%)	2,2,2	0.34	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NGA	A	303	4	-	0/6/26/26	0/1/1/1
4	GAL	A	304	3	-	0/2/15/22	0/1/1/1
5	EDO	A	305	-	-	0/1/1/1	0/0/0/0
5	EDO	A	306	-	-	0/1/1/1	0/0/0/0
5	EDO	A	307[A]	-	-	0/1/1/1	0/0/0/0
5	EDO	A	307[B]	-	-	0/1/1/1	0/0/0/0
5	EDO	A	308	-	-	0/1/1/1	0/0/0/0
5	EDO	A	309	-	-	0/1/1/1	0/0/0/0
5	EDO	A	310[A]	-	-	0/1/1/1	0/0/0/0
5	EDO	A	310[B]	-	-	0/1/1/1	0/0/0/0
5	EDO	A	311	-	-	0/1/1/1	0/0/0/0
3	NGA	B	303	6	-	0/6/26/26	0/1/1/1
6	GLA	B	304	3,4	-	0/2/15/22	0/1/1/1
4	GAL	B	305	7,6	-	0/2/22/22	0/1/1/1
7	BGC	B	306	4	-	0/2/18/22	0/1/1/1
5	EDO	B	307	-	-	0/1/1/1	0/0/0/0
5	EDO	B	308	-	-	0/1/1/1	0/0/0/0
5	EDO	B	309	-	-	0/1/1/1	0/0/0/0

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	B	309	EDO	O2-C2	-4.17	1.19	1.42
4	A	304	GAL	C3-C4	-2.63	1.48	1.52
6	B	304	GLA	C4-C5	2.29	1.56	1.52
3	B	303	NGA	C4-C3	2.37	1.58	1.52
6	B	304	GLA	C1-C2	2.41	1.53	1.51
3	B	303	NGA	C1-C2	3.76	1.57	1.53
4	A	304	GAL	C4-C5	4.53	1.60	1.52
4	A	304	GAL	C1-C2	4.65	1.55	1.51

All (17) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	304	GAL	O5-C1-C2	-11.69	97.65	110.85
4	A	304	GAL	O4-C4-C3	-8.43	89.45	110.02
6	B	304	GLA	O4-C4-C3	-4.29	99.55	110.02
4	A	304	GAL	C4-C3-C2	-3.91	106.02	111.54
4	A	304	GAL	O5-C5-C4	-3.22	104.22	109.70
6	B	304	GLA	O5-C5-C4	-2.98	104.63	109.70
6	B	304	GLA	O6-C6-C5	-2.94	101.47	111.30
4	B	305	GAL	C4-C3-C2	-2.68	105.84	110.79
5	A	310[B]	EDO	O1-C1-C2	-2.16	97.44	112.23
3	B	303	NGA	O5-C5-C6	2.21	112.12	106.38
3	A	303	NGA	O3-C3-C4	2.26	115.46	110.36
4	B	305	GAL	O5-C1-C2	2.54	114.44	110.00
4	A	304	GAL	O5-C5-C6	3.30	114.41	107.34
6	B	304	GLA	O4-C4-C5	3.31	116.28	109.70
6	B	304	GLA	O5-C5-C6	3.84	115.56	107.34
7	B	306	BGC	O5-C5-C6	3.99	111.56	106.67
4	A	304	GAL	O4-C4-C5	4.83	119.29	109.70

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 18 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	307[B]	EDO	2	0
5	A	311	EDO	14	0
5	B	309	EDO	2	0

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	229/242 (94%)	-0.55	3 (1%) 79 80	9, 12, 28, 56	0
1	B	229/242 (94%)	-0.56	1 (0%) 93 95	9, 14, 29, 58	0
All	All	458/484 (94%)	-0.55	4 (0%) 85 86	9, 13, 29, 58	0

All (4) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	229	ASN	3.0
1	B	229	ASN	2.8
1	A	57	ARG	2.1
1	A	37	ASN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
5	EDO	A	306	4/4	0.89	0.17	13.58	36,40,49,51	0
5	EDO	A	311	4/4	0.96	0.14	13.19	19,31,32,32	0
5	EDO	A	310[B]	4/4	0.92	0.15	8.20	13,14,15,17	4
5	EDO	A	310[A]	4/4	0.92	0.15	8.20	14,16,20,20	4
5	EDO	B	308	4/4	0.87	0.11	4.33	24,29,30,33	0
5	EDO	A	307[B]	4/4	0.92	0.17	3.02	20,20,23,26	4
5	EDO	A	307[A]	4/4	0.92	0.17	3.02	24,26,30,34	4
5	EDO	A	305	4/4	0.94	0.08	2.52	25,27,29,30	0
5	EDO	A	308	4/4	0.97	0.11	2.32	19,28,29,42	0
5	EDO	B	309	4/4	0.98	0.09	2.03	18,18,20,25	0
5	EDO	B	307	4/4	0.97	0.11	1.49	21,25,32,35	0
2	CA	A	302	1/1	1.00	0.05	0.21	10,10,10,10	0
3	NGA	A	303	15/15	0.98	0.04	-0.52	10,11,16,18	0
2	CA	B	302	1/1	1.00	0.05	-0.62	11,11,11,11	0
3	NGA	B	303	15/15	0.99	0.04	-1.12	11,13,18,19	0
2	CA	B	301	1/1	1.00	0.04	-1.65	12,12,12,12	0
2	CA	A	301	1/1	1.00	0.03	-2.28	12,12,12,12	0
7	BGC	B	306	11/12	0.88	0.20	-	33,36,45,52	0
4	GAL	A	304	10/12	0.84	0.19	-	28,36,48,71	0
5	EDO	A	309	4/4	0.95	0.17	-	36,38,40,41	0
4	GAL	B	305	12/12	0.94	0.19	-	28,32,35,53	0
8	CL	B	310	1/1	1.00	0.03	-	17,17,17,17	0
6	GLA	B	304	10/12	0.92	0.15	-	25,30,53,74	0

6.5 Other polymers ⓘ

There are no such residues in this entry.