



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 01:39 PM GMT

PDB ID : 3UGF
Title : Crystal structure of a 6-SST/6-SFT from *Pachysandra terminalis*
Authors : Lammens, W.; Rabijs, A.; Van Laere, A.; Strelkov, S.V.; Van den Ende, W.
Deposited on : 2011-11-02
Resolution : 1.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	3190 (1.70-1.70)
Clashscore	102246	3585 (1.70-1.70)
Ramachandran outliers	100387	3527 (1.70-1.70)
Sidechain outliers	100360	3527 (1.70-1.70)
RSRZ outliers	91569	3200 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	546	<div> <div style="width: 100%; height: 10px; background-color: red; position: relative;"> % </div> <div style="width: 95%; height: 10px; background-color: green; position: relative;"> 95% </div> </div>
1	B	546	<div> <div style="width: 100%; height: 10px; background-color: red; position: relative;"> 2% </div> <div style="width: 95%; height: 10px; background-color: green; position: relative;"> 95% </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	NAG	A	547	-	-	-	X
3	NAG	B	550	-	-	-	X
6	GOL	A	560	-	-	-	X
6	GOL	A	562	-	-	-	X
6	GOL	A	564	-	-	-	X
6	GOL	B	565	-	-	-	X
6	GOL	B	566	-	-	-	X
6	GOL	B	570	-	-	-	X
9	NAG	B	562	-	-	-	X

2 Entry composition [i](#)

There are 10 unique types of molecules in this entry. The entry contains 9809 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Sucrose:(Sucrose/fructan) 6-fructosyltransferase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	524	Total	C	N	O	S	0	0	0
			4164	2687	679	784	14			
1	B	530	Total	C	N	O	S	0	0	0
			4203	2709	685	795	14			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	133	VAL	ALA	SEE REMARK 999	UNP E3PQS3
A	361	GLY	ARG	SEE REMARK 999	UNP E3PQS3
B	133	VAL	ALA	SEE REMARK 999	UNP E3PQS3
B	361	GLY	ARG	SEE REMARK 999	UNP E3PQS3

- Molecule 2 is a polymer of unknown type called SUGAR (2-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	2	Total	C	N	O	0	0
			24	14	1	9		

- Molecule 3 is a polymer of unknown type called SUGAR (3-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	3	Total	C	N	O	0	0
			38	22	2	14		
3	B	3	Total	C	N	O	0	0
			38	22	2	14		

- Molecule 4 is a polymer of unknown type called SUGAR (3-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	3	Total	C	N	O	0	0
			39	22	2	15		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	B	3	Total	C	N	O	0	0
			39	22	2	15		

- Molecule 5 is a polymer of unknown type called SUGAR (4-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	4	Total	C	N	O	0	0
			49	28	2	19		

- Molecule 6 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			6	3	3		
6	A	1	Total	C	O	0	0
			6	3	3		
6	A	1	Total	C	O	0	0
			6	3	3		
6	A	1	Total	C	O	0	0
			6	3	3		
6	A	1	Total	C	O	0	0
			6	3	3		
6	B	1	Total	C	O	0	0
			6	3	3		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	B	1	Total	C	O	0	0
			6	3	3		
6	B	1	Total	C	O	0	0
			6	3	3		
6	B	1	Total	C	O	0	0
			6	3	3		
6	B	1	Total	C	O	0	0
			6	3	3		
6	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 7 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	1	Total	O	S	0	0
			5	4	1		
7	B	1	Total	O	S	0	0
			5	4	1		

- Molecule 8 is a polymer of unknown type called SUGAR (5-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
8	B	5	Total	C	N	O	0	0
			60	34	2	24		

- Molecule 9 is a polymer of unknown type called SUGAR (6-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
9	B	6	Total	C	N	O	0	0
			74	42	3	29		

- Molecule 10 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
10	A	478	Total	O	0	0
			478	478		
10	B	515	Total	O	0	0
			515	515		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Sucrose:(Sucrose/fructan) 6-fructosyltransferase



- Molecule 1: Sucrose:(Sucrose/fructan) 6-fructosyltransferase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	89.08Å 125.27Å 148.95Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.45 – 1.70 48.45 – 1.70	Depositor EDS
% Data completeness (in resolution range)	100.0 (48.45-1.70) 100.0 (48.45-1.70)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.10	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.70 (at 1.70Å)	Xtriage
Refinement program	REFMAC 5.5.0102	Depositor
R, R_{free}	0.171 , 0.186 0.185 , 0.200	Depositor DCC
R_{free} test set	9166 reflections (5.28%)	DCC
Wilson B-factor (Å ²)	17.6	Xtriage
Anisotropy	0.212	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 42.5	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Outliers	0 of 182827 reflections	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	9809	wwPDB-VP
Average B, all atoms (Å ²)	20.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.81% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, BMA, NAG, SO4, FUC, MAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.42	0/4287	0.53	0/5841
1	B	0.43	0/4327	0.55	0/5898
All	All	0.43	0/8614	0.54	0/11739

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4164	0	4021	6	0
1	B	4203	0	4053	5	0
2	A	24	0	22	0	0
3	A	38	0	34	1	0
3	B	38	0	34	0	0
4	A	39	0	34	0	0
4	B	39	0	34	0	0
5	A	49	0	43	0	0
6	A	36	0	48	0	0
6	B	42	0	56	2	0
7	A	5	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
7	B	5	0	0	0	0
8	B	60	0	52	0	0
9	B	74	0	64	0	0
10	A	478	0	0	1	0
10	B	515	0	0	0	0
All	All	9809	0	8495	11	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (11) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:35:ASP:OD1	1:B:93:SER:HA	1.80	0.80
1:A:260:THR:HG23	1:A:262:LYS:H	1.53	0.73
1:A:257:ASP:HB3	1:A:260:THR:HG22	1.72	0.71
1:A:260:THR:HG21	10:A:633:HOH:O	1.97	0.65
1:B:543:PRO:HB3	6:B:565:GOL:H31	1.78	0.65
1:A:177:TYR:HB3	3:A:549:NAG:H82	1.87	0.55
1:A:62:TRP:CH2	1:A:94:GLY:HA3	2.42	0.54
1:A:62:TRP:HH2	1:A:94:GLY:HA3	1.80	0.47
1:B:76:LEU:HA	6:B:565:GOL:H12	1.98	0.44
1:B:346:GLU:HG2	1:B:349:ARG:CZ	2.49	0.43
1:B:382:PHE:HE2	1:B:480:MET:HG2	1.85	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	518/546 (95%)	498 (96%)	18 (4%)	2 (0%)	39 20

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	526/546 (96%)	505 (96%)	19 (4%)	2 (0%)	39	20
All	All	1044/1092 (96%)	1003 (96%)	37 (4%)	4 (0%)	39	20

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	441	ASP
1	A	93	SER
1	B	93	SER
1	A	29	SER

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	447/463 (96%)	447 (100%)	0	100	100
1	B	451/463 (97%)	449 (100%)	2 (0%)	93	90
All	All	898/926 (97%)	896 (100%)	2 (0%)	95	92

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	29	SER
1	B	91	VAL

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

29 carbohydrates are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	NAG	A	547	1,2	14,14,15	0.52	0	15,19,21	0.80	0
2	FUC	A	548	2	10,10,11	0.48	0	14,14,16	1.09	2 (14%)
3	NAG	A	549	1,3	14,14,15	0.45	0	15,19,21	0.96	1 (6%)
3	NAG	A	550	3	14,14,15	0.51	0	15,19,21	0.68	0
3	FUC	A	551	3	10,10,11	0.52	0	14,14,16	0.72	1 (7%)
4	NAG	A	552	1,4	14,14,15	0.55	0	15,19,21	0.78	0
4	NAG	A	553	4	14,14,15	0.56	0	15,19,21	0.77	0
4	BMA	A	554	4	11,11,12	0.49	0	14,15,17	0.80	1 (7%)
5	NAG	A	555	1,5	14,14,15	0.44	0	15,19,21	0.94	1 (6%)
5	NAG	A	556	5	14,14,15	0.48	0	15,19,21	0.76	1 (6%)
5	BMA	A	557	5	11,11,12	0.60	0	14,15,17	0.74	1 (7%)
5	FUC	A	558	5	10,10,11	0.45	0	14,14,16	0.90	1 (7%)
4	NAG	B	547	1,4	14,14,15	0.50	0	15,19,21	0.72	0
4	NAG	B	548	4	14,14,15	0.51	0	15,19,21	0.84	0
4	BMA	B	549	4	11,11,12	0.55	0	14,15,17	0.69	0
3	NAG	B	550	1,3	14,14,15	0.55	0	15,19,21	0.76	0
3	NAG	B	551	3	14,14,15	0.48	0	15,19,21	0.80	0
3	FUC	B	552	3	10,10,11	0.47	0	14,14,16	0.98	2 (14%)
8	NAG	B	553	1,8	14,14,15	0.47	0	15,19,21	0.98	1 (6%)
8	NAG	B	554	8	14,14,15	0.49	0	15,19,21	0.89	1 (6%)
8	BMA	B	555	8	11,11,12	0.57	0	14,15,17	0.81	1 (7%)
8	MAN	B	556	8	11,11,12	0.51	0	14,15,17	1.17	1 (7%)
8	FUC	B	557	8	10,10,11	0.48	0	14,14,16	0.78	0
9	NAG	B	558	1,9	14,14,15	0.47	0	15,19,21	1.15	2 (13%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
9	NAG	B	559	9	14,14,15	0.49	0	15,19,21	0.70	0
9	BMA	B	560	9	11,11,12	0.68	0	14,15,17	0.85	0
9	MAN	B	561	9	11,11,12	0.51	0	14,15,17	1.09	1 (7%)
9	NAG	B	562	9	14,14,15	0.49	0	15,19,21	0.82	0
9	FUC	B	563	9	10,10,11	0.46	0	14,14,16	0.89	1 (7%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	A	547	1,2	-	0/6/23/26	0/1/1/1
2	FUC	A	548	2	-	0/0/17/20	0/1/1/1
3	NAG	A	549	1,3	-	0/6/23/26	0/1/1/1
3	NAG	A	550	3	-	0/6/23/26	0/1/1/1
3	FUC	A	551	3	-	0/0/17/20	0/1/1/1
4	NAG	A	552	1,4	-	0/6/23/26	0/1/1/1
4	NAG	A	553	4	-	0/6/23/26	0/1/1/1
4	BMA	A	554	4	-	0/2/19/22	0/1/1/1
5	NAG	A	555	1,5	-	0/6/23/26	0/1/1/1
5	NAG	A	556	5	-	0/6/23/26	0/1/1/1
5	BMA	A	557	5	-	0/2/19/22	0/1/1/1
5	FUC	A	558	5	-	0/0/17/20	0/1/1/1
4	NAG	B	547	1,4	-	0/6/23/26	0/1/1/1
4	NAG	B	548	4	-	0/6/23/26	0/1/1/1
4	BMA	B	549	4	-	0/2/19/22	0/1/1/1
3	NAG	B	550	1,3	-	0/6/23/26	0/1/1/1
3	NAG	B	551	3	-	0/6/23/26	0/1/1/1
3	FUC	B	552	3	-	0/0/17/20	0/1/1/1
8	NAG	B	553	1,8	-	0/6/23/26	0/1/1/1
8	NAG	B	554	8	-	0/6/23/26	0/1/1/1
8	BMA	B	555	8	-	0/2/19/22	0/1/1/1
8	MAN	B	556	8	-	0/2/19/22	0/1/1/1
8	FUC	B	557	8	-	0/0/17/20	0/1/1/1
9	NAG	B	558	1,9	-	0/6/23/26	0/1/1/1
9	NAG	B	559	9	-	0/6/23/26	0/1/1/1
9	BMA	B	560	9	-	0/2/19/22	0/1/1/1
9	MAN	B	561	9	-	0/2/19/22	0/1/1/1
9	NAG	B	562	9	-	0/6/23/26	0/1/1/1
9	FUC	B	563	9	-	0/0/17/20	0/1/1/1

There are no bond length outliers.

All (19) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	556	NAG	C1-O5-C5	2.01	114.80	112.25
3	B	552	FUC	O5-C5-C6	2.03	109.49	106.13
3	A	551	FUC	O5-C5-C6	2.04	109.50	106.13
9	B	558	NAG	C2-N2-C7	2.05	125.67	123.04
5	A	558	FUC	O5-C5-C6	2.12	109.64	106.13
5	A	557	BMA	C1-O5-C5	2.15	114.98	112.25
8	B	554	NAG	C1-O5-C5	2.16	114.99	112.25
9	B	563	FUC	O5-C5-C6	2.29	109.91	106.13
4	A	554	BMA	C1-O5-C5	2.30	115.16	112.25
2	A	548	FUC	C1-C2-C3	2.33	112.30	109.54
2	A	548	FUC	O5-C5-C6	2.52	110.30	106.13
3	B	552	FUC	C1-O5-C5	2.56	116.34	112.38
8	B	555	BMA	C1-O5-C5	2.58	115.53	112.25
8	B	553	NAG	C1-O5-C5	2.91	115.95	112.25
9	B	558	NAG	C1-O5-C5	2.92	115.95	112.25
5	A	555	NAG	C1-O5-C5	2.92	115.96	112.25
3	A	549	NAG	C1-O5-C5	3.03	116.10	112.25
9	B	561	MAN	C1-O5-C5	3.36	116.51	112.25
8	B	556	MAN	C1-O5-C5	3.38	116.54	112.25

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	549	NAG	1	0

5.6 Ligand geometry ⓘ

15 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	GOL	A	559	-	5,5,5	0.29	0	5,5,5	0.31	0
6	GOL	A	560	-	5,5,5	0.33	0	5,5,5	0.49	0
6	GOL	A	561	-	5,5,5	0.36	0	5,5,5	0.18	0
6	GOL	A	562	-	5,5,5	0.35	0	5,5,5	0.25	0
6	GOL	A	563	-	5,5,5	0.30	0	5,5,5	0.21	0
6	GOL	A	564	-	5,5,5	0.36	0	5,5,5	0.24	0
7	SO4	A	565	-	4,4,4	0.24	0	6,6,6	0.12	0
6	GOL	B	564	-	5,5,5	0.32	0	5,5,5	0.32	0
6	GOL	B	565	-	5,5,5	0.35	0	5,5,5	0.42	0
6	GOL	B	566	-	5,5,5	0.37	0	5,5,5	0.25	0
6	GOL	B	567	-	5,5,5	0.36	0	5,5,5	0.30	0
6	GOL	B	568	-	5,5,5	0.36	0	5,5,5	0.39	0
6	GOL	B	569	-	5,5,5	0.29	0	5,5,5	0.29	0
6	GOL	B	570	-	5,5,5	0.34	0	5,5,5	0.26	0
7	SO4	B	571	-	4,4,4	0.25	0	6,6,6	0.08	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	GOL	A	559	-	-	0/4/4/4	0/0/0/0
6	GOL	A	560	-	-	0/4/4/4	0/0/0/0
6	GOL	A	561	-	-	0/4/4/4	0/0/0/0
6	GOL	A	562	-	-	0/4/4/4	0/0/0/0
6	GOL	A	563	-	-	0/4/4/4	0/0/0/0
6	GOL	A	564	-	-	0/4/4/4	0/0/0/0
7	SO4	A	565	-	-	0/0/0/0	0/0/0/0
6	GOL	B	564	-	-	0/4/4/4	0/0/0/0
6	GOL	B	565	-	-	0/4/4/4	0/0/0/0
6	GOL	B	566	-	-	0/4/4/4	0/0/0/0
6	GOL	B	567	-	-	0/4/4/4	0/0/0/0
6	GOL	B	568	-	-	0/4/4/4	0/0/0/0
6	GOL	B	569	-	-	0/4/4/4	0/0/0/0
6	GOL	B	570	-	-	0/4/4/4	0/0/0/0
7	SO4	B	571	-	-	0/0/0/0	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	B	565	GOL	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	524/546 (95%)	-0.13	8 (1%) 76 80	12, 19, 28, 37	0
1	B	530/546 (97%)	-0.08	12 (2%) 64 68	11, 18, 28, 44	0
All	All	1054/1092 (96%)	-0.10	20 (1%) 70 74	11, 18, 28, 44	0

All (20) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	440	THR	10.1
1	B	233	PRO	6.5
1	B	441	ASP	5.7
1	B	234	LYS	4.6
1	B	442	GLY	4.5
1	A	233	PRO	4.2
1	A	386	LYS	3.9
1	B	222	GLY	3.3
1	B	443	ASN	3.2
1	A	7	PRO	3.2
1	B	474	ASP	3.1
1	A	438	LYS	2.8
1	B	439	GLY	2.6
1	B	387	GLU	2.4
1	A	443	ASN	2.2
1	A	234	LYS	2.2
1	B	541	ILE	2.1
1	B	488	ILE	2.1
1	A	545	HIS	2.1
1	A	424	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
9	NAG	B	562	14/15	0.66	0.32	13.93	47,48,49,50	0
3	NAG	B	550	14/15	0.82	0.15	9.74	25,30,37,40	0
2	NAG	A	547	14/15	0.86	0.17	4.51	26,29,34,35	0
4	NAG	A	552	14/15	0.93	0.08	0.22	19,21,23,26	0
4	NAG	B	547	14/15	0.97	0.07	-0.32	16,19,21,24	0
4	BMA	B	549	11/12	0.68	0.40	-	43,46,47,47	0
4	NAG	B	548	14/15	0.88	0.20	-	28,32,34,38	0
8	NAG	B	554	14/15	0.94	0.10	-	20,23,28,29	0
4	NAG	A	553	14/15	0.90	0.17	-	25,28,31,35	0
5	FUC	A	558	10/11	0.93	0.11	-	35,36,37,37	0
5	NAG	A	556	14/15	0.90	0.10	-	35,36,40,40	0
9	NAG	B	559	14/15	0.89	0.12	-	29,31,34,35	0
8	NAG	B	553	14/15	0.94	0.10	-	18,21,22,24	0
2	FUC	A	548	10/11	0.73	0.33	-	39,41,42,42	0
9	BMA	B	560	11/12	0.83	0.18	-	38,40,41,43	0
3	NAG	A	550	14/15	0.88	0.13	-	19,24,26,28	0
9	FUC	B	563	10/11	0.93	0.12	-	29,30,31,32	0
9	NAG	B	558	14/15	0.93	0.09	-	23,27,28,29	0
3	FUC	B	552	10/11	0.79	0.26	-	40,42,42,43	0
9	MAN	B	561	11/12	0.70	0.28	-	45,46,47,47	0
8	FUC	B	557	10/11	0.96	0.07	-	20,21,22,23	0
3	FUC	A	551	10/11	0.92	0.08	-	23,24,26,26	0
5	BMA	A	557	11/12	0.65	0.26	-	41,42,43,44	0
3	NAG	A	549	14/15	0.93	0.10	-	18,22,23,26	0
3	NAG	B	551	14/15	0.68	0.37	-	43,46,47,47	0
8	MAN	B	556	11/12	0.78	0.32	-	40,41,42,44	0
5	NAG	A	555	14/15	0.96	0.09	-	27,30,33,33	0
4	BMA	A	554	11/12	0.62	0.18	-	38,41,42,43	0
8	BMA	B	555	11/12	0.89	0.23	-	33,35,37,39	0

6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
6	GOL	B	565	6/6	0.56	0.32	10.82	46,47,47,47	0
6	GOL	B	570	6/6	0.76	0.24	8.01	55,55,55,55	0
6	GOL	B	566	6/6	0.73	0.29	7.61	39,42,42,43	0
6	GOL	A	564	6/6	0.88	0.14	6.89	52,52,52,52	0
6	GOL	A	562	6/6	0.88	0.13	3.93	49,49,49,49	0
6	GOL	A	560	6/6	0.86	0.17	3.09	36,37,38,38	0
6	GOL	A	563	6/6	0.81	0.17	1.98	37,37,38,38	0
6	GOL	B	569	6/6	0.96	0.08	-0.13	17,19,21,23	0
6	GOL	A	559	6/6	0.93	0.08	-0.56	18,20,22,24	0
7	SO4	A	565	5/5	0.93	0.17	-	53,53,53,53	0
6	GOL	A	561	6/6	0.50	0.31	-	44,45,46,47	0
7	SO4	B	571	5/5	0.94	0.09	-	49,49,50,50	0
6	GOL	B	564	6/6	0.83	0.17	-	43,43,44,44	0
6	GOL	B	567	6/6	0.86	0.12	-	48,48,48,48	0
6	GOL	B	568	6/6	0.78	0.20	-	38,39,39,40	0

6.5 Other polymers

There are no such residues in this entry.