



wwPDB X-ray Structure Validation Summary Report ⓘ

Apr 7, 2016 – 04:58 PM EDT

PDB ID : 4UIQ
Title : Isolated globin domain of the Bordetella pertussis globin-coupled sensor with a heme at the dimer interface
Authors : Germani, F.; Donne, J.; De Schutter, A.; Pesce, A.; Berghmans, H.; Van Hauwaert, M.-L.; Moens, L.; Van Doorslaer, S.; Bolognesi, M.; Nardini, M.; Dewilde, S.
Deposited on : 2015-03-31
Resolution : 1.55 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.1 (RC1), CSD as537be (2016)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20027107
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20027107


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.55 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1665 (1.58-1.54)
Clashscore	102246	1014 (1.56-1.56)
Ramachandran outliers	100387	1704 (1.58-1.54)
Sidechain outliers	100360	1702 (1.58-1.54)
RSRZ outliers	91569	1668 (1.58-1.54)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	177	<div> <div>5%</div> <div> <div></div> <div>76%</div> <div>10%</div> <div>•</div> <div>13%</div> </div> </div>
1	B	177	<div> <div>2%</div> <div> <div></div> <div>75%</div> <div>10%</div> <div>••</div> <div>14%</div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	HEM	B	201	-	-	-	X
3	GOL	A	1159	-	-	X	-
3	GOL	A	1160	-	-	-	X

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 2855 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called GLOBIN-COUPLED SENSOR WITH DIGUANYLATE CYCLASE ACTIVITY.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	154	Total	C	N	O	S	0	9	0
			1252	797	215	231	9			
1	B	153	Total	C	N	O	S	0	7	0
			1235	782	218	226	9			

There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	170	LEU	-	EXPRESSION TAG	UNP Q7VTL8
A	171	GLU	-	EXPRESSION TAG	UNP Q7VTL8
A	172	HIS	-	EXPRESSION TAG	UNP Q7VTL8
A	173	HIS	-	EXPRESSION TAG	UNP Q7VTL8
A	174	HIS	-	EXPRESSION TAG	UNP Q7VTL8
A	175	HIS	-	EXPRESSION TAG	UNP Q7VTL8
A	176	HIS	-	EXPRESSION TAG	UNP Q7VTL8
A	177	HIS	-	EXPRESSION TAG	UNP Q7VTL8
A	16	SER	CYS	ENGINEERED MUTATION	UNP Q7VTL8
A	45	SER	CYS	ENGINEERED MUTATION	UNP Q7VTL8
A	114	SER	CYS	ENGINEERED MUTATION	UNP Q7VTL8
A	154	SER	CYS	ENGINEERED MUTATION	UNP Q7VTL8
B	170	LEU	-	EXPRESSION TAG	UNP Q7VTL8
B	171	GLU	-	EXPRESSION TAG	UNP Q7VTL8
B	172	HIS	-	EXPRESSION TAG	UNP Q7VTL8
B	173	HIS	-	EXPRESSION TAG	UNP Q7VTL8
B	174	HIS	-	EXPRESSION TAG	UNP Q7VTL8
B	175	HIS	-	EXPRESSION TAG	UNP Q7VTL8
B	176	HIS	-	EXPRESSION TAG	UNP Q7VTL8
B	177	HIS	-	EXPRESSION TAG	UNP Q7VTL8
B	16	SER	CYS	ENGINEERED MUTATION	UNP Q7VTL8
B	45	SER	CYS	ENGINEERED MUTATION	UNP Q7VTL8
B	114	SER	CYS	ENGINEERED MUTATION	UNP Q7VTL8
B	154	SER	CYS	ENGINEERED MUTATION	UNP Q7VTL8

- Molecule 2 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula: $C_{34}H_{32}FeN_4O_4$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	Fe	N	O	0	1
			44	35	1	4	4		
2	A	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		
2	B	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		
2	B	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		

- Molecule 4 is IMIDAZOLE (three-letter code: IMD) (formula: $C_3H_5N_2$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	C	N	0	0
			5	3	2		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	82	Total 82	O 82	0	0
5	B	89	Total 96	O 96	0	7

• Molecule 1: GLOBIN-COUPLED SENSOR WITH DIGUANYLATE CYCLASE ACTIVITY



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	69.96Å 53.11Å 81.17Å 90.00° 93.23° 90.00°	Depositor
Resolution (Å)	40.52 – 1.55 40.52 – 1.55	Depositor EDS
% Data completeness (in resolution range)	97.2 (40.52-1.55) 97.4 (40.52-1.55)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.30 (at 1.55Å)	Xtriage
Refinement program	PHENIX (PHENIX.REFINE)	Depositor
R, R_{free}	0.156 , 0.204 0.167 , 0.211	Depositor DCC
R_{free} test set	2116 reflections (5.03%)	DCC
Wilson B-factor (Å ²)	14.0	Xtriage
Anisotropy	0.702	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 45.1	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	3 of 42164 reflections (0.007%)	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	2855	wwPDB-VP
Average B, all atoms (Å ²)	24.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 9.50% of the height of the origin peak. No significant pseudotranslation is detected.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, IMD, HEM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.73	0/1306	0.84	1/1776 (0.1%)
1	B	0.74	0/1284	0.84	3/1746 (0.2%)
All	All	0.73	0/2590	0.84	4/3522 (0.1%)

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	5	PRO	CA-N-CD	-9.70	97.92	111.50
1	B	49	ASP	CB-CG-OD2	-6.32	112.61	118.30
1	B	119	ARG	NE-CZ-NH1	-5.49	117.56	120.30
1	B	89	VAL	CG1-CB-CG2	5.22	119.25	110.90

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1252	0	1256	20	1
1	B	1235	0	1232	16	0
2	A	87	0	36	5	0
2	B	86	0	60	9	0
3	A	12	0	16	5	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	B	5	0	4	0	0
5	A	82	0	0	2	1
5	B	96	0	0	2	0
All	All	2855	0	2604	45	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

The worst 5 of 45 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:126:ARG:HG3	5:A:2068:HOH:O	1.70	0.90
1:B:154:SER:HB3	2:B:201:HEM:HMB3	1.63	0.80
2:B:201:HEM:HMA1	2:B:201:HEM:HBA1	1.69	0.75
1:A:122:GLU:HG2	3:A:1159:GOL:O2	1.88	0.72
1:A:122:GLU:HA	3:A:1159:GOL:H2	1.72	0.71

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:15:THR:OG1	5:A:2064:HOH:O[2_556]	2.18	0.02

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	161/177 (91%)	158 (98%)	3 (2%)	0	100	100
1	B	158/177 (89%)	156 (99%)	2 (1%)	0	100	100
All	All	319/354 (90%)	314 (98%)	5 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	133/145 (92%)	130 (98%)	3 (2%)	58	25
1	B	131/145 (90%)	130 (99%)	1 (1%)	86	71
All	All	264/290 (91%)	260 (98%)	4 (2%)	78	44

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	73[A]	LEU
1	A	73[B]	LEU
1	A	91	PHE
1	B	89	VAL

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	GOL	A	1159	-	5,5,5	0.45	0	5,5,5	1.49	0
3	GOL	A	1160	-	5,5,5	0.37	0	5,5,5	0.30	0
2	HEM	A	200[A]	-	24,50,50	2.35	8 (33%)	16,82,82	2.04	6 (37%)
2	HEM	A	200[B]	-	24,50,50	2.35	8 (33%)	16,82,82	2.04	6 (37%)
2	HEM	A	201	1	24,50,50	2.27	6 (25%)	16,82,82	1.86	5 (31%)
4	IMD	B	1156	2	3,5,5	0.59	0	4,5,5	1.62	1 (25%)
2	HEM	B	200	1,4	24,50,50	2.54	6 (25%)	16,82,82	1.44	2 (12%)
2	HEM	B	201	1	24,50,50	2.30	7 (29%)	16,82,82	2.33	7 (43%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	A	1159	-	-	0/4/4/4	0/0/0/0
3	GOL	A	1160	-	-	0/4/4/4	0/0/0/0
2	HEM	A	200[A]	-	-	0/6/54/54	0/0/8/8
2	HEM	A	200[B]	-	-	0/6/54/54	0/0/8/8
2	HEM	A	201	1	-	0/6/54/54	0/0/8/8
4	IMD	B	1156	2	-	0/0/0/0	0/1/1/1
2	HEM	B	200	1,4	-	0/6/54/54	0/0/8/8
2	HEM	B	201	1	-	1/6/54/54	0/0/8/8

The worst 5 of 35 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	200	HEM	C3B-C2B	-6.66	1.31	1.40
2	A	201	HEM	C3B-C2B	-5.01	1.34	1.40
2	B	200	HEM	C3C-C2C	-4.93	1.34	1.40

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	200[A]	HEM	C3C-C2C	-4.85	1.34	1.40
2	A	200[B]	HEM	C3C-C2C	-4.85	1.34	1.40

The worst 5 of 27 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	201	HEM	CMA-C3A-C4A	-5.46	119.02	128.31
2	B	200	HEM	CBD-CAD-C3D	-3.92	105.59	112.47
2	A	200[A]	HEM	CMD-C2D-C1D	-3.51	122.35	128.31
2	A	200[B]	HEM	CMD-C2D-C1D	-3.51	122.35	128.31
2	A	201	HEM	CMA-C3A-C4A	-3.47	122.41	128.31

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	201	HEM	C1A-C2A-CAA-CBA

There are no ring outliers.

6 monomers are involved in 19 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	1159	GOL	5	0
2	A	200[A]	HEM	1	0
2	A	200[B]	HEM	1	0
2	A	201	HEM	3	0
2	B	200	HEM	4	0
2	B	201	HEM	5	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	154/177 (87%)	0.19	9 (5%) 26 27	14, 23, 34, 49	0
1	B	153/177 (86%)	-0.05	4 (2%) 59 63	14, 21, 33, 48	1 (0%)
All	All	307/354 (86%)	0.08	13 (4%) 40 41	14, 22, 33, 49	1 (0%)

The worst 5 of 13 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	5	PRO	5.9
1	A	6	GLU	5.1
1	B	6	GLU	3.9
1	A	157	TYR	3.7
1	A	126	ARG	3.6

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	HEM	B	201	43/43	0.87	0.31	6.08	37,41,44,45	43
3	GOL	A	1160	6/6	0.66	0.27	5.39	51,53,53,53	0
2	HEM	B	200	43/43	0.98	0.09	0.36	14,17,28,33	0
2	HEM	A	201	43/43	0.97	0.12	-0.03	26,28,35,37	43
2	HEM	A	200[B]	43/43	0.97	0.10	-0.14	16,19,32,34	1
2	HEM	A	200[A]	43/43	0.97	0.10	-0.24	16,19,32,34	1
4	IMD	B	1156	5/5	0.97	0.07	-1.15	18,19,19,19	0
3	GOL	A	1159	6/6	0.59	0.41	-	45,46,47,48	0

6.5 Other polymers [i](#)

There are no such residues in this entry.