



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 01:49 PM GMT

PDB ID : 3V3V
Title : Structural and functional analysis of quercetagenin, a natural JNK1 inhibitor
Authors : Baek, S.; Kang, N.J.; Popowicz, G.M.; Arciniega, M.; Jung, S.K.; Byun, S.;
Song, N.R.; Heo, Y.S.; Kim, B.Y.; Lee, H.J.; Holak, T.A.; Augustin, M.; Bode,
A.M.; Huber, R.; Dong, Z.; Lee, K.W.
Deposited on : 2011-12-14
Resolution : 2.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	2103 (2.70-2.70)
Clashscore	102246	2422 (2.70-2.70)
Ramachandran outliers	100387	2382 (2.70-2.70)
Sidechain outliers	100360	2382 (2.70-2.70)
RSRZ outliers	91569	2107 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	379	<div> <div></div> <div>58% 31% 7%</div> </div>
2	B	11	<div> <div></div> <div>55% 36% 9%</div> </div>

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 2952 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Mitogen-activated protein kinase 8.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	351	Total	C	N	O	S	0	0	0
			2782	1783	466	514	19			


There are 17 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	185	GLU	TYR	CONFLICT	UNP P45983
A	288	ALA	LYS	CONFLICT	UNP P45983
A	364	PRO	GLU	CONFLICT	UNP P45983
A	365	LYS	GLU	CONFLICT	UNP P45983
A	367	PRO	-	EXPRESSION TAG	UNP P45983
A	368	THR	-	EXPRESSION TAG	UNP P45983
A	369	THR	-	EXPRESSION TAG	UNP P45983
A	370	LEU	-	EXPRESSION TAG	UNP P45983
A	371	ASN	-	EXPRESSION TAG	UNP P45983
A	372	LEU	-	EXPRESSION TAG	UNP P45983
A	373	PHE	-	EXPRESSION TAG	UNP P45983
A	374	HIS	-	EXPRESSION TAG	UNP P45983
A	375	HIS	-	EXPRESSION TAG	UNP P45983
A	376	HIS	-	EXPRESSION TAG	UNP P45983
A	377	HIS	-	EXPRESSION TAG	UNP P45983
A	378	HIS	-	EXPRESSION TAG	UNP P45983
A	379	HIS	-	EXPRESSION TAG	UNP P45983

- Molecule 2 is a protein called C-Jun-amino-terminal kinase-interacting protein 1.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	B	10	Total	C	N	O	0	0	0
			84	55	15	14			

- Molecule 3 is 3,5,6,7-TETRAHYDROXY-2-(3,4-DIHYDROXYPHENYL)-4H-CHROMEN-4-ONE (three-letter code: MYU) (formula: C₁₅H₁₀O₈).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			23	15	8		

- Molecule 4 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	O	S	0	0
			5	4	1		
4	A	1	Total	O	S	0	0
			5	4	1		
4	A	1	Total	O	S	0	0
			5	4	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	O	S	0	0
			5	4	1		
4	A	1	Total	O	S	0	0
			5	4	1		
4	A	1	Total	O	S	0	0
			5	4	1		
4	B	1	Total	O	S	0	0
			5	4	1		

- Molecule 5 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total	Cl	0	0
			1	1		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	23	Total	O	0	0
			23	23		
6	B	4	Total	O	0	0
			4	4		

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Mitogen-activated protein kinase 8



- Molecule 2: C-Jun-amino-terminal kinase-interacting protein 1



4 Data and refinement statistics

Property	Value	Source
Space group	I 4 2 2	Depositor
Cell constants a, b, c, α , β , γ	172.37Å 172.37Å 86.04Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 2.70 43.09 – 2.50	Depositor EDS
% Data completeness (in resolution range)	77.0 (20.00-2.70) 67.5 (43.09-2.50)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.48 (at 2.51Å)	Xtriage
Refinement program	REFMAC 5.5.0072	Depositor
R, R_{free}	0.227 , 0.266 0.220 , 0.256	Depositor DCC
R_{free} test set	612 reflections (4.61%)	DCC
Wilson B-factor (Å ²)	37.5	Xtriage
Anisotropy	0.054	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.27 , 43.8	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.44$, $\langle L^2 \rangle = 0.27$	Xtriage
Outliers	0 of 15327 reflections	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	2952	wwPDB-VP
Average B, all atoms (Å ²)	64.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.50% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MYU, SO4, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.62	0/2842	0.69	0/3848
2	B	0.60	0/86	0.70	0/114
All	All	0.62	0/2928	0.69	0/3962

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2782	0	2758	73	1
2	B	84	0	91	2	0
3	A	23	0	9	2	0
4	A	30	0	0	0	1
4	B	5	0	0	0	0
5	A	1	0	0	0	0
6	A	23	0	0	2	0
6	B	4	0	0	1	0
All	All	2952	0	2858	75	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 13.

All (75) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:253:GLN:OE1	1:A:253:GLN:HA	1.72	0.88
1:A:90:ASN:HD22	1:A:91:VAL:N	1.85	0.75
1:A:189:ARG:HG3	1:A:189:ARG:HH11	1.56	0.71
1:A:263:ARG:HE	1:A:263:ARG:HA	1.54	0.71
1:A:187:VAL:O	1:A:192:ARG:NH2	2.24	0.69
1:A:38:GLY:HA3	1:A:55:LYS:HE2	1.76	0.67
1:A:348:THR:HG23	1:A:351:GLU:H	1.58	0.66
1:A:29:LEU:H	1:A:29:LEU:HD12	1.61	0.66
1:A:90:ASN:HD22	1:A:91:VAL:H	1.43	0.64
1:A:25:ARG:HG3	1:A:47:ILE:HG12	1.80	0.63
1:A:97:SER:HA	1:A:357:TYR:OH	1.98	0.62
1:A:57:LEU:O	1:A:60:PRO:HD3	2.00	0.62
1:A:193:ALA:HA	1:A:209:TRP:CD1	2.34	0.62
1:A:224:LEU:O	1:A:226:PRO:HD3	2.00	0.61
1:A:54:ILE:HG12	1:A:107:VAL:HG22	1.84	0.58
2:B:154:PRO:N	6:B:304:HOH:O	2.36	0.58
3:A:401:MYU:H5	3:A:401:MYU:O19	2.04	0.58
1:A:259:TYR:CZ	1:A:263:ARG:HD2	2.40	0.57
1:A:134:GLN:HE22	1:A:164:THR:HA	1.70	0.56
1:A:271:PHE:HA	1:A:274:LEU:HD12	1.89	0.55
1:A:12:SER:OG	1:A:21:THR:HG22	2.07	0.55
1:A:153:LYS:NZ	1:A:188:THR:HG21	2.22	0.55
1:A:63:ASN:HB2	1:A:66:HIS:HB2	1.88	0.54
1:A:338:PRO:HB2	1:A:341:GLN:OE1	2.06	0.54
1:A:348:THR:HG22	1:A:351:GLU:CD	2.28	0.54
1:A:238:ILE:HA	6:A:516:HOH:O	2.07	0.54
1:A:253:GLN:OE1	1:A:253:GLN:CA	2.50	0.53
1:A:271:PHE:CZ	1:A:299:SER:HA	2.43	0.53
1:A:140:LYS:HB2	1:A:312:VAL:HG11	1.91	0.53
1:A:59:ARG:CD	1:A:102:GLN:HE21	2.23	0.52
1:A:295:ARG:HA	1:A:298:LEU:HD12	1.91	0.52
1:A:81:ASN:HD22	1:A:81:ASN:C	2.14	0.51
1:A:350:GLU:OE2	1:A:350:GLU:HA	2.08	0.51
1:A:119:ILE:HG22	1:A:221:GLY:HA2	1.93	0.51
1:A:190:TYR:HA	1:A:233:GLN:NE2	2.26	0.51
1:A:162:ASP:OD1	1:A:162:ASP:N	2.44	0.50
1:A:32:ILE:HG13	1:A:33:GLY:H	1.77	0.50
1:A:127:ARG:HH21	2:B:159:THR:HG22	1.76	0.50
1:A:59:ARG:HD2	1:A:102:GLN:HE21	1.78	0.49
1:A:132:LEU:O	1:A:136:LEU:HG	2.12	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:294:ALA:O	1:A:297:LEU:N	2.45	0.49
1:A:145:ALA:HB1	1:A:337:ILE:HD11	1.95	0.49
1:A:154:PRO:HD3	1:A:214:ILE:HG12	1.95	0.49
1:A:56:LYS:NZ	6:A:502:HOH:O	2.46	0.49
1:A:193:ALA:O	1:A:197:ILE:HG13	2.13	0.48
1:A:71:TYR:O	1:A:72:ARG:C	2.53	0.47
1:A:32:ILE:HG13	1:A:33:GLY:N	2.29	0.47
1:A:194:PRO:HA	1:A:197:ILE:HD12	1.96	0.46
1:A:342:LEU:O	1:A:343:ASP:C	2.54	0.46
1:A:168:LEU:HD22	3:A:401:MYU:C5	2.45	0.46
1:A:253:GLN:O	1:A:257:ARG:HB2	2.17	0.45
1:A:13:VAL:HG21	1:A:29:LEU:HD13	1.98	0.45
1:A:194:PRO:HG2	1:A:304:ILE:HA	1.98	0.45
1:A:56:LYS:HB2	1:A:105:TYR:CE2	2.52	0.45
1:A:153:LYS:O	1:A:154:PRO:C	2.53	0.44
1:A:95:GLN:HG2	1:A:100:GLU:O	2.17	0.44
1:A:348:THR:HG22	1:A:351:GLU:CG	2.47	0.44
1:A:139:ILE:HD11	1:A:152:LEU:HD13	1.99	0.43
1:A:153:LYS:HZ2	1:A:188:THR:HG21	1.83	0.43
1:A:150:ARG:HD3	1:A:202:TYR:CZ	2.54	0.43
1:A:224:LEU:HD23	1:A:224:LEU:HA	1.80	0.42
1:A:150:ARG:HD3	1:A:202:TYR:CE1	2.55	0.42
1:A:86:ILE:HA	1:A:166:LYS:HD3	2.02	0.42
1:A:149:HIS:O	1:A:150:ARG:HB2	2.19	0.42
1:A:244:PRO:HB3	1:A:248:PHE:CD2	2.55	0.41
1:A:217:GLU:HG3	1:A:222:GLY:O	2.20	0.41
1:A:10:PHE:CE2	1:A:94:PRO:HA	2.55	0.41
1:A:118:VAL:O	1:A:121:MET:HB2	2.21	0.41
1:A:283:ASP:HB3	1:A:290:LYS:HG3	2.01	0.41
1:A:327:PRO:HA	1:A:330:ALA:HB3	2.03	0.41
1:A:106:ILE:CG2	1:A:108:MET:HE3	2.51	0.41
1:A:189:ARG:HG3	1:A:189:ARG:NH1	2.30	0.40
1:A:249:MET:SD	1:A:260:VAL:HG12	2.61	0.40
1:A:111:MET:CE	1:A:160:LYS:HB2	2.51	0.40
1:A:14:GLU:OE1	1:A:14:GLU:HA	2.20	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:189:ARG:NH1	4:A:405:SO4:O1[5_555]	2.15	0.05

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	345/379 (91%)	292 (85%)	43 (12%)	10 (3%)	6	14
2	B	8/11 (73%)	8 (100%)	0	0	100	100
All	All	353/390 (90%)	300 (85%)	43 (12%)	10 (3%)	6	15

All (10) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	281	PRO
1	A	343	ASP
1	A	34	SER
1	A	284	SER
1	A	36	ALA
1	A	169	ASP
1	A	319	PRO
1	A	8	ASN
1	A	268	GLY
1	A	349	ILE

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	302/340 (89%)	268 (89%)	34 (11%)	7	16
2	B	10/11 (91%)	8 (80%)	2 (20%)	1	4
All	All	312/351 (89%)	276 (88%)	36 (12%)	7	16

All (36) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	8	ASN
1	A	23	LEU
1	A	62	GLN
1	A	65	THR
1	A	81	ASN
1	A	90	ASN
1	A	97	SER
1	A	100	GLU
1	A	103	ASP
1	A	124	ASP
1	A	129	SER
1	A	142	LEU
1	A	143	HIS
1	A	158	VAL
1	A	161	SER
1	A	178	THR
1	A	188	THR
1	A	189	ARG
1	A	198	LEU
1	A	204	GLU
1	A	224	LEU
1	A	241	LEU
1	A	243	THR
1	A	253	GLN
1	A	255	THR
1	A	257	ARG
1	A	263	ARG
1	A	289	LEU
1	A	290	LYS
1	A	337	ILE
1	A	339	ASP
1	A	340	LYS
1	A	349	ILE
1	A	356	ILE
2	B	155	LYS
2	B	162	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	62	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	63	ASN
1	A	81	ASN
1	A	84	ASN
1	A	90	ASN
1	A	102	GLN
1	A	117	GLN
1	A	120	GLN
1	A	134	GLN
1	A	143	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 9 ligands modelled in this entry, 1 is monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	MYU	A	401	-	21,25,25	1.14	2 (9%)	28,38,38	1.52	5 (17%)
4	SO4	A	402	-	4,4,4	0.17	0	6,6,6	0.19	0
4	SO4	A	403	-	4,4,4	0.18	0	6,6,6	0.11	0
4	SO4	A	404	-	4,4,4	0.17	0	6,6,6	0.47	0
4	SO4	A	405	-	4,4,4	0.22	0	6,6,6	0.52	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	SO4	A	406	-	4,4,4	0.18	0	6,6,6	0.20	0
4	SO4	A	407	-	4,4,4	0.17	0	6,6,6	0.25	0
4	SO4	B	201	-	4,4,4	0.18	0	6,6,6	0.13	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	MYU	A	401	-	-	0/4/4/4	0/3/3/3
4	SO4	A	402	-	-	0/0/0/0	0/0/0/0
4	SO4	A	403	-	-	0/0/0/0	0/0/0/0
4	SO4	A	404	-	-	0/0/0/0	0/0/0/0
4	SO4	A	405	-	-	0/0/0/0	0/0/0/0
4	SO4	A	406	-	-	0/0/0/0	0/0/0/0
4	SO4	A	407	-	-	0/0/0/0	0/0/0/0
4	SO4	B	201	-	-	0/0/0/0	0/0/0/0

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	401	MYU	C6-C7	-3.47	1.41	1.46
3	A	401	MYU	C16-C15	2.51	1.39	1.37

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	401	MYU	C1-C2-C3	-2.77	117.65	120.49
3	A	401	MYU	C16-C15-C14	-2.56	118.78	120.41
3	A	401	MYU	C12-C11-C10	-2.09	118.22	121.25
3	A	401	MYU	C11-C12-C7	2.05	122.94	119.41
3	A	401	MYU	C6-C7-C12	3.94	127.51	120.04

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	401	MYU	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	405	SO4	0	1

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	351/379 (92%)	-0.39	2 (0%) 90 91	43, 60, 93, 100	0
2	B	10/11 (90%)	-0.22	0 100 100	55, 59, 61, 63	0
All	All	361/390 (92%)	-0.39	2 (0%) 90 91	43, 60, 92, 100	0

All (2) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	363	LEU	3.0
1	A	99	GLU	2.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
4	SO4	A	403	5/5	0.95	0.16	-0.56	132,132,132,132	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	MYU	A	401	23/23	0.94	0.12	-1.04	63,66,67,70	0
4	SO4	A	405	5/5	0.99	0.09	-1.52	38,38,41,43	0
4	SO4	A	404	5/5	0.98	0.09	-1.53	92,92,93,93	0
4	SO4	A	406	5/5	0.82	0.23	-	161,162,162,162	0
4	SO4	A	402	5/5	0.96	0.12	-	93,94,94,95	0
5	CL	A	408	1/1	0.94	0.08	-	67,67,67,67	0
4	SO4	B	201	5/5	0.89	0.17	-	120,120,121,121	0
4	SO4	A	407	5/5	0.96	0.12	-	70,70,70,70	5

6.5 Other polymers

There are no such residues in this entry.