



# wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 05:56 AM GMT

PDB ID : 2VD9  
Title : The crystal structure of alanine racemase from Bacillus anthracis (BA0252) with bound L-Ala-P  
Authors : Au, K.; Ren, J.; Walter, T.S.; Harlos, K.; Nettleship, J.E.; Owens, R.J.; Stuart, D.I.; Esnouf, R.M.; Oxford Protein Production Facility (OPPF); Structural Proteomics in Europe (SPINE)  
Deposited on : 2007-10-01  
Resolution : 2.10 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	3939 (2.10-2.10)
Clashscore	102246	4460 (2.10-2.10)
Ramachandran outliers	100387	4413 (2.10-2.10)
Sidechain outliers	100360	4414 (2.10-2.10)
RSRZ outliers	91569	3948 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	391	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 1%, orange 1%, yellow 1%, green 98%);"></div> <div style="display: flex; justify-content: space-between; width: 90%; margin: 0 auto;"> <span>%</span> <span>92%</span> <span>7% .</span> </div> </div>
1	B	391	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 0%, orange 0%, yellow 0%, green 90%, grey 10%);"></div> <div style="display: flex; justify-content: space-between; width: 90%; margin: 0 auto;"> <span></span> <span>90%</span> <span>8% ..</span> </div> </div>

## 2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 7161 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ALANINE RACEMASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	386	Total	C	N	O	S	0	0	0
			3098	2021	511	560	6			
1	B	386	Total	C	N	O	S	0	0	0
			3098	2021	511	560	6			


- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total	Mg	0	0
			1	1		
2	A	2	Total	Mg	0	0
			2	2		

- Molecule 3 is CHLORIDE ION (three-letter code: CL) (formula: Cl).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total	Cl	0	0
			1	1		
3	A	2	Total	Cl	0	0
			2	2		

- Molecule 4 is {1-[(3-HYDROXY-METHYL-5-PHOSPHONOXY-METHYL-PYRIDIN-4-YLMETHYL)-AMINO]-ETHYL}-PHOSPHONIC ACID (three-letter code: IN5) (formula: C<sub>10</sub>H<sub>18</sub>N<sub>2</sub>O<sub>8</sub>P<sub>2</sub>).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O	P	0	0
			22	10	2	8	2		
4	B	1	Total	C	N	O	P	0	0
			22	10	2	8	2		

- Molecule 5 is (1S)-1-[(1E)-{3-HYDROXY-2-METHYL-5-[(PHOSPHONOOXY)METHYL]PYRIDIN-4-YL}METHYLENE)AMINO]ETHYLPHOSPHONIC ACID (three-letter code: EPC) (formula: C<sub>10</sub>H<sub>16</sub>N<sub>2</sub>O<sub>8</sub>P<sub>2</sub>).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	A	1	Total	C	N	O	P	0	0
			22	10	2	8	2		

Continued on next page...

*Continued from previous page...*

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	B	1	Total	C	N	O	P	0	0
			22	10	2	8	2		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	424	Total	O	0	0
			424	424		
6	B	447	Total	O	0	0
			447	447		


### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: ALANINE RACEMASE



#### • Molecule 1: ALANINE RACEMASE



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	59.69 Å 96.50 Å 140.66 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	45.64 – 2.10 45.64 – 2.10	Depositor EDS
% Data completeness (in resolution range)	99.7 (45.64-2.10) 99.7 (45.64-2.10)	Depositor EDS
$R_{merge}$	0.23	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.47 (at 2.10 Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, $R_{free}$	0.188 , 0.239 0.191 , 0.244	Depositor DCC
$R_{free}$ test set	2434 reflections (5.34%)	DCC
Wilson B-factor (Å <sup>2</sup> )	19.1	Xtriage
Anisotropy	0.093	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 56.0	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	1 of 48030 reflections (0.002%)	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	7161	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	19.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 35.50 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 5.7772e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup> Intensities estimated from amplitudes.

<sup>2</sup> Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: IN5, MG, MLY, EPC, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.40	0/2973	0.58	0/4065
1	B	0.39	0/2973	0.58	0/4065
All	All	0.40	0/5946	0.58	0/8130

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3098	0	3119	20	0
1	B	3098	0	3121	33	0
2	A	2	0	0	0	0
2	B	1	0	0	0	0
3	A	2	0	0	0	0
3	B	1	0	0	0	0
4	A	22	0	7	4	0
4	B	22	0	8	2	0
5	A	22	0	6	5	0
5	B	22	0	6	0	0
6	A	424	0	0	1	0

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	B	447	0	0	4	0
All	All	7161	0	6267	54	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 54 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:208:ASN:O	1:B:212:THR:HG23	1.70	0.91
1:A:4:ALA:HB1	1:A:5:PRO:HD2	1.71	0.73
1:B:86:VAL:O	1:B:106:THR:HG23	1.90	0.70
1:B:4:ALA:CB	1:B:5:PRO:CD	2.69	0.70
1:B:86:VAL:O	1:B:106:THR:CG2	2.41	0.67

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	366/391 (94%)	353 (96%)	12 (3%)	1 (0%)	46	45
1	B	366/391 (94%)	352 (96%)	13 (4%)	1 (0%)	46	45
All	All	732/782 (94%)	705 (96%)	25 (3%)	2 (0%)	46	45

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	48	ASP
1	B	48	ASP

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	308/312 (99%)	306 (99%)	2 (1%)	90	94
1	B	308/312 (99%)	302 (98%)	6 (2%)	65	70
All	All	616/624 (99%)	608 (99%)	8 (1%)	76	82

5 of 8 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	106	THR
1	B	279	GLU
1	B	177	THR
1	B	95	ILE
1	B	176	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

36 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
1	MLY	A	110	1	9,10,11	0.43	0	9,11,13	1.85	3 (33%)
1	MLY	A	118	1	9,10,11	0.46	0	9,11,13	2.01	5 (55%)
1	MLY	A	127	1	9,10,11	0.54	0	9,11,13	1.87	3 (33%)
1	MLY	A	145	1	9,10,11	0.37	0	9,11,13	1.81	3 (33%)
1	MLY	A	148	1	9,10,11	0.44	0	9,11,13	1.84	3 (33%)
1	MLY	A	152	1	9,10,11	0.42	0	9,11,13	1.99	5 (55%)
1	MLY	A	182	1	9,10,11	0.41	0	9,11,13	2.07	5 (55%)
1	MLY	A	195	1	9,10,11	0.49	0	9,11,13	2.10	4 (44%)
1	MLY	A	202	1	9,10,11	0.45	0	9,11,13	1.90	2 (22%)
1	MLY	A	245	1	9,10,11	0.47	0	9,11,13	1.80	3 (33%)
1	MLY	A	255	1	9,10,11	0.46	0	9,11,13	1.79	2 (22%)
1	MLY	A	264	1	9,10,11	0.40	0	9,11,13	1.97	4 (44%)
1	MLY	A	27	1	9,10,11	0.41	0	9,11,13	1.88	3 (33%)
1	MLY	A	277	1	9,10,11	0.34	0	9,11,13	2.12	5 (55%)
1	MLY	A	307	1	9,10,11	0.35	0	9,11,13	1.92	3 (33%)
1	MLY	A	333	1	9,10,11	0.62	0	9,11,13	2.10	3 (33%)
1	MLY	A	378	1	9,10,11	0.41	0	9,11,13	1.85	2 (22%)
1	MLY	A	54	1	9,10,11	0.40	0	9,11,13	1.90	3 (33%)
1	MLY	B	110	1	9,10,11	0.47	0	9,11,13	1.93	3 (33%)
1	MLY	B	118	1	9,10,11	0.35	0	9,11,13	1.90	4 (44%)
1	MLY	B	127	1	9,10,11	0.50	0	9,11,13	1.79	2 (22%)
1	MLY	B	145	1	9,10,11	0.45	0	9,11,13	1.91	4 (44%)
1	MLY	B	148	1	9,10,11	0.37	0	9,11,13	2.01	5 (55%)
1	MLY	B	152	1	9,10,11	0.38	0	9,11,13	1.97	5 (55%)
1	MLY	B	182	1	9,10,11	0.45	0	9,11,13	1.93	4 (44%)
1	MLY	B	195	1	9,10,11	0.37	0	9,11,13	1.99	4 (44%)
1	MLY	B	202	1	9,10,11	0.31	0	9,11,13	2.01	4 (44%)
1	MLY	B	245	1	9,10,11	0.46	0	9,11,13	2.02	4 (44%)
1	MLY	B	255	1	9,10,11	0.44	0	9,11,13	1.77	2 (22%)
1	MLY	B	264	1	9,10,11	0.56	0	9,11,13	2.46	3 (33%)
1	MLY	B	27	1	9,10,11	0.43	0	9,11,13	2.11	5 (55%)
1	MLY	B	277	1	9,10,11	0.43	0	9,11,13	1.88	2 (22%)
1	MLY	B	307	1	9,10,11	0.45	0	9,11,13	1.78	3 (33%)
1	MLY	B	333	1	9,10,11	0.47	0	9,11,13	1.99	4 (44%)
1	MLY	B	378	1	9,10,11	0.37	0	9,11,13	2.08	4 (44%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
1	MLY	B	54	1	9,10,11	0.42	0	9,11,13	1.85	3 (33%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	MLY	A	110	1	-	0/7/9/11	0/0/0/0
1	MLY	A	118	1	-	0/7/9/11	0/0/0/0
1	MLY	A	127	1	-	0/7/9/11	0/0/0/0
1	MLY	A	145	1	-	0/7/9/11	0/0/0/0
1	MLY	A	148	1	-	0/7/9/11	0/0/0/0
1	MLY	A	152	1	-	0/7/9/11	0/0/0/0
1	MLY	A	182	1	-	0/7/9/11	0/0/0/0
1	MLY	A	195	1	-	0/7/9/11	0/0/0/0
1	MLY	A	202	1	-	0/7/9/11	0/0/0/0
1	MLY	A	245	1	-	0/7/9/11	0/0/0/0
1	MLY	A	255	1	-	0/7/9/11	0/0/0/0
1	MLY	A	264	1	-	0/7/9/11	0/0/0/0
1	MLY	A	27	1	-	0/7/9/11	0/0/0/0
1	MLY	A	277	1	-	0/7/9/11	0/0/0/0
1	MLY	A	307	1	-	0/7/9/11	0/0/0/0
1	MLY	A	333	1	-	0/7/9/11	0/0/0/0
1	MLY	A	378	1	-	0/7/9/11	0/0/0/0
1	MLY	A	54	1	-	0/7/9/11	0/0/0/0
1	MLY	B	110	1	-	0/7/9/11	0/0/0/0
1	MLY	B	118	1	-	0/7/9/11	0/0/0/0
1	MLY	B	127	1	-	0/7/9/11	0/0/0/0
1	MLY	B	145	1	-	0/7/9/11	0/0/0/0
1	MLY	B	148	1	-	0/7/9/11	0/0/0/0
1	MLY	B	152	1	-	0/7/9/11	0/0/0/0
1	MLY	B	182	1	-	0/7/9/11	0/0/0/0
1	MLY	B	195	1	-	0/7/9/11	0/0/0/0
1	MLY	B	202	1	-	0/7/9/11	0/0/0/0
1	MLY	B	245	1	-	0/7/9/11	0/0/0/0
1	MLY	B	255	1	-	0/7/9/11	0/0/0/0
1	MLY	B	264	1	-	0/7/9/11	0/0/0/0
1	MLY	B	27	1	-	0/7/9/11	0/0/0/0
1	MLY	B	277	1	-	0/7/9/11	0/0/0/0
1	MLY	B	307	1	-	0/7/9/11	0/0/0/0
1	MLY	B	333	1	-	0/7/9/11	0/0/0/0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	MLY	B	378	1	-	0/7/9/11	0/0/0/0
1	MLY	B	54	1	-	0/7/9/11	0/0/0/0

There are no bond length outliers.

The worst 5 of 126 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	264	MLY	CD-CE-NZ	-4.18	103.16	113.92
1	B	378	MLY	CD-CE-NZ	-2.98	106.26	113.92
1	A	195	MLY	CD-CE-NZ	-2.86	106.56	113.92
1	A	182	MLY	CD-CE-NZ	-2.77	106.80	113.92
1	B	333	MLY	CD-CE-NZ	-2.72	106.91	113.92

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	277	MLY	1	0
1	B	145	MLY	1	0
1	B	152	MLY	1	0
1	B	195	MLY	2	0

## 5.5 Carbohydrates

There are no carbohydrates in this entry.

## 5.6 Ligand geometry

Of 10 ligands modelled in this entry, 6 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	IN5	A	1394	-	20,22,22	3.23	6 (30%)	26,33,33	1.54	4 (15%)
5	EPC	A	1395	-	22,22,22	2.29	6 (27%)	25,33,33	1.45	2 (8%)
4	IN5	B	1392	-	20,22,22	3.32	6 (30%)	26,33,33	1.45	4 (15%)
5	EPC	B	1393	-	22,22,22	2.28	7 (31%)	25,33,33	1.33	2 (8%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	IN5	A	1394	-	-	0/17/17/17	0/1/1/1
5	EPC	A	1395	-	-	0/15/17/17	0/1/1/1
4	IN5	B	1392	-	-	1/17/17/17	0/1/1/1
5	EPC	B	1393	-	-	0/15/17/17	0/1/1/1

The worst 5 of 25 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	1392	IN5	C8-C4	-10.57	1.39	1.51
4	A	1394	IN5	C8-C4	-10.09	1.40	1.51
4	A	1394	IN5	P2-O7	-4.62	1.47	1.54
4	B	1392	IN5	C8-N2	-4.51	1.32	1.46
4	A	1394	IN5	C8-N2	-4.42	1.32	1.46

The worst 5 of 12 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	1395	EPC	O8-P2-O7	-3.07	105.32	113.49
5	B	1393	EPC	O8-P2-O7	-3.07	105.33	113.49
4	B	1392	IN5	O7-P2-O6	-2.46	106.95	113.49
4	A	1394	IN5	O8-P2-O6	-2.42	107.06	113.49
4	A	1394	IN5	C8-C4-C5	-2.19	117.75	119.71

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	B	1392	IN5	P2-C9-N2-C8

There are no ring outliers.

3 monomers are involved in 10 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	1394	IN5	4	0
5	A	1395	EPC	5	0
4	B	1392	IN5	2	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	368/391 (94%)	-0.37	3 (0%) 87 90	10, 17, 26, 29	0
1	B	368/391 (94%)	-0.44	0 100 100	11, 17, 25, 30	0
All	All	736/782 (94%)	-0.40	3 (0%) 93 94	10, 17, 25, 30	0

All (3) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	124	SER	3.0
1	A	4	ALA	2.8
1	A	342	ASP	2.5

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
1	MLY	A	195	11/12	0.83	0.21	-	25,28,33,33	0
1	MLY	B	118	11/12	0.85	0.28	-	24,26,32,32	0
1	MLY	B	148	11/12	0.90	0.17	-	19,22,27,28	0
1	MLY	A	202	11/12	0.89	0.19	-	23,25,29,29	0
1	MLY	A	127	11/12	0.87	0.16	-	18,21,25,26	0
1	MLY	B	255	11/12	0.95	0.11	-	12,13,15,16	0
1	MLY	A	27	11/12	0.81	0.18	-	23,26,30,30	0
1	MLY	A	245	11/12	0.84	0.24	-	22,25,31,31	0
1	MLY	B	264	11/12	0.94	0.11	-	20,21,23,23	0

*Continued on next page...*



*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
1	MLY	B	378	11/12	0.83	0.24	-	25,28,34,34	0
1	MLY	B	202	11/12	0.86	0.17	-	22,24,29,29	0
1	MLY	A	255	11/12	0.95	0.10	-	12,13,17,17	0
1	MLY	A	333	11/12	0.91	0.12	-	15,18,22,22	0
1	MLY	B	127	11/12	0.87	0.16	-	17,19,24,24	0
1	MLY	B	195	11/12	0.71	0.29	-	24,27,33,33	0
1	MLY	B	54	11/12	0.93	0.15	-	19,21,27,28	0
1	MLY	A	182	11/12	0.82	0.28	-	20,23,29,29	0
1	MLY	A	152	11/12	0.80	0.20	-	21,24,29,29	0
1	MLY	A	110	11/12	0.90	0.19	-	18,20,26,26	0
1	MLY	B	245	11/12	0.87	0.19	-	22,25,31,31	0
1	MLY	B	145	11/12	0.77	0.25	-	20,24,30,30	0
1	MLY	B	333	11/12	0.92	0.11	-	16,18,22,22	0
1	MLY	A	277	11/12	0.74	0.36	-	25,28,33,33	0
1	MLY	A	118	11/12	0.85	0.25	-	25,27,32,32	0
1	MLY	B	110	11/12	0.87	0.18	-	16,20,26,27	0
1	MLY	B	182	11/12	0.85	0.23	-	18,22,27,28	0
1	MLY	A	378	11/12	0.81	0.28	-	27,29,33,34	0
1	MLY	B	27	11/12	0.91	0.17	-	23,25,30,30	0
1	MLY	B	307	11/12	0.83	0.24	-	20,22,27,27	0
1	MLY	A	54	11/12	0.86	0.17	-	19,22,27,27	0
1	MLY	A	148	11/12	0.89	0.20	-	20,22,27,28	0
1	MLY	B	277	11/12	0.82	0.21	-	24,26,30,31	0
1	MLY	B	152	11/12	0.85	0.18	-	21,23,29,29	0
1	MLY	A	145	11/12	0.91	0.13	-	20,23,28,28	0
1	MLY	A	307	11/12	0.92	0.15	-	20,22,26,27	0
1	MLY	A	264	11/12	0.89	0.18	-	21,23,27,27	0

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
5	EPC	A	1395	22/22	0.96	0.13	1.39	11,12,14,14	22
4	IN5	B	1392	22/22	0.96	0.12	1.21	10,13,16,17	22
5	EPC	B	1393	22/22	0.97	0.12	1.14	11,12,13,14	22
4	IN5	A	1394	22/22	0.96	0.12	1.14	10,11,12,13	22
3	CL	B	1391	1/1	0.99	0.07	-1.29	11,11,11,11	0
2	MG	B	1390	1/1	0.57	0.28	-	81,81,81,81	0
3	CL	A	1392	1/1	0.99	0.04	-	15,15,15,15	0
3	CL	A	1393	1/1	0.90	0.11	-	53,53,53,53	0
2	MG	A	1390	1/1	0.99	0.10	-	24,24,24,24	0
2	MG	A	1391	1/1	0.24	0.41	-	73,73,73,73	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.