



# Full wwPDB X-ray Structure Validation Report ⓘ

Jan 31, 2016 – 11:28 PM GMT

PDB ID : 1XCC  
Title : 1-Cys peroxidoxin from Plasmodium Yoelli  
Authors : Vedadi, M.; Sharma, S.; Houston, S.; Lew, J.; Wasney, G.; Amani, M.; Xu, X.;  
Bray, J.; Sundstrom, M.; Arrowsmith, C.; Edwards, A.; Hui, R.; Bochkarev,  
A.; Structural Genomics Consortium (SGC)  
Deposited on : 2004-09-01  
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	3852 (2.30-2.30)
Clashscore	102246	4452 (2.30-2.30)
Ramachandran outliers	100387	4410 (2.30-2.30)
Sidechain outliers	100360	4409 (2.30-2.30)
RSRZ outliers	91569	3857 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	220	<div> <div>2%</div> <div> <div></div> <div>85%</div> <div>13%</div> <div>.</div> </div> </div>
1	B	220	<div> <div>3%</div> <div> <div></div> <div>83%</div> <div>12%</div> <div>..</div> </div> </div>
1	C	220	<div> <div>%</div> <div> <div></div> <div>90%</div> <div>7%</div> <div>..</div> </div> </div>
1	D	220	<div> <div></div> <div> <div></div> <div>85%</div> <div>13%</div> <div>..</div> </div> </div>

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 7362 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 1-Cys peroxiredoxin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	219	Total	C	N	O	S	0	0	0
			1769	1140	289	329	11			
1	B	218	Total	C	N	O	S	0	0	0
			1765	1138	288	328	11			
1	C	218	Total	C	N	O	S	0	0	0
			1765	1138	288	328	11			
1	D	216	Total	C	N	O	S	0	0	0
			1743	1123	284	325	11			


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	92	Total	O	0	0
			92	92		
2	B	63	Total	O	0	0
			63	63		
2	C	94	Total	O	0	0
			94	94		
2	D	71	Total	O	0	0
			71	71		



### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



- Molecule 1: 1-Cys peroxiredoxin




- Molecule 1: 1-Cys peroxiredoxin



- Molecule 1: 1-Cys peroxiredoxin



- Molecule 1: 1-Cys peroxiredoxin



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	90.39Å 156.84Å 178.07Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 2.30 89.04 – 2.30	Depositor EDS
% Data completeness (in resolution range)	99.7 (20.00-2.30) 99.6 (89.04-2.30)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.06	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	4.24 (at 2.29Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, $R_{free}$	0.183 , 0.226 0.181 , 0.223	Depositor DCC
$R_{free}$ test set	2863 reflections (5.36%)	DCC
Wilson B-factor (Å <sup>2</sup> )	36.0	Xtriage
Anisotropy	0.361	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 40.1	EDS
Estimated twinning fraction	0.015 for 1/2*h-1/2*k,-3/2*h-1/2*k,-l 0.024 for 1/2*h+1/2*k,3/2*h-1/2*k,-l	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 56471 reflections	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	7362	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	37.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.89% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	1.05	0/1812	0.86	1/2451 (0.0%)
1	B	0.95	1/1808 (0.1%)	0.86	2/2446 (0.1%)
1	C	1.04	2/1808 (0.1%)	0.91	2/2446 (0.1%)
1	D	1.05	1/1784 (0.1%)	0.83	2/2413 (0.1%)
All	All	1.02	4/7212 (0.1%)	0.87	7/9756 (0.1%)

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	184	CYS	CB-SG	-13.68	1.58	1.82
1	B	184	CYS	CB-SG	-5.88	1.72	1.81
1	C	184	CYS	CB-SG	-5.17	1.73	1.81
1	C	103	GLU	CD-OE1	5.03	1.31	1.25

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	114	MET	CG-SD-CE	-10.17	83.94	100.20
1	C	105	ARG	NE-CZ-NH1	9.18	124.89	120.30
1	C	105	ARG	NE-CZ-NH2	-8.02	116.29	120.30
1	D	51	LEU	CA-CB-CG	-6.11	101.24	115.30
1	A	51	LEU	CA-CB-CG	-5.82	101.91	115.30
1	D	184	CYS	CB-CA-C	-5.38	99.63	110.40
1	B	208	MET	CG-SD-CE	5.33	108.73	100.20

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1769	0	1771	18	0
1	B	1765	0	1768	35	0
1	C	1765	0	1768	8	0
1	D	1743	0	1752	24	0
2	A	92	0	0	5	0
2	B	63	0	0	3	0
2	C	94	0	0	1	0
2	D	71	0	0	2	0
All	All	7362	0	7059	72	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (72) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:114:MET:CE	1:B:125:PRO:HB2	1.81	1.09
1:B:114:MET:HE1	1:B:125:PRO:HB2	1.02	0.99
1:A:191:GLN:HB2	2:A:311:HOH:O	1.62	0.97
1:B:114:MET:HE1	1:B:125:PRO:CB	1.96	0.92
1:B:213:LYS:O	1:B:213:LYS:HG3	1.78	0.83
1:D:211:LYS:HD3	2:D:290:HOH:O	1.79	0.82
1:B:201:ASN:HD22	1:B:202:GLU:H	1.27	0.80
1:A:104:SER:OG	1:A:106:GLU:HG2	1.83	0.78
1:B:49:THR:HG23	1:D:176:VAL:HG12	1.65	0.78
1:B:49:THR:CG2	1:D:176:VAL:H	2.03	0.71
1:B:94:LYS:HD3	1:B:95:TRP:H	1.55	0.70
1:D:191:GLN:HE21	1:D:193:ASP:HB2	1.55	0.69
1:B:213:LYS:O	1:B:213:LYS:CG	2.44	0.65
1:D:206:VAL:HB	1:D:215:LEU:HD23	1.80	0.63
1:A:13:THR:HG22	1:A:24:GLU:HG3	1.81	0.61
1:B:195:ILE:HG21	1:B:203:ILE:HD12	1.83	0.59
1:B:154:ALA:H	1:D:177:ASN:HD21	1.50	0.59
1:B:49:THR:HG23	1:D:176:VAL:CG1	2.33	0.58
1:A:195:ILE:HG12	1:A:203:ILE:CD1	2.34	0.57
1:A:30:GLU:HG3	2:A:290:HOH:O	2.04	0.57
1:B:194:GLU:O	1:B:194:GLU:HG2	2.05	0.56
1:A:195:ILE:HG12	1:A:203:ILE:HD13	1.86	0.56
1:C:159:ARG:HD2	1:C:178:TRP:O	2.06	0.56

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:139:LYS:HD2	2:B:281:HOH:O	2.06	0.55
1:B:49:THR:HG21	1:D:176:VAL:H	1.72	0.54
1:C:3:TYR:HA	2:C:302:HOH:O	2.08	0.54
1:A:206:VAL:HB	1:A:215:LEU:HD23	1.89	0.53
1:B:191:GLN:CD	1:B:191:GLN:H	2.10	0.53
1:B:206:VAL:HB	1:B:215:LEU:HD23	1.91	0.53
1:B:183:LYS:HG2	2:B:257:HOH:O	2.09	0.52
1:B:201:ASN:HD22	1:B:202:GLU:N	2.04	0.52
1:B:192:ASP:HA	1:B:195:ILE:HD12	1.90	0.52
1:A:4:HIS:HB3	2:A:312:HOH:O	2.10	0.51
1:B:31:ASN:N	2:B:271:HOH:O	2.31	0.50
1:A:124:LEU:HB3	1:A:125:PRO:HD2	1.94	0.50
1:C:30:GLU:HG2	1:C:30:GLU:O	2.11	0.49
1:A:191:GLN:HG3	1:A:192:ASP:H	1.77	0.48
1:B:215:LEU:HD21	1:D:89:TYR:CD1	2.48	0.48
1:C:26:TYR:CE1	1:C:138:LYS:HG2	2.48	0.48
1:C:26:TYR:CZ	1:C:138:LYS:HG2	2.48	0.47
1:C:94:LYS:HE2	1:C:94:LYS:HB3	1.68	0.47
1:D:13:THR:HA	1:D:23:PHE:O	2.16	0.46
1:B:85:ASP:O	1:D:208:MET:CE	2.63	0.46
1:B:85:ASP:O	1:D:208:MET:HE1	2.16	0.46
1:A:88:TYR:O	1:A:91:LYS:HE2	2.16	0.46
1:B:195:ILE:HD11	1:B:216:ARG:NH1	2.31	0.45
1:A:117:GLN:HG3	1:A:118:GLU:N	2.32	0.45
1:A:138:LYS:HE3	2:A:260:HOH:O	2.15	0.45
1:B:4:HIS:NE2	1:D:112:LYS:HD2	2.31	0.45
1:A:129:ARG:HB3	1:A:152:ARG:CZ	2.47	0.45
1:B:94:LYS:CD	1:B:95:TRP:H	2.25	0.45
1:B:195:ILE:CG2	1:B:203:ILE:HD12	2.48	0.44
1:B:94:LYS:HD3	1:B:95:TRP:N	2.29	0.44
1:B:85:ASP:CA	1:D:208:MET:CE	2.96	0.44
1:A:170:THR:CG2	1:A:171:PRO:HD2	2.48	0.44
1:C:192:ASP:OD1	1:C:192:ASP:N	2.50	0.43
1:D:128:CYS:HB2	1:D:146:TYR:O	2.18	0.43
1:D:195:ILE:HD11	1:D:216:ARG:NH1	2.34	0.43
1:A:191:GLN:NE2	2:A:311:HOH:O	2.48	0.42
1:D:120:ASP:HB3	1:D:126:LEU:HD21	2.02	0.41
1:A:104:SER:O	1:A:105:ARG:HB2	2.21	0.41
1:C:141:LYS:HD2	1:C:141:LYS:HA	1.84	0.41
1:B:153:ASN:HA	1:D:177:ASN:HD22	1.86	0.41
1:B:154:ALA:H	1:D:177:ASN:ND2	2.16	0.41

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:85:ASP:HB3	1:D:208:MET:HE1	2.03	0.41
1:A:91:LYS:HD3	1:A:91:LYS:HA	1.57	0.41
1:B:164:LEU:O	1:B:168:TYR:HB2	2.21	0.41
1:B:128:CYS:HB2	1:B:146:TYR:O	2.21	0.40
1:D:71:PHE:C	1:D:71:PHE:CD2	2.95	0.40
1:D:192:ASP:HA	1:D:195:ILE:HD12	2.04	0.40
1:D:20:ASP:HB2	2:D:284:HOH:O	2.21	0.40
1:D:159:ARG:HD2	1:D:178:TRP:O	2.22	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	217/220 (99%)	211 (97%)	6 (3%)	0	100	100
1	B	216/220 (98%)	204 (94%)	11 (5%)	1 (0%)	34	41
1	C	216/220 (98%)	208 (96%)	7 (3%)	1 (0%)	34	41
1	D	214/220 (97%)	205 (96%)	9 (4%)	0	100	100
All	All	863/880 (98%)	828 (96%)	33 (4%)	2 (0%)	52	64

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	31	ASN
1	C	40	PRO

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar

resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	199/200 (100%)	189 (95%)	10 (5%)	30	41
1	B	199/200 (100%)	187 (94%)	12 (6%)	24	31
1	C	199/200 (100%)	190 (96%)	9 (4%)	34	46
1	D	197/200 (98%)	190 (96%)	7 (4%)	42	57
All	All	794/800 (99%)	756 (95%)	38 (5%)	31	42

All (38) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	13	THR
1	A	32	SER
1	A	59	GLU
1	A	116	GLU
1	A	193	ASP
1	A	194	GLU
1	A	195	ILE
1	A	196	SER
1	A	202	GLU
1	A	215	LEU
1	B	27	LYS
1	B	49	THR
1	B	71	PHE
1	B	72	SER
1	B	94	LYS
1	B	117	GLN
1	B	138	LYS
1	B	170	THR
1	B	191	GLN
1	B	201	ASN
1	B	202	GLU
1	B	213	LYS
1	C	5	LEU
1	C	30	GLU
1	C	62	LEU
1	C	105	ARG
1	C	137	GLU
1	C	192	ASP

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	C	195	ILE
1	C	203	ILE
1	C	215	LEU
1	D	13	THR
1	D	36	LEU
1	D	44	THR
1	D	96	GLU
1	D	100	VAL
1	D	193	ASP
1	D	207	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	117	GLN
1	A	198	HIS
1	A	201	ASN
1	B	201	ASN
1	C	201	ASN
1	D	177	ASN
1	D	191	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

There are no ligands in this entry.

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	219/220 (99%)	-0.13	4 (1%) 71 78	22, 31, 65, 78	0
1	B	218/220 (99%)	-0.04	6 (2%) 56 66	26, 38, 73, 87	0
1	C	218/220 (99%)	-0.17	3 (1%) 78 83	22, 31, 60, 70	0
1	D	216/220 (98%)	-0.17	1 (0%) 91 94	25, 35, 65, 77	0
All	All	871/880 (98%)	-0.13	14 (1%) 74 80	22, 34, 64, 87	0

All (14) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	193	ASP	4.3
1	B	20	ASP	4.1
1	B	198	HIS	3.0
1	B	201	ASN	2.8
1	A	197	LYS	2.8
1	C	200	LYS	2.7
1	C	202	GLU	2.6
1	A	200	LYS	2.6
1	B	199	PHE	2.4
1	A	4	HIS	2.4
1	D	20	ASP	2.4
1	B	202	GLU	2.2
1	B	19	ILE	2.1
1	A	193	ASP	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

### 6.5 Other polymers [i](#)

There are no such residues in this entry.