



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 19, 2016 – 07:56 PM GMT

PDB ID : 4XCR  
Title : Monomeric Human Cu,Zn Superoxide dismutase, loops IV and VII deleted, apo form, mutant I35A  
Authors : Wang, H.; Logan, D.T.; Danielsson, J.; Mu, X.; Binolfi, A.; Theillet, F.; Bekei, B.; Lang, L.; Wennerstrom, H.; Selenko, P.; Oliveberg, M.  
Deposited on : 2014-12-18  
Resolution : 3.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : unknown  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026982  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20026982


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*



The reported resolution of this entry is 3.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	1408 (3.80-3.40)
Clashscore	102246	1010 (3.74-3.46)
Ramachandran outliers	100387	1007 (3.76-3.44)
Sidechain outliers	100360	1007 (3.76-3.44)
RSRZ outliers	91569	1003 (3.78-3.42)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	110	 91% 8% •
1	B	110	 90% 10%

## 2 Entry composition

There is only 1 type of molecule in this entry. The entry contains 3085 atoms, of which 1543 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Superoxide dismutase [Cu-Zn].

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	110	Total	C	H	N	O	0	0	0
			1542	477	771	140	154			
1	B	110	Total	C	H	N	O	0	0	0
			1543	477	772	140	154			

There are 104 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	6	ALA	CYS	conflict	UNP P00441
A	35	ALA	ILE	engineered mutation	UNP P00441
A	49	GLY	GLU	conflict	UNP P00441
A	50	ALA	PHE	conflict	UNP P00441
A	?	-	ASP	deletion	UNP P00441
A	?	-	ASN	deletion	UNP P00441
A	?	-	THR	deletion	UNP P00441
A	?	-	ALA	deletion	UNP P00441
A	?	-	GLY	deletion	UNP P00441
A	?	-	CYS	deletion	UNP P00441
A	?	-	THR	deletion	UNP P00441
A	?	-	SER	deletion	UNP P00441
A	?	-	ALA	deletion	UNP P00441
A	?	-	GLY	deletion	UNP P00441
A	?	-	PRO	deletion	UNP P00441
A	?	-	HIS	deletion	UNP P00441
A	?	-	PHE	deletion	UNP P00441
A	?	-	ASN	deletion	UNP P00441
A	?	-	PRO	deletion	UNP P00441
A	?	-	LEU	deletion	UNP P00441
A	?	-	SER	deletion	UNP P00441
A	?	-	ARG	deletion	UNP P00441
A	?	-	LYS	deletion	UNP P00441
A	?	-	HIS	deletion	UNP P00441
A	?	-	GLY	deletion	UNP P00441

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
A	?	-	GLY	deletion	UNP P00441
A	?	-	PRO	deletion	UNP P00441
A	?	-	LYS	deletion	UNP P00441
A	?	-	ASP	deletion	UNP P00441
A	?	-	GLU	deletion	UNP P00441
A	?	-	GLU	deletion	UNP P00441
A	?	-	ARG	deletion	UNP P00441
A	?	-	HIS	deletion	UNP P00441
A	?	-	VAL	deletion	UNP P00441
A	81	SER	CYS	conflict	UNP P00441
A	?	-	ASP	deletion	UNP P00441
A	?	-	ASP	deletion	UNP P00441
A	?	-	LEU	deletion	UNP P00441
A	?	-	GLY	deletion	UNP P00441
A	?	-	LYS	deletion	UNP P00441
A	?	-	GLY	deletion	UNP P00441
A	?	-	GLY	deletion	UNP P00441
A	?	-	ASN	deletion	UNP P00441
A	?	-	GLU	deletion	UNP P00441
A	?	-	GLU	deletion	UNP P00441
A	?	-	SER	deletion	UNP P00441
A	?	-	THR	deletion	UNP P00441
A	?	-	LYS	deletion	UNP P00441
A	94	GLY	THR	conflict	UNP P00441
A	95	ALA	GLY	conflict	UNP P00441
A	96	GLY	ASN	conflict	UNP P00441
A	103	SER	CYS	conflict	UNP P00441
B	6	ALA	CYS	conflict	UNP P00441
B	35	ALA	ILE	engineered mutation	UNP P00441
B	49	GLY	GLU	conflict	UNP P00441
B	50	ALA	PHE	conflict	UNP P00441
B	?	-	ASP	deletion	UNP P00441
B	?	-	ASN	deletion	UNP P00441
B	?	-	THR	deletion	UNP P00441
B	?	-	ALA	deletion	UNP P00441
B	?	-	GLY	deletion	UNP P00441
B	?	-	CYS	deletion	UNP P00441
B	?	-	THR	deletion	UNP P00441
B	?	-	SER	deletion	UNP P00441
B	?	-	ALA	deletion	UNP P00441
B	?	-	GLY	deletion	UNP P00441
B	?	-	PRO	deletion	UNP P00441

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
B	?	-	HIS	deletion	UNP P00441
B	?	-	PHE	deletion	UNP P00441
B	?	-	ASN	deletion	UNP P00441
B	?	-	PRO	deletion	UNP P00441
B	?	-	LEU	deletion	UNP P00441
B	?	-	SER	deletion	UNP P00441
B	?	-	ARG	deletion	UNP P00441
B	?	-	LYS	deletion	UNP P00441
B	?	-	HIS	deletion	UNP P00441
B	?	-	GLY	deletion	UNP P00441
B	?	-	GLY	deletion	UNP P00441
B	?	-	PRO	deletion	UNP P00441
B	?	-	LYS	deletion	UNP P00441
B	?	-	ASP	deletion	UNP P00441
B	?	-	GLU	deletion	UNP P00441
B	?	-	GLU	deletion	UNP P00441
B	?	-	ARG	deletion	UNP P00441
B	?	-	HIS	deletion	UNP P00441
B	?	-	VAL	deletion	UNP P00441
B	81	SER	CYS	conflict	UNP P00441
B	?	-	ASP	deletion	UNP P00441
B	?	-	ASP	deletion	UNP P00441
B	?	-	LEU	deletion	UNP P00441
B	?	-	GLY	deletion	UNP P00441
B	?	-	LYS	deletion	UNP P00441
B	?	-	GLY	deletion	UNP P00441
B	?	-	GLY	deletion	UNP P00441
B	?	-	ASN	deletion	UNP P00441
B	?	-	GLU	deletion	UNP P00441
B	?	-	GLU	deletion	UNP P00441
B	?	-	SER	deletion	UNP P00441
B	?	-	THR	deletion	UNP P00441
B	?	-	LYS	deletion	UNP P00441
B	94	GLY	THR	conflict	UNP P00441
B	95	ALA	GLY	conflict	UNP P00441
B	96	GLY	ASN	conflict	UNP P00441
B	103	SER	CYS	conflict	UNP P00441

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $\text{RSRZ} > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Superoxide dismutase [Cu-Zn]

Chain A:  91% 8% •



- Molecule 1: Superoxide dismutase [Cu-Zn]

Chain B:  90% 10%



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 65	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	70.82Å 70.82Å 70.01Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	46.13 – 3.60 46.13 – 3.60	Depositor EDS
% Data completeness (in resolution range)	93.0 (46.13-3.60) 92.8 (46.13-3.60)	Depositor EDS
$R_{merge}$	0.33	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.16 (at 3.57Å)	Xtriage
Refinement program	PHENIX	Depositor
R, $R_{free}$	0.185 , 0.239 0.182 , 0.236	Depositor DCC
$R_{free}$ test set	219 reflections (9.97%)	DCC
Wilson B-factor (Å <sup>2</sup> )	61.8	Xtriage
Anisotropy	0.001	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , -7.3	EDS
Estimated twinning fraction	0.791 for H, K, L 0.209 for K, H, -L 0.106 for h,-h-k,-l	Xtriage
Reported twinning fraction	0.791 for H, K, L 0.209 for K, H, -L	Depositor
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.44$ , $\langle L^2 \rangle = 0.27$	Xtriage
Outliers	0 of 2203 reflections	Xtriage
$F_o, F_c$ correlation	0.92	EDS
Total number of atoms	3085	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	58.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.54% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.24	0/781	0.44	0/1053
1	B	0.25	0/781	0.47	0/1053
All	All	0.25	0/1562	0.45	0/2106

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	771	771	771	5	0
1	B	771	772	771	4	0
All	All	1542	1543	1542	9	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (9) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:3:LYS:NZ	1:A:21:GLU:OE1	2.25	0.69
1:B:3:LYS:NZ	1:B:110:GLN:O	2.29	0.62

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:29:VAL:N	1:A:71:ASP:O	2.43	0.51
1:A:8:LEU:N	1:A:16:GLY:O	2.43	0.49
1:A:29:VAL:O	1:A:71:ASP:N	2.52	0.43
1:A:46:HIS:HB3	1:A:53:ASP:HA	2.02	0.42
1:B:13:PRO:O	1:B:15:GLN:N	2.52	0.42
1:B:43:HIS:ND1	1:B:90:HIS:O	2.53	0.41
1:B:22:GLN:HB2	1:B:76:LEU:HD12	2.03	0.41

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	108/110 (98%)	96 (89%)	11 (10%)	1 (1%)	21	67
1	B	108/110 (98%)	100 (93%)	7 (6%)	1 (1%)	21	67
All	All	216/220 (98%)	196 (91%)	18 (8%)	2 (1%)	21	67

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	14	VAL
1	A	52	GLY

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	78/78 (100%)	76 (97%)	2 (3%)	54	83
1	B	78/78 (100%)	76 (97%)	2 (3%)	54	83
All	All	156/156 (100%)	152 (97%)	4 (3%)	54	83

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	46	HIS
1	A	60	ASP
1	B	2	THR
1	B	69	ILE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	110/110 (100%)	-0.26	0 100 100	45, 58, 73, 84	1 (0%)
1	B	110/110 (100%)	-0.15	0 100 100	48, 64, 83, 90	1 (0%)
All	All	220/220 (100%)	-0.21	0 100 100	45, 61, 80, 90	2 (0%)

There are no RSRZ outliers to report.

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

### 6.4 Ligands ⓘ

There are no ligands in this entry.

### 6.5 Other polymers ⓘ

There are no such residues in this entry.