



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 06:32 AM GMT

PDB ID : 2XFP
Title : ISATIN-INHIBITED HUMAN MONOAMINE OXIDASE B IN COMPLEX WITH 2-(2-BENZOFURANYL)-2-IMIDAZOLINE
Authors : Bonivento, D.; Milczek, E.M.; Mcdonald, G.R.; Binda, C.; Holt, A.; Edmondson, D.E.; Mattevi, A.
Deposited on : 2010-05-26
Resolution : 1.66 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.66 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1226 (1.66-1.66)
Clashscore	102246	1323 (1.66-1.66)
Ramachandran outliers	100387	1295 (1.66-1.66)
Sidechain outliers	100360	1295 (1.66-1.66)
RSRZ outliers	91569	1227 (1.66-1.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	520	 2% 86% 9% • •
1	B	520	 3% 83% 11% • 5%

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 9049 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called AMINE OXIDASE [FLAVIN-CONTAINING] B.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	499	Total	C	N	O	S	0	6	0
			4003	2564	681	731	27			
1	B	494	Total	C	N	O	S	0	2	0
			3950	2527	676	721	26			

- Molecule 2 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula: $C_{27}H_{33}N_9O_{15}P_2$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
2	B	1	Total	C	N	O	P	0	0
			53	27	9	15	2		

- Molecule 3 is ISATIN (three-letter code: ISN) (formula: $C_8H_5NO_2$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			11	8	1	2		
3	B	1	Total	C	N	O	0	0
			11	8	1	2		

- Molecule 4 is 2-(2-BENZOFURANYL)-2-IMIDAZOLINE (three-letter code: XCG) (formula: $C_{11}H_8N_2O$).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			14	11	2	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	B	1	Total	C	N	O	0	0
			14	11	2	1		

- Molecule 5 is N-DODECYL-N,N-DIMETHYL-3-AMMONIO-1-PROPANESULFONATE (three-letter code: C15) (formula: $C_{17}H_{38}NO_3S$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	A	1	Total	C	N	O	S	0	0
			22	17	1	3	1		

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	435	Total	O	0	0
			435	435		
6	B	483	Total	O	0	0
			483	483		

4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	131.08Å 222.61Å 86.40Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	27.88 – 1.66 27.88 – 1.66	Depositor EDS
% Data completeness (in resolution range)	95.7 (27.88-1.66) 95.7 (27.88-1.66)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.15 (at 1.66Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.161 , 0.192 0.163 , 0.195	Depositor DCC
R_{free} test set	3657 reflections (2.65%)	DCC
Wilson B-factor (Å ²)	13.2	Xtriage
Anisotropy	0.049	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 50.3	EDS
Estimated twinning fraction	0.010 for 1/2*h-1/2*k,-3/2*h-1/2*k,-l 0.016 for 1/2*h+1/2*k,3/2*h-1/2*k,-l	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 141444 reflections	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	9049	wwPDB-VP
Average B, all atoms (Å ²)	15.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.20% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: XCG, C15, FAD, ISN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.38	14/4118 (0.3%)	1.15	18/5587 (0.3%)
1	B	1.42	13/4053 (0.3%)	1.19	15/5499 (0.3%)
All	All	1.40	27/8171 (0.3%)	1.17	33/11086 (0.3%)

All (27) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	413	VAL	CB-CG2	-7.64	1.36	1.52
1	B	305	PHE	CE1-CZ	6.29	1.49	1.37
1	A	358	GLU	CG-CD	6.22	1.61	1.51
1	B	106	VAL	CB-CG2	6.13	1.65	1.52
1	A	390	GLU	CG-CD	6.05	1.61	1.51
1	A	484	ARG	CG-CD	5.96	1.66	1.51
1	A	247	VAL	CB-CG1	5.93	1.65	1.52
1	B	321	GLU	CG-CD	5.91	1.60	1.51
1	A	321	GLU	CG-CD	5.86	1.60	1.51
1	A	435	TYR	CG-CD2	5.84	1.46	1.39
1	A	300	TYR	CD1-CE1	5.80	1.48	1.39
1	B	410	TYR	CD1-CE1	5.71	1.48	1.39
1	A	60	TYR	CD1-CE1	5.70	1.48	1.39
1	A	337	TYR	CD2-CE2	5.58	1.47	1.39
1	B	128	GLU	CG-CD	5.55	1.60	1.51
1	A	453	ALA	CA-CB	5.54	1.64	1.52
1	B	415	ARG	CB-CG	-5.53	1.37	1.52
1	B	413	VAL	CA-CB	5.52	1.66	1.54
1	A	388	TRP	CB-CG	-5.37	1.40	1.50
1	B	150	GLU	CD-OE1	5.27	1.31	1.25
1	A	294	VAL	CB-CG1	5.25	1.63	1.52
1	A	280	MET	CB-CG	5.24	1.68	1.51
1	B	112	TYR	CD2-CE2	5.19	1.47	1.39
1	B	256	GLU	CB-CG	-5.16	1.42	1.52

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	320	GLU	CG-CD	5.13	1.59	1.51
1	A	112	TYR	CD2-CE2	5.13	1.47	1.39
1	B	106	VAL	CB-CG1	-5.12	1.42	1.52

All (33) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	227	ASP	CB-CG-OD1	10.04	127.34	118.30
1	B	222	MET	CG-SD-CE	9.67	115.67	100.20
1	B	47	ARG	NE-CZ-NH2	-9.58	115.51	120.30
1	A	127	ARG	NE-CZ-NH2	-8.78	115.91	120.30
1	B	227	ASP	CB-CG-OD2	-7.86	111.23	118.30
1	B	36	ARG	NE-CZ-NH2	-7.61	116.49	120.30
1	A	47	ARG	NE-CZ-NH1	-7.52	116.54	120.30
1	B	329	ASP	CB-CG-OD2	-7.23	111.79	118.30
1	A	124	ASP	CB-CG-OD2	-7.05	111.95	118.30
1	B	305	PHE	CB-CG-CD2	-6.91	115.96	120.80
1	B	167	LEU	CB-CG-CD2	-6.86	99.34	111.00
1	A	448	ARG	NE-CZ-NH1	-6.64	116.98	120.30
1	A	127	ARG	CG-CD-NE	-6.38	98.41	111.80
1	A	308	LYS	CD-CE-NZ	6.25	126.06	111.70
1	A	233	ARG	NE-CZ-NH2	-5.94	117.33	120.30
1	B	280	MET	CG-SD-CE	-5.92	90.72	100.20
1	A	36	ARG	NE-CZ-NH1	5.89	123.25	120.30
1	A	288	ARG	NE-CZ-NH2	-5.84	117.38	120.30
1	A	132	ASP	CB-CG-OD2	-5.84	113.05	118.30
1	B	124	ASP	CB-CG-OD2	-5.62	113.24	118.30
1	B	208	ARG	NE-CZ-NH2	-5.56	117.52	120.30
1	B	494	ARG	NE-CZ-NH1	5.54	123.07	120.30
1	B	460	ASP	CB-CG-OD2	5.53	123.28	118.30
1	B	239	ASP	CB-CG-OD2	-5.43	113.41	118.30
1	A	254[A]	MET	CG-SD-CE	5.38	108.81	100.20
1	A	254[B]	MET	CG-SD-CE	5.38	108.81	100.20
1	A	120	ARG	NE-CZ-NH2	5.36	122.98	120.30
1	A	132	ASP	CB-CG-OD1	5.34	123.11	118.30
1	A	233	ARG	NE-CZ-NH1	5.29	122.94	120.30
1	B	254	MET	CG-SD-CE	-5.27	91.77	100.20
1	A	309	LYS	CD-CE-NZ	-5.25	99.64	111.70
1	A	420	ARG	NE-CZ-NH2	-5.11	117.75	120.30
1	A	227	ASP	CB-CG-OD1	5.07	122.87	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4003	0	4025	22	2
1	B	3950	0	3955	41	0
2	A	53	0	29	2	0
2	B	53	0	29	0	0
3	A	11	0	5	0	0
3	B	11	0	5	1	0
4	A	14	0	8	0	0
4	B	14	0	8	1	0
5	A	22	0	38	0	0
6	A	435	0	0	3	0
6	B	483	0	0	19	2
All	All	9049	0	8102	60	3

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (60) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:76:GLY:HA3	6:B:2096:HOH:O	1.74	0.85
1:B:227:ASP:HB3	6:B:2257:HOH:O	1.79	0.83
1:A:163:GLN:OE1	6:A:2197:HOH:O	1.98	0.81
1:A:244:ASN:ND2	1:A:256:GLU:OE1	2.14	0.77
1:B:414:LEU:HD22	6:B:2408:HOH:O	1.87	0.74
1:A:117:ASN:HD22	1:A:120:ARG:HH21	1.35	0.73
1:B:117:ASN:HD22	1:B:120:ARG:HH21	1.38	0.71
1:B:232:GLU:OE1	6:B:2263:HOH:O	2.09	0.70
1:A:251:ASN:O	1:A:252:HIS:HB2	1.94	0.67
1:B:227:ASP:N	6:B:2256:HOH:O	2.29	0.64
1:A:280:MET:CE	6:B:2386:HOH:O	2.47	0.63
1:B:410:TYR:O	1:B:413:VAL:HG13	1.99	0.63
1:B:17:MET:CE	1:B:39:VAL:HG11	2.29	0.62

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:219:GLU:OE2	6:B:2249:HOH:O	2.15	0.61
1:B:220:ARG:NH2	6:B:2250:HOH:O	2.24	0.60
1:B:451:LEU:HD23	1:B:454:MET:CE	2.31	0.60
1:B:48:ASN:HA	6:B:2061:HOH:O	2.01	0.59
1:B:285[B]:MET:SD	1:B:413:VAL:HG22	2.43	0.59
1:B:414:LEU:CD2	6:B:2408:HOH:O	2.47	0.59
1:A:305:PHE:HA	1:A:308:LYS:HE3	1.85	0.58
1:B:451:LEU:HD23	1:B:454:MET:HE2	1.87	0.56
1:A:376[A]:GLU:HG3	6:A:2337:HOH:O	2.04	0.56
1:A:321:GLU:HG2	6:A:2291:HOH:O	2.06	0.56
1:A:280:MET:HE1	1:B:353:ALA:HB1	1.88	0.55
1:B:100:ARG:NH1	6:B:2127:HOH:O	2.40	0.55
1:B:142:GLU:OE1	6:B:2182:HOH:O	2.18	0.54
1:B:220:ARG:HG3	6:B:2252:HOH:O	2.08	0.53
1:B:223:ASP:OD1	6:B:2254:HOH:O	2.19	0.53
1:B:88:LEU:N	1:B:88:LEU:HD12	2.27	0.50
1:A:3:ASN:HD22	1:A:3:ASN:N	2.09	0.50
1:A:410:TYR:OH	1:B:350:ARG:HD2	2.14	0.48
1:B:136:LYS:HE3	6:B:2169:HOH:O	2.13	0.48
1:A:500:THR:O	1:A:500:THR:HG22	2.15	0.47
1:B:454:MET:HE1	1:B:456:LYS:HD2	1.96	0.47
1:B:454:MET:HB2	1:B:454:MET:HE3	1.71	0.47
1:A:500:THR:O	1:A:501:ILE:HG13	2.15	0.47
1:A:499:THR:C	1:A:501:ILE:H	2.19	0.46
1:B:456:LYS:HD3	6:B:2022:HOH:O	2.16	0.46
1:B:309:LYS:NZ	1:B:373:GLY:O	2.49	0.46
1:A:58:GLY:HA2	2:A:600:FAD:C4X	2.46	0.45
1:A:117:ASN:HD22	1:A:120:ARG:NH2	2.09	0.45
1:A:280:MET:HE3	1:A:280:MET:HB3	1.32	0.45
1:B:28:LEU:HD11	1:B:454:MET:HE1	1.99	0.44
1:B:76:GLY:CA	6:B:2096:HOH:O	2.49	0.44
1:B:148:MET:O	1:B:152:LEU:HG	2.18	0.43
1:B:199:ILE:HD12	3:B:601:ISN:H3	1.99	0.43
1:B:332:LYS:HB3	1:B:333:PRO:HD2	1.99	0.43
1:B:332:LYS:HG3	1:B:337:TYR:CD1	2.54	0.43
1:B:102:PRO:O	4:B:602:XCG:HAE	2.18	0.43
1:A:270[B]:MET:SD	1:B:270[B]:MET:SD	3.17	0.43
1:B:28:LEU:HD11	1:B:454:MET:CE	2.49	0.43
1:B:117:ASN:HD22	1:B:120:ARG:NH2	2.11	0.43
1:B:228:ARG:NH1	6:B:2260:HOH:O	2.41	0.43
1:A:81:LYS:HE3	1:A:208:ARG:NH2	2.34	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:280:MET:HE1	6:B:2386:HOH:O	2.13	0.41
1:B:454:MET:CE	1:B:456:LYS:HD2	2.49	0.41
1:B:451:LEU:HA	1:B:454:MET:CE	2.51	0.41
1:A:58:GLY:HA2	2:A:600:FAD:C5X	2.50	0.41
1:A:486:LEU:HD23	1:A:486:LEU:HA	1.87	0.41
1:B:17:MET:CE	1:B:39:VAL:CG1	2.98	0.40

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
6:B:2254:HOH:O	6:B:2254:HOH:O[4_565]	1.68	0.52
1:A:475:GLN:NE2	1:A:475:GLN:NE2[3_656]	1.79	0.41
1:A:95:LYS:CE	6:B:2182:HOH:O[6_565]	2.18	0.02

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	503/520 (97%)	491 (98%)	11 (2%)	1 (0%)	52	30
1	B	494/520 (95%)	481 (97%)	13 (3%)	0	100	100
All	All	997/1040 (96%)	972 (98%)	24 (2%)	1 (0%)	56	33

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	500	THR

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	434/444 (98%)	426 (98%)	8 (2%)	66	43
1	B	426/444 (96%)	421 (99%)	5 (1%)	78	60
All	All	860/888 (97%)	847 (98%)	13 (2%)	72	51

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	3	ASN
1	A	52	LYS
1	A	93	LYS
1	A	167	LEU
1	A	280	MET
1	A	350	ARG
1	A	495	LEU
1	A	498	LEU
1	B	92	VAL
1	B	350	ARG
1	B	397	CYS
1	B	495	LEU
1	B	496	ILE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	117	ASN
1	A	163	GLN
1	A	452	HIS
1	A	475	GLN
1	A	485	HIS
1	B	116	ASN
1	B	117	ASN
1	B	452	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

7 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
5	C15	A	1503	-	20,21,21	0.73	0	23,26,26	1.33	3 (13%)
2	FAD	A	600	1	48,58,58	1.95	7 (14%)	54,89,89	2.30	12 (22%)
3	ISN	A	601	-	12,12,12	2.73	2 (16%)	17,17,17	4.17	9 (52%)
4	XCG	A	602	-	12,16,16	2.42	4 (33%)	10,22,22	2.42	5 (50%)
2	FAD	B	600	1	48,58,58	1.91	9 (18%)	54,89,89	1.77	10 (18%)
3	ISN	B	601	-	12,12,12	2.83	5 (41%)	17,17,17	4.79	11 (64%)
4	XCG	B	602	-	12,16,16	2.66	6 (50%)	10,22,22	2.51	6 (60%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	C15	A	1503	-	-	0/21/21/21	0/0/0/0
2	FAD	A	600	1	-	0/30/50/50	0/6/6/6

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	ISN	A	601	-	-	0/0/12/12	0/2/2/2
4	XCG	A	602	-	-	0/0/4/4	0/2/3/3
2	FAD	B	600	1	-	0/30/50/50	0/6/6/6
3	ISN	B	601	-	-	0/0/12/12	0/2/2/2
4	XCG	B	602	-	-	0/0/4/4	0/2/3/3

All (33) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	601	ISN	C10-C7	-7.85	1.41	1.56
3	B	601	ISN	C10-C7	-6.71	1.44	1.56
4	B	602	XCG	CAL-CAK	-5.11	1.39	1.49
4	A	602	XCG	CAL-CAK	-4.93	1.40	1.49
3	B	601	ISN	C1-C7	-4.86	1.37	1.48
3	A	601	ISN	C1-C7	-4.59	1.38	1.48
3	B	601	ISN	C10-N1	-2.93	1.33	1.36
2	B	600	FAD	C8A-N7A	-2.67	1.29	1.34
3	B	601	ISN	C5-C6	-2.19	1.35	1.39
4	B	602	XCG	CAK-NAI	-2.14	1.32	1.35
2	A	600	FAD	C8M-C8	2.02	1.55	1.51
2	A	600	FAD	C7M-C7	2.13	1.55	1.51
3	B	601	ISN	C4-C5	2.22	1.43	1.38
4	A	602	XCG	CAB-CAA	2.25	1.43	1.38
2	A	600	FAD	C1'-N10	2.32	1.50	1.48
2	B	600	FAD	O4B-C1B	2.33	1.44	1.41
2	B	600	FAD	C5B-C4B	2.36	1.59	1.51
4	A	602	XCG	CAD-CAN	2.38	1.43	1.38
2	B	600	FAD	C4A-N3A	2.42	1.39	1.35
2	B	600	FAD	O3B-C3B	2.58	1.49	1.43
4	B	602	XCG	CAG-NAI	2.71	1.48	1.37
2	A	600	FAD	C2B-C3B	2.73	1.60	1.53
2	B	600	FAD	C4-N3	2.81	1.38	1.33
4	B	602	XCG	CAB-CAD	3.03	1.43	1.36
2	B	600	FAD	C10-N1	3.11	1.40	1.35
4	B	602	XCG	CAF-NAH	3.27	1.51	1.37
4	B	602	XCG	CAG-CAF	4.11	1.52	1.36
2	A	600	FAD	C4-N3	4.68	1.41	1.33
4	A	602	XCG	CAG-CAF	4.74	1.54	1.36
2	B	600	FAD	C4X-N5	4.79	1.40	1.33
2	A	600	FAD	C10-N1	6.35	1.46	1.35
2	A	600	FAD	C4X-N5	6.61	1.43	1.33
2	B	600	FAD	C1'-N10	7.30	1.56	1.48

All (56) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	601	ISN	C6-N1-C10	-10.72	104.78	111.35
3	B	601	ISN	C6-N1-C10	-8.87	105.91	111.35
2	A	600	FAD	N3A-C2A-N1A	-8.17	122.64	128.89
3	B	601	ISN	C6-C1-C7	-7.27	102.26	107.29
2	A	600	FAD	C4X-C10-N10	-5.29	117.40	120.52
3	A	601	ISN	C6-C1-C7	-5.12	103.75	107.29
2	B	600	FAD	N3A-C2A-N1A	-4.45	125.49	128.89
5	A	1503	C15	O2S-S1-C1	-4.07	103.43	106.91
2	A	600	FAD	C1B-N9A-C4A	-3.71	121.35	126.94
2	A	600	FAD	C2B-C1B-N9A	-3.68	108.67	114.29
2	B	600	FAD	C1B-N9A-C4A	-3.61	121.50	126.94
4	A	602	XCG	CAA-CAC-CAM	-3.41	114.97	120.44
2	A	600	FAD	C4B-O4B-C1B	-3.31	106.08	109.72
4	B	602	XCG	CAA-CAC-CAM	-3.20	115.31	120.44
3	B	601	ISN	O11-C10-N1	-3.10	121.60	126.38
3	B	601	ISN	C5-C6-C1	-3.02	118.93	122.19
3	A	601	ISN	C3-C2-C1	-2.96	114.15	119.87
2	B	600	FAD	C4X-C10-N10	-2.88	118.82	120.52
2	A	600	FAD	O3'-C3'-C4'	-2.87	101.50	108.75
4	B	602	XCG	CAG-CAF-NAH	-2.55	102.83	109.30
4	A	602	XCG	CAG-CAF-NAH	-2.50	102.96	109.30
3	B	601	ISN	C4-C3-C2	-2.40	116.68	120.19
4	B	602	XCG	CAF-CAG-NAI	-2.36	103.31	109.30
2	B	600	FAD	C1'-C2'-C3'	-2.34	103.12	109.82
2	A	600	FAD	O4B-C4B-C5B	-2.34	100.96	109.32
2	A	600	FAD	C4X-C4-N3	-2.28	120.47	123.59
5	A	1503	C15	C2N-N1-C3	-2.15	103.97	109.43
2	B	600	FAD	O2'-C2'-C1'	-2.14	104.69	109.94
2	B	600	FAD	C4X-C4-N3	-2.07	120.76	123.59
2	B	600	FAD	O3'-C3'-C4'	-2.07	103.55	108.75
3	A	601	ISN	C5-C6-C1	-2.04	119.98	122.19
5	A	1503	C15	O1S-S1-C1	2.02	108.63	106.91
4	A	602	XCG	CAF-NAH-CAK	2.24	110.23	104.33
4	B	602	XCG	CAG-NAI-CAK	2.37	110.57	104.33
2	A	600	FAD	C2A-N1A-C6A	2.85	123.86	118.77
3	B	601	ISN	C7-C10-N1	2.96	108.55	106.07
4	A	602	XCG	CAB-CAA-CAC	3.09	124.95	120.45
2	A	600	FAD	C4-C4X-N5	3.37	122.81	118.72
3	B	601	ISN	O11-C10-C7	3.54	129.84	126.00
4	B	602	XCG	CAB-CAA-CAC	3.75	125.91	120.45
3	A	601	ISN	C2-C1-C6	3.76	123.26	118.86
3	A	601	ISN	C1-C6-N1	3.81	112.82	108.28

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	602	XCG	CAE-CAM-CAN	3.82	109.60	106.27
2	B	600	FAD	C4-C4X-N5	3.85	123.39	118.72
3	A	601	ISN	O8-C7-C10	3.86	127.76	123.95
3	B	601	ISN	C3-C4-C5	3.98	126.01	120.19
3	B	601	ISN	C1-C6-N1	4.11	113.17	108.28
4	A	602	XCG	CAE-CAM-CAN	4.35	110.07	106.27
2	B	600	FAD	C4-N3-C2	4.91	119.49	115.25
2	A	600	FAD	C4-N3-C2	4.94	119.52	115.25
2	B	600	FAD	C5X-C9A-N10	5.20	121.57	117.62
3	A	601	ISN	C7-C10-N1	5.20	110.42	106.07
3	B	601	ISN	C2-C1-C6	5.65	125.47	118.86
2	A	600	FAD	C1'-N10-C9A	6.41	126.06	118.86
3	A	601	ISN	C1-C7-C10	7.37	108.22	105.07
3	B	601	ISN	C1-C7-C10	11.63	110.05	105.07

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	600	FAD	2	0
3	B	601	ISN	1	0
4	B	602	XCG	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	499/520 (95%)	-0.17	11 (2%) 65 67	7, 12, 27, 56	0
1	B	494/520 (95%)	-0.17	15 (3%) 54 55	6, 11, 28, 61	0
All	All	993/1040 (95%)	-0.17	26 (2%) 59 60	6, 12, 28, 61	0

All (26) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	495	LEU	7.0
1	B	107	TRP	6.0
1	B	496	ILE	5.2
1	A	498	LEU	4.8
1	B	493	LEU	4.4
1	A	501	ILE	4.1
1	A	107	TRP	4.1
1	B	494	ARG	4.0
1	A	481	PHE	3.9
1	A	500	THR	3.4
1	B	243	GLU	3.0
1	B	3	ASN	2.8
1	B	252	HIS	2.7
1	A	242	ARG	2.6
1	A	354	ARG	2.5
1	B	242	ARG	2.4
1	A	499	THR	2.3
1	B	354	ARG	2.3
1	B	4	LYS	2.3
1	A	494	ARG	2.3
1	B	481	PHE	2.3
1	B	241	THR	2.2
1	A	3	ASN	2.2
1	B	93	LYS	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	252	HIS	2.1
1	B	482	LEU	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
5	C15	A	1503	22/22	0.85	0.18	1.50	17,38,55,58	0
3	ISN	A	601	11/11	0.94	0.12	1.15	13,19,25,26	0
3	ISN	B	601	11/11	0.92	0.11	0.81	12,16,23,26	0
2	FAD	A	600	53/53	0.98	0.08	-0.16	5,8,11,12	0
4	XCG	B	602	14/14	0.97	0.07	-0.50	9,11,14,16	0
2	FAD	B	600	53/53	0.98	0.07	-0.82	4,7,10,11	0
4	XCG	A	602	14/14	0.98	0.06	-1.82	9,11,12,13	0

6.5 Other polymers [i](#)

There are no such residues in this entry.