



# wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 10:21 PM GMT

PDB ID : 4XGI  
Title : Crystal structure of Glutamate dehydrogenase from Burkholderia thailandensis  
Authors : Seattle Structural Genomics Center for Infectious Disease (SSGCID)  
Deposited on : 2014-12-30  
Resolution : 2.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	6249 (2.00-2.00)
Clashscore	102246	7340 (2.00-2.00)
Ramachandran outliers	100387	7248 (2.00-2.00)
Sidechain outliers	100360	7247 (2.00-2.00)
RSRZ outliers	91569	6262 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	436	<div> <div>90%</div> <div>5% 5%</div> </div>
1	B	436	<div> <div>87%</div> <div>7% 5%</div> </div>
1	C	436	<div> <div>89%</div> <div>6% 5%</div> </div>
1	D	436	<div> <div>88%</div> <div>7% 5%</div> </div>
1	E	436	<div> <div>89%</div> <div>6% 5%</div> </div>

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
1	F	436	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	AKG	A	501	-	-	-	X
3	AKG	C	501	-	-	-	X
3	AKG	D	501	-	-	-	X
3	AKG	E	501	-	-	-	X
4	EDO	A	502	-	-	-	X
4	EDO	D	502	-	-	-	X
4	EDO	E	502	-	-	-	X
4	EDO	F	502	-	-	-	X

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 20777 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Glutamate dehydrogenase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	416	Total	C	N	O	S	0	5	0
			3138	1977	559	590	12			
1	B	415	Total	C	N	O	S	0	4	0
			3135	1975	554	594	12			
1	C	416	Total	C	N	O	S	0	7	0
			3152	1981	564	595	12			
1	D	415	Total	C	N	O	S	0	3	0
			3127	1969	555	591	12			
1	E	415	Total	C	N	O	S	0	5	0
			3127	1967	555	593	12			
1	F	415	Total	C	N	O	S	0	4	0
			3107	1954	554	587	12			

There are 12 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	GLY	-	expression tag	UNP Q2SZ78
A	0	SER	-	expression tag	UNP Q2SZ78
B	-1	GLY	-	expression tag	UNP Q2SZ78
B	0	SER	-	expression tag	UNP Q2SZ78
C	-1	GLY	-	expression tag	UNP Q2SZ78
C	0	SER	-	expression tag	UNP Q2SZ78
D	-1	GLY	-	expression tag	UNP Q2SZ78
D	0	SER	-	expression tag	UNP Q2SZ78
E	-1	GLY	-	expression tag	UNP Q2SZ78
E	0	SER	-	expression tag	UNP Q2SZ78
F	-1	GLY	-	expression tag	UNP Q2SZ78
F	0	SER	-	expression tag	UNP Q2SZ78

- Molecule 2 is NICOTINAMIDE-ADENINE-DINUCLEOTIDE (three-letter code: NAD) (formula: C<sub>21</sub>H<sub>27</sub>N<sub>7</sub>O<sub>14</sub>P<sub>2</sub>).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			44	21	7	14	2		
2	B	1	Total	C	N	O	P	0	0
			44	21	7	14	2		
2	C	1	Total	C	N	O	P	0	0
			44	21	7	14	2		
2	D	1	Total	C	N	O	P	0	0
			44	21	7	14	2		
2	E	1	Total	C	N	O	P	0	0
			44	21	7	14	2		
2	F	1	Total	C	N	O	P	0	0
			44	21	7	14	2		

- Molecule 3 is 2-OXOGLUTARIC ACID (three-letter code: AKG) (formula: C<sub>5</sub>H<sub>6</sub>O<sub>5</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			10	5	5		
3	B	1	Total	C	O	0	0
			10	5	5		
3	C	1	Total	C	O	0	0
			10	5	5		
3	D	1	Total	C	O	0	0
			10	5	5		
3	E	1	Total	C	O	0	0
			10	5	5		
3	F	1	Total	C	O	0	0
			10	5	5		

- Molecule 4 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C<sub>2</sub>H<sub>6</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			4	2	2		
4	B	1	Total	C	O	0	0
			4	2	2		
4	C	1	Total	C	O	0	0
			4	2	2		
4	D	1	Total	C	O	0	0
			4	2	2		
4	E	1	Total	C	O	0	0
			4	2	2		
4	F	1	Total	C	O	0	0
			4	2	2		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	273	Total	O	0	0
			273	273		
5	B	291	Total	O	0	1
			292	292		
5	C	259	Total	O	0	0
			259	259		
5	D	321	Total	O	0	0
			321	321		
5	E	269	Total	O	0	1
			270	270		
5	F	228	Total	O	0	0
			228	228		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Glutamate dehydrogenase




- Molecule 1: Glutamate dehydrogenase




- Molecule 1: Glutamate dehydrogenase





- Molecule 1: Glutamate dehydrogenase









● Molecule 1: Glutamate dehydrogenase



● Molecule 1: Glutamate dehydrogenase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	95.49Å 93.52Å 158.37Å 90.00° 106.23° 90.00°	Depositor
Resolution (Å)	45.84 – 2.00 45.84 – 2.00	Depositor EDS
% Data completeness (in resolution range)	96.8 (45.84-2.00) 96.8 (45.84-2.00)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.07 (at 2.00Å)	Xtriage
Refinement program	PHENIX	Depositor
R, $R_{free}$	0.176 , 0.222 0.182 , 0.227	Depositor DCC
$R_{free}$ test set	1996 reflections (1.16%)	DCC
Wilson B-factor (Å <sup>2</sup> )	22.3	Xtriage
Anisotropy	0.202	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.31 , 48.7	EDS
Estimated twinning fraction	0.026 for h,-k,-h-l	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.29$	Xtriage
Outliers	1 of 174735 reflections (0.001%)	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	20777	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	30.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.03% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: AKG, EDO, NAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.37	0/3211	0.51	0/4362
1	B	0.39	0/3206	0.53	0/4354
1	C	0.37	0/3230	0.51	0/4384
1	D	0.38	0/3195	0.51	0/4340
1	E	0.37	0/3200	0.52	0/4347
1	F	0.35	0/3177	0.51	0/4318
All	All	0.37	0/19219	0.51	0/26105

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3138	0	3134	14	0
1	B	3135	0	3116	23	0
1	C	3152	0	3148	16	0
1	D	3127	0	3110	21	0
1	E	3127	0	3100	15	0
1	F	3107	0	3075	13	0
2	A	44	0	26	1	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	44	0	26	2	0
2	C	44	0	26	0	0
2	D	44	0	26	0	0
2	E	44	0	26	2	0
2	F	44	0	26	0	0
3	A	10	0	4	1	0
3	B	10	0	4	0	0
3	C	10	0	4	0	0
3	D	10	0	4	2	0
3	E	10	0	4	0	0
3	F	10	0	4	0	0
4	A	4	0	6	0	0
4	B	4	0	6	0	0
4	C	4	0	6	0	0
4	D	4	0	6	0	0
4	E	4	0	6	0	0
4	F	4	0	6	0	0
5	A	273	0	0	1	0
5	B	292	0	0	2	0
5	C	259	0	0	1	0
5	D	321	0	0	4	0
5	E	270	0	0	2	0
5	F	228	0	0	1	0
All	All	20777	0	18899	96	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 96 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:64[A]:GLU:HG2	1:C:432:LEU:HD12	1.74	0.69
1:A:365:THR:HG21	1:A:392:VAL:HG21	1.77	0.66
1:A:321:ASN:HA	1:A:324:LYS:HD3	1.79	0.64
1:B:58:ILE:HB	1:C:56:ILE:HB	1.80	0.63
1:D:83[A]:SER:OG	5:D:810:HOH:O	2.16	0.62

There are no symmetry-related clashes.

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	417/436 (96%)	404 (97%)	13 (3%)	0	100	100
1	B	415/436 (95%)	402 (97%)	13 (3%)	0	100	100
1	C	419/436 (96%)	408 (97%)	11 (3%)	0	100	100
1	D	414/436 (95%)	403 (97%)	11 (3%)	0	100	100
1	E	416/436 (95%)	408 (98%)	8 (2%)	0	100	100
1	F	415/436 (95%)	402 (97%)	13 (3%)	0	100	100
All	All	2496/2616 (95%)	2427 (97%)	69 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	323/346 (93%)	323 (100%)	0	100	100
1	B	323/346 (93%)	319 (99%)	4 (1%)	78	81
1	C	325/346 (94%)	323 (99%)	2 (1%)	90	93
1	D	322/346 (93%)	320 (99%)	2 (1%)	90	93
1	E	321/346 (93%)	320 (100%)	1 (0%)	94	96
1	F	317/346 (92%)	316 (100%)	1 (0%)	94	96
All	All	1931/2076 (93%)	1921 (100%)	10 (0%)	92	94

5 of 10 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	228	GLU
1	C	367	SER
1	D	293	GLU
1	B	428	GLU
1	D	196	ILE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 11 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	182	ASN
1	D	183	GLN
1	E	315	ASN
1	B	407	HIS
1	E	31	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

18 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	NAD	A	500	-	38,48,48	1.53	8 (21%)	47,73,73	1.94	15 (31%)
3	AKG	A	501	-	3,9,9	1.30	0	4,11,11	1.00	0
4	EDO	A	502	-	3,3,3	0.54	0	2,2,2	0.17	0
2	NAD	B	500	-	38,48,48	1.38	5 (13%)	47,73,73	1.91	13 (27%)
3	AKG	B	501	-	3,9,9	1.36	0	4,11,11	1.45	1 (25%)
4	EDO	B	502	-	3,3,3	0.52	0	2,2,2	0.33	0
2	NAD	C	500	-	38,48,48	1.41	6 (15%)	47,73,73	1.96	14 (29%)
3	AKG	C	501	-	3,9,9	1.35	0	4,11,11	1.39	1 (25%)
4	EDO	C	502	-	3,3,3	0.54	0	2,2,2	0.15	0
2	NAD	D	500	-	38,48,48	1.60	10 (26%)	47,73,73	1.91	11 (23%)
3	AKG	D	501	-	3,9,9	1.42	1 (33%)	4,11,11	1.02	0
4	EDO	D	502	-	3,3,3	0.63	0	2,2,2	0.18	0
2	NAD	E	500	-	38,48,48	1.44	4 (10%)	47,73,73	1.75	12 (25%)
3	AKG	E	501	-	3,9,9	1.48	1 (33%)	4,11,11	1.17	1 (25%)
4	EDO	E	502	-	3,3,3	0.55	0	2,2,2	0.28	0
2	NAD	F	500	-	38,48,48	1.37	4 (10%)	47,73,73	1.95	13 (27%)
3	AKG	F	501	-	3,9,9	1.40	0	4,11,11	0.79	0
4	EDO	F	502	-	3,3,3	0.55	0	2,2,2	0.33	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAD	A	500	-	-	0/22/62/62	0/5/5/5
3	AKG	A	501	-	-	0/3/9/9	0/0/0/0
4	EDO	A	502	-	-	0/1/1/1	0/0/0/0
2	NAD	B	500	-	-	0/22/62/62	0/5/5/5
3	AKG	B	501	-	-	0/3/9/9	0/0/0/0
4	EDO	B	502	-	-	0/1/1/1	0/0/0/0
2	NAD	C	500	-	-	0/22/62/62	0/5/5/5
3	AKG	C	501	-	-	0/3/9/9	0/0/0/0
4	EDO	C	502	-	-	0/1/1/1	0/0/0/0
2	NAD	D	500	-	-	0/22/62/62	0/5/5/5
3	AKG	D	501	-	-	0/3/9/9	0/0/0/0
4	EDO	D	502	-	-	0/1/1/1	0/0/0/0
2	NAD	E	500	-	-	0/22/62/62	0/5/5/5
3	AKG	E	501	-	-	0/3/9/9	0/0/0/0
4	EDO	E	502	-	-	0/1/1/1	0/0/0/0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAD	F	500	-	-	0/22/62/62	0/5/5/5
3	AKG	F	501	-	-	0/3/9/9	0/0/0/0
4	EDO	F	502	-	-	0/1/1/1	0/0/0/0

The worst 5 of 39 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	500	NAD	O5D-C5D	-2.26	1.35	1.44
3	E	501	AKG	O5-C2	-2.14	1.18	1.22
3	D	501	AKG	O5-C2	-2.05	1.18	1.22
2	B	500	NAD	O5D-C5D	-2.05	1.36	1.44
2	D	500	NAD	O5D-C5D	-2.03	1.36	1.44

The worst 5 of 81 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	500	NAD	PN-O3-PA	-5.70	116.72	132.73
2	F	500	NAD	PN-O3-PA	-5.60	117.01	132.73
2	D	500	NAD	PN-O3-PA	-5.43	117.47	132.73
2	B	500	NAD	PN-O3-PA	-5.08	118.47	132.73
2	E	500	NAD	PN-O3-PA	-4.80	119.25	132.73

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5 monomers are involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	500	NAD	1	0
3	A	501	AKG	1	0
2	B	500	NAD	2	0
3	D	501	AKG	2	0
2	E	500	NAD	2	0

## 5.7 Other polymers

There are no such residues in this entry.



## 5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	416/436 (95%)	-0.57	1 (0%) 95 95	15, 29, 51, 71	0
1	B	415/436 (95%)	-0.53	0 100 100	14, 25, 48, 77	0
1	C	416/436 (95%)	-0.58	0 100 100	14, 28, 51, 65	0
1	D	415/436 (95%)	-0.57	1 (0%) 95 95	14, 26, 49, 66	0
1	E	415/436 (95%)	-0.60	1 (0%) 95 95	14, 27, 49, 75	0
1	F	415/436 (95%)	-0.44	7 (1%) 73 73	15, 32, 55, 71	0
All	All	2492/2616 (95%)	-0.55	10 (0%) 93 93	14, 28, 51, 77	0

The worst 5 of 10 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	F	201	SER	3.0
1	F	387	HIS	2.7
1	D	200	GLY	2.6
1	E	202	LEU	2.3
1	F	202	LEU	2.3

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
4	EDO	E	502	4/4	0.77	0.16	8.43	42,44,48,54	0
4	EDO	D	502	4/4	0.21	0.27	6.28	40,43,43,45	0
3	AKG	E	501	10/10	0.88	0.17	5.35	31,40,49,52	10
3	AKG	A	501	10/10	0.82	0.19	5.21	35,41,46,47	10
3	AKG	C	501	10/10	0.88	0.18	4.05	29,36,42,42	10
4	EDO	A	502	4/4	0.70	0.18	3.97	45,49,55,57	0
4	EDO	F	502	4/4	0.57	0.27	3.77	51,51,52,55	0
3	AKG	D	501	10/10	0.88	0.14	2.49	26,32,38,41	10
4	EDO	B	502	4/4	0.82	0.20	1.79	35,39,48,55	0
3	AKG	B	501	10/10	0.94	0.12	1.63	24,29,38,41	10
4	EDO	C	502	4/4	0.63	0.17	1.05	43,43,47,48	0
3	AKG	F	501	10/10	0.91	0.11	0.73	24,31,37,39	10
2	NAD	A	500	44/44	0.97	0.08	0.51	14,23,28,29	0
2	NAD	F	500	44/44	0.96	0.08	0.22	18,26,34,43	0
2	NAD	B	500	44/44	0.98	0.08	0.05	14,21,26,28	0
2	NAD	C	500	44/44	0.98	0.08	-0.05	16,21,27,30	0
2	NAD	E	500	44/44	0.98	0.07	-0.38	14,22,27,30	0
2	NAD	D	500	44/44	0.98	0.07	-0.43	12,20,25,27	0

## 6.5 Other polymers

There are no such residues in this entry.