



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 19, 2016 – 08:06 PM GMT

PDB ID : 4XJM  
Title : Crystal structure of 7,8-diaminopelargonic acid synthase (BioA) from Mycobacterium tuberculosis, complexed with a HTS lead compound  
Authors : Finzel, B.C.; Dai, R.  
Deposited on : 2015-01-08  
Resolution : 1.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.1 (RC1), CSD as537be (2016)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026982  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20026982


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	2475 (1.60-1.60)
Clashscore	102246	2732 (1.60-1.60)
Ramachandran outliers	100387	2654 (1.60-1.60)
Sidechain outliers	100360	2653 (1.60-1.60)
RSRZ outliers	91569	2479 (1.60-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	457	
1	B	457	

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 6871 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Adenosylmethionine-8-amino-7-oxononanoate aminotransferase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	429	Total	C	N	O	S	0	8	0
			3253	2071	572	588	22			
1	B	429	Total	C	N	O	S	0	6	0
			3232	2061	566	584	21			

There are 40 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-19	MET	-	initiating methionine	UNP P9WQ81
A	-18	GLY	-	expression tag	UNP P9WQ81
A	-17	SER	-	expression tag	UNP P9WQ81
A	-16	SER	-	expression tag	UNP P9WQ81
A	-15	HIS	-	expression tag	UNP P9WQ81
A	-14	HIS	-	expression tag	UNP P9WQ81
A	-13	HIS	-	expression tag	UNP P9WQ81
A	-12	HIS	-	expression tag	UNP P9WQ81
A	-11	HIS	-	expression tag	UNP P9WQ81
A	-10	HIS	-	expression tag	UNP P9WQ81
A	-9	SER	-	expression tag	UNP P9WQ81
A	-8	SER	-	expression tag	UNP P9WQ81
A	-7	GLY	-	expression tag	UNP P9WQ81
A	-6	LEU	-	expression tag	UNP P9WQ81
A	-5	VAL	-	expression tag	UNP P9WQ81
A	-4	PRO	-	expression tag	UNP P9WQ81
A	-3	ARG	-	expression tag	UNP P9WQ81
A	-2	GLY	-	expression tag	UNP P9WQ81
A	-1	SER	-	expression tag	UNP P9WQ81
A	0	HIS	-	expression tag	UNP P9WQ81
B	-19	MET	-	initiating methionine	UNP P9WQ81
B	-18	GLY	-	expression tag	UNP P9WQ81
B	-17	SER	-	expression tag	UNP P9WQ81
B	-16	SER	-	expression tag	UNP P9WQ81

*Continued on next page...*

Chain	Residue	Modelled	Actual	Comment	Reference
B	-15	HIS	-	expression tag	UNP P9WQ81
B	-14	HIS	-	expression tag	UNP P9WQ81
B	-13	HIS	-	expression tag	UNP P9WQ81
B	-12	HIS	-	expression tag	UNP P9WQ81
B	-11	HIS	-	expression tag	UNP P9WQ81
B	-10	HIS	-	expression tag	UNP P9WQ81
B	-9	SER	-	expression tag	UNP P9WQ81
B	-8	SER	-	expression tag	UNP P9WQ81
B	-7	GLY	-	expression tag	UNP P9WQ81
B	-6	LEU	-	expression tag	UNP P9WQ81
B	-5	VAL	-	expression tag	UNP P9WQ81
B	-4	PRO	-	expression tag	UNP P9WQ81
B	-3	ARG	-	expression tag	UNP P9WQ81
B	-2	GLY	-	expression tag	UNP P9WQ81
B	-1	SER	-	expression tag	UNP P9WQ81
B	0	HIS	-	expression tag	UNP P9WQ81

- [illegible]

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total 29	C 22	N 2	O 4	S 1	0	0
2	B	1	Total 29	C 22	N 2	O 4	S 1	0	0

- 



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			15	8	1	5	1		
3	B	1	Total	C	N	O	P	0	0
			15	8	1	5	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	165	Total	O	0	0
			165	165		
4	B	133	Total	O	0	0
			133	133		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Adenosylmethionine-8-amino-7-oxononanoate aminotransferase



- Molecule 1: Adenosylmethionine-8-amino-7-oxononanoate aminotransferase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	63.06 Å 66.05 Å 203.89 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	101.94 – 1.60 101.94 – 1.60	Depositor EDS
% Data completeness (in resolution range)	99.9 (101.94-1.60) 99.9 (101.94-1.60)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.46 (at 1.60 Å)	Xtriage
Refinement program	REFMAC	Depositor
R, $R_{free}$	0.203 , 0.221 0.201 , 0.218	Depositor DCC
$R_{free}$ test set	5669 reflections (5.28%)	DCC
Wilson B-factor (Å <sup>2</sup> )	17.5	Xtriage
Anisotropy	0.074	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.37 , 40.2	EDS
Estimated twinning fraction	0.028 for k,h,-l	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.46$ , $\langle L^2 \rangle = 0.28$	Xtriage
Outliers	0 of 113109 reflections	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	6871	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	18.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.54% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 41E, PLP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	1.87	7/3348 (0.2%)	1.16	8/4576 (0.2%)
1	B	0.51	3/3330 (0.1%)	0.54	0/4553
All	All	1.37	10/6678 (0.1%)	0.91	8/9129 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	403[A]	ARG	CZ-NH1	-71.54	0.40	1.33
1	A	403[B]	ARG	CZ-NH1	-71.54	0.40	1.33
1	A	403[A]	ARG	CZ-NH2	16.04	1.53	1.33
1	A	403[B]	ARG	CZ-NH2	16.04	1.53	1.33
1	A	64	TRP	CD2-CE2	5.24	1.47	1.41
1	B	153	TRP	CD2-CE2	5.11	1.47	1.41
1	A	398	TRP	CD2-CE2	5.10	1.47	1.41
1	B	398	TRP	CD2-CE2	5.09	1.47	1.41
1	B	139	TRP	CD2-CE2	5.08	1.47	1.41
1	A	153	TRP	CD2-CE2	5.01	1.47	1.41

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	403[A]	ARG	NE-CZ-NH1	-38.98	100.81	120.30

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	403[B]	ARG	NE-CZ-NH1	-38.98	100.81	120.30
1	A	403[A]	ARG	NE-CZ-NH2	-23.10	108.75	120.30
1	A	403[B]	ARG	NE-CZ-NH2	-23.10	108.75	120.30
1	A	403[A]	ARG	NH1-CZ-NH2	-15.34	102.53	119.40
1	A	403[B]	ARG	NH1-CZ-NH2	-15.34	102.53	119.40
1	A	403[A]	ARG	CD-NE-CZ	-11.40	107.64	123.60
1	A	403[B]	ARG	CD-NE-CZ	-11.40	107.64	123.60

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	403[A]	ARG	Sidechain

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3253	0	3242	20	0
1	B	3232	0	3230	25	0
2	A	29	0	26	8	0
2	B	29	0	26	3	0
3	A	15	0	6	5	0
3	B	15	0	6	5	0
4	A	165	0	0	0	0
4	B	133	0	0	0	0
All	All	6871	0	6536	51	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (51) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:283:LYS:NZ	3:A:502:PLP:C4A	1.84	1.41

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:283:LYS:NZ	3:B:502:PLP:C4A	1.91	1.31
1:A:283:LYS:HZ1	3:A:502:PLP:C4A	1.44	1.26
1:B:283:LYS:HZ1	3:B:502:PLP:C4A	1.53	1.16
1:B:283:LYS:HZ2	3:B:502:PLP:C4A	1.62	0.99
1:A:283:LYS:HZ2	3:A:502:PLP:C4A	1.58	0.98
2:A:501:41E:H20	2:A:501:41E:OAD	1.76	0.84
1:A:389:THR:HB	1:A:390:PRO:HD3	1.69	0.73
1:B:65:TRP:HB2	1:B:283:LYS:HD3	1.74	0.69
1:A:65:TRP:HB2	1:A:283:LYS:HD3	1.75	0.68
1:B:178:TRP:HE3	1:B:181:VAL:HG21	1.67	0.58
2:A:501:41E:CAJ	2:A:501:41E:H7	2.32	0.57
1:A:286:THR:HG22	1:A:329[A]:SER:OG	2.06	0.56
1:B:66:THR:HB	1:B:288:GLY:HA2	1.88	0.56
1:B:179:THR:O	1:B:180:ASP:HB2	2.06	0.54
2:B:501:41E:H7	2:B:501:41E:CAJ	2.37	0.54
1:B:286:THR:HG22	1:B:329[A]:SER:OG	2.08	0.53
1:B:178:TRP:CE3	1:B:181:VAL:HG21	2.43	0.53
2:A:501:41E:H5	2:A:501:41E:H7	1.92	0.52
1:A:66:THR:HB	1:A:288:GLY:HA2	1.91	0.51
1:A:178:TRP:HE1	2:A:501:41E:H23	1.75	0.51
1:A:174:MET:HB3	1:A:177:LEU:HD13	1.92	0.51
1:A:88:ASN:HD22	1:A:89:HIS:H	1.58	0.49
1:B:88:ASN:HD22	1:B:89:HIS:H	1.60	0.49
1:A:348:LEU:HD11	1:A:412:TYR:HA	1.97	0.47
1:A:88:ASN:ND2	1:A:89:HIS:H	2.14	0.46
1:B:89:HIS:HA	1:B:323:PRO:HD2	1.97	0.46
1:A:89:HIS:HA	1:A:323:PRO:HD2	1.97	0.46
1:B:170:PRO:HA	1:B:175:HIS:HD2	1.82	0.45
1:B:173:GLY:HA2	2:B:501:41E:CAX	2.47	0.45
1:B:244:CYS:HB3	1:B:249:VAL:O	2.18	0.44
1:B:80:LEU:HD13	1:B:328[B]:VAL:CG2	2.46	0.44
1:B:88:ASN:ND2	1:B:89:HIS:H	2.14	0.43
1:A:110:ILE:HG21	1:A:337:LEU:HD11	2.00	0.43
2:A:501:41E:H13	2:A:501:41E:H17	1.66	0.43
2:A:501:41E:CAK	2:A:501:41E:OAD	2.47	0.43
1:B:88:ASN:N	1:B:88:ASN:HD22	2.16	0.43
1:A:283:LYS:NZ	3:A:502:PLP:C4	2.70	0.42
1:B:97:HIS:CE1	1:B:100:ALA:HB2	2.55	0.42
1:A:283:LYS:HZ2	3:A:502:PLP:C4	2.26	0.41
1:B:283:LYS:NZ	3:B:502:PLP:C4	2.73	0.41
1:A:173:GLY:HA2	2:A:501:41E:CAX	2.51	0.41

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:88:ASN:ND2	1:B:89:HIS:N	2.69	0.41
1:A:218:VAL:HG22	1:A:252:ILE:HD12	2.03	0.41
1:B:283:LYS:HZ2	3:B:502:PLP:C4	2.26	0.41
1:B:252:ILE:HG12	1:B:278:ILE:HB	2.02	0.40
1:A:97:HIS:CE1	1:A:100:ALA:HB2	2.56	0.40
2:A:501:41E:H26	1:B:314:MET:O	2.21	0.40
1:B:174:MET:HB2	2:B:501:41E:CAF	2.51	0.40
1:A:178:TRP:HE3	1:A:181:VAL:HG21	1.87	0.40
1:B:286:THR:HG22	1:B:329[A]:SER:HG	1.85	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	435/457 (95%)	422 (97%)	10 (2%)	3 (1%)	26	8
1	B	433/457 (95%)	420 (97%)	11 (2%)	2 (0%)	34	12
All	All	868/914 (95%)	842 (97%)	21 (2%)	5 (1%)	30	9

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	179	THR
1	A	282	GLY
1	B	282	GLY
1	A	316	GLY
1	B	316	GLY

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	334/346 (96%)	330 (99%)	4 (1%)	78	60
1	B	332/346 (96%)	323 (97%)	9 (3%)	52	23
All	All	666/692 (96%)	653 (98%)	13 (2%)	63	36

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	88	ASN
1	A	153	TRP
1	A	400	ARG
1	A	409	MET
1	B	88	ASN
1	B	122	ASP
1	B	153	TRP
1	B	174	MET
1	B	176	SER
1	B	179	THR
1	B	182	LEU
1	B	274	VAL
1	B	409	MET

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	88	ASN
1	B	88	ASN
1	B	175	HIS
1	B	209	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
2	41E	A	501	-	27,31,31	1.03	1 (3%)	33,42,42	1.46	5 (15%)
3	PLP	A	502	-	15,15,16	3.40	3 (20%)	21,22,23	1.33	3 (14%)
2	41E	B	501	-	27,31,31	0.99	1 (3%)	33,42,42	1.17	2 (6%)
3	PLP	B	502	-	15,15,16	3.26	3 (20%)	21,22,23	1.36	3 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	41E	A	501	-	-	0/15/33/33	0/3/3/3
3	PLP	A	502	-	-	0/6/6/8	0/1/1/1
2	41E	B	501	-	-	0/15/33/33	0/3/3/3
3	PLP	B	502	-	-	0/6/6/8	0/1/1/1

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	501	41E	CAX-NAR	-3.74	1.34	1.41
2	B	501	41E	CAX-NAR	-3.61	1.34	1.41

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	502	PLP	C3-C4	3.91	1.48	1.40
3	B	502	PLP	C3-C4	4.16	1.49	1.40
3	B	502	PLP	C5-C4	6.21	1.47	1.40
3	A	502	PLP	C5-C4	6.27	1.47	1.40
3	B	502	PLP	C3-C2	9.86	1.47	1.40
3	A	502	PLP	C3-C2	10.61	1.48	1.40

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	501	41E	CAX-NAR-CAV	-3.96	119.91	127.49
2	A	501	41E	CAP-CAN-CBB	-2.74	107.17	111.85
3	B	502	PLP	C4A-C4-C5	-2.57	118.26	120.90
2	A	501	41E	CAM-CAL-CAV	-2.19	108.26	113.26
2	A	501	41E	CAA-OAS-CAY	-2.09	112.67	117.51
3	A	502	PLP	C4A-C4-C5	-2.05	118.79	120.90
2	B	501	41E	CAM-CBB-CAO	-2.03	107.79	112.03
3	B	502	PLP	C6-N1-C2	2.10	123.48	119.26
3	B	502	PLP	O3-C3-C4	2.13	123.91	118.11
3	A	502	PLP	C6-N1-C2	2.26	123.80	119.26
3	A	502	PLP	O3P-P-O2P	2.37	116.13	107.44
2	B	501	41E	CAQ-NBC-CAP	3.00	118.07	112.57
2	A	501	41E	CAQ-NBC-CAP	3.73	119.40	112.57

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 21 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	501	41E	8	0
3	A	502	PLP	5	0
2	B	501	41E	3	0
3	B	502	PLP	5	0

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	429/457 (93%)	0.06	16 (3%) 45 43	10, 15, 29, 49	3 (0%)
1	B	429/457 (93%)	0.05	15 (3%) 48 44	10, 17, 33, 57	4 (0%)
All	All	858/914 (93%)	0.05	31 (3%) 46 43	10, 16, 32, 57	7 (0%)

All (31) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	175	HIS	7.4
1	A	176	SER	7.2
1	B	436	LEU	6.7
1	A	436	LEU	6.4
1	A	174	MET	6.3
1	B	173	GLY	6.2
1	A	173	GLY	6.2
1	A	177	LEU	5.2
1	A	175	HIS	5.1
1	A	32	ALA	5.0
1	A	180	ASP	4.4
1	A	33	VAL	4.2
1	B	177	LEU	4.0
1	B	171	HIS	3.9
1	A	65	TRP	3.7
1	B	309	ALA	3.5
1	B	176	SER	3.4
1	B	179	THR	3.4
1	B	385	LEU	3.0
1	A	179	THR	3.0
1	A	171	HIS	3.0
1	B	172	GLY	2.9
1	B	174	MET	2.9
1	A	64	TRP	2.8

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	180	ASP	2.6
1	B	361	ALA	2.4
1	A	178	TRP	2.4
1	A	181	VAL	2.2
1	B	33	VAL	2.2
1	A	25	TYR	2.0
1	B	181	VAL	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
2	41E	A	501	29/29	0.86	0.27	1.50	20,28,36,37	0
2	41E	B	501	29/29	0.90	0.16	0.41	16,23,30,31	0
3	PLP	B	502	15/16	0.97	0.08	0.26	11,11,12,13	0
3	PLP	A	502	15/16	0.97	0.09	0.09	10,11,11,12	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.