



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 10:26 PM GMT

PDB ID : 4XTR
Title : Structure of Get3 bound to the transmembrane domain of Pep12
Authors : Mateja, A.; Paduch, M.; Chang, H.-Y.; Szydlowska, A.; Kossiakoff, A.A.;
Hegde, R.S.; Keenan, R.J.
Deposited on : 2015-01-23
Resolution : 2.05 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.05 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1192 (2.04-2.04)
Clashscore	102246	1269 (2.04-2.04)
Ramachandran outliers	100387	1258 (2.04-2.04)
Sidechain outliers	100360	1258 (2.04-2.04)
RSRZ outliers	91569	1194 (2.04-2.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	354	<div> <div>7%</div> <div>82%</div> <div>14%</div> </div>
1	B	354	<div> <div>4%</div> <div>83%</div> <div>5%</div> <div>12%</div> </div>
2	C	230	<div> <div>8%</div> <div>92%</div> <div>6%</div> </div>
2	E	230	<div> <div>96%</div> <div>2%</div> </div>
3	D	217	<div> <div>10%</div> <div>95%</div> <div>5%</div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	F	217	<div><div></div><div>2%</div><div>96%</div><div>.</div></div>
4	G	37	<div><div></div><div>49%</div><div>46%</div><div>14%</div><div>41%</div></div>

2 Entry composition [i](#)

There are 9 unique types of molecules in this entry. The entry contains 24293 atoms, of which 11564 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ATPase GET3.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	305	Total	C	H	N	O	S	0	2	0
			4815	1530	2400	401	466	18			
1	B	311	Total	C	H	N	O	S	0	1	0
			4873	1543	2432	404	476	18			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	57	ASN	ASP	engineered mutation	UNP Q12154
B	57	ASN	ASP	engineered mutation	UNP Q12154

- Molecule 2 is a protein called Antibody Heavy chain.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	C	217	Total	C	H	N	O	S	0	1	0
			3241	1037	1600	278	320	6			
2	E	222	Total	C	H	N	O	S	0	1	0
			3306	1055	1633	284	328	6			

- Molecule 3 is a protein called Antibody Light chain.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
3	D	216	Total	C	H	N	O	S	0	0	0
			3269	1038	1611	276	338	6			
3	F	216	Total	C	H	N	O	S	0	0	0
			3269	1038	1611	276	338	6			


- Molecule 4 is a protein called Pep12p.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
4	G	22	Total	C	H	N	O	S	0	0	0
			427	141	231	30	23	2			

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
G	252	MET	-	initiating methionine	UNP E7M086
G	253	GLY	-	expression tag	UNP E7M086
G	254	SER	-	expression tag	UNP E7M086
G	255	HIS	-	expression tag	UNP E7M086
G	256	HIS	-	expression tag	UNP E7M086
G	257	HIS	-	expression tag	UNP E7M086
G	258	HIS	-	expression tag	UNP E7M086
G	259	HIS	-	expression tag	UNP E7M086
G	260	HIS	-	expression tag	UNP E7M086
G	261	SER	-	expression tag	UNP E7M086

- Molecule 5 is ADENOSINE-5'-DIPHOSPHATE (three-letter code: ADP) (formula: $C_{10}H_{15}N_5O_{10}P_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	A	1	Total	C	H	N	O	P	
			38	10	11	5	10	2	1
5	B	1	Total	C	H	N	O	P	
			39	10	12	5	10	2	1

- Molecule 6 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	B	1	Total	Mg		
			1	1	0	0

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	1	Total	Mg	0	0
			1	1		

- Molecule 7 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	1	Total	Zn	0	0
			1	1		

- Molecule 8 is ADENOSINE-5'-TRIPHOSPHATE (three-letter code: ATP) (formula: $C_{10}H_{16}N_5O_{13}P_3$).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
8	A	1	Total	C	H	N	O	P	0	1
			42	10	11	5	13	3		
8	B	1	Total	C	H	N	O	P	0	1
			43	10	12	5	13	3		

- Molecule 9 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	A	104	Total	O	0	0
			104	104		
9	B	149	Total	O	0	0
			149	149		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	C	162	Total 162	O 162	0	0
9	D	164	Total 164	O 164	0	0
9	E	211	Total 211	O 211	0	0
9	F	137	Total 137	O 137	0	0
9	G	1	Total 1	O 1	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: ATPase GET3




- Molecule 1: ATPase GET3



- Molecule 2: Antibody Heavy chain




- Molecule 2: Antibody Heavy chain




- Molecule 3: Antibody Light chain







● Molecule 3: Antibody Light chain



● Molecule 4: Pep12p



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	102.87Å 112.03Å 153.69Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	53.95 – 2.05 61.56 – 2.05	Depositor EDS
% Data completeness (in resolution range)	97.8 (53.95-2.05) 92.1 (61.56-2.05)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.39 (at 2.05Å)	Xtriage
Refinement program	PHENIX (phenix.refine: dev_1839)	Depositor
R, R_{free}	0.187 , 0.216 0.185 , 0.213	Depositor DCC
R_{free} test set	5177 reflections (5.03%)	DCC
Wilson B-factor (Å ²)	35.4	Xtriage
Anisotropy	0.458	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 48.4	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Outliers	0 of 109287 reflections	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	24293	wwPDB-VP
Average B, all atoms (Å ²)	56.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.39% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ATP, MG, ZN, ADP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.23	0/2457	0.40	0/3310
1	B	0.24	0/2482	0.41	0/3345
2	C	0.23	0/1682	0.43	0/2293
2	E	0.23	0/1715	0.44	0/2338
3	D	0.23	0/1694	0.41	0/2299
3	F	0.23	0/1694	0.41	0/2299
4	G	0.24	0/200	0.47	0/269
All	All	0.23	0/11924	0.42	0/16153

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2415	2400	2397	6	0
1	B	2441	2432	2431	6	0
2	C	1641	1600	1598	4	0
2	E	1673	1633	1632	0	0
3	D	1658	1611	1611	4	0
3	F	1658	1611	1611	5	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	G	196	231	230	3	0
5	A	27	11	12	0	0
5	B	27	12	12	0	0
6	A	1	0	0	0	0
6	B	1	0	0	0	0
7	A	1	0	0	0	0
8	A	31	11	12	0	0
8	B	31	12	12	0	0
9	A	104	0	0	0	0
9	B	149	0	0	2	0
9	C	162	0	0	2	0
9	D	164	0	0	0	0
9	E	211	0	0	0	0
9	F	137	0	0	2	0
9	G	1	0	0	0	0
All	All	12729	11564	11558	25	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

The worst 5 of 25 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:320:GLU:OE1	1:B:322:ARG:NH1	2.01	0.93
3:D:111:ARG:NH1	3:D:112:THR:O	2.10	0.83
1:A:277:ALA:O	1:A:279:ASN:N	2.17	0.77
3:F:192:HIS:O	3:F:214:ARG:NH1	2.24	0.70
1:B:310:HIS:NE2	9:B:577:HOH:O	2.30	0.63

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	297/354 (84%)	285 (96%)	10 (3%)	2 (1%)	26	15
1	B	304/354 (86%)	297 (98%)	5 (2%)	2 (1%)	26	15
2	C	214/230 (93%)	208 (97%)	6 (3%)	0	100	100
2	E	221/230 (96%)	218 (99%)	3 (1%)	0	100	100
3	D	214/217 (99%)	209 (98%)	5 (2%)	0	100	100
3	F	214/217 (99%)	210 (98%)	4 (2%)	0	100	100
4	G	20/37 (54%)	19 (95%)	1 (5%)	0	100	100
All	All	1484/1639 (90%)	1446 (97%)	34 (2%)	4 (0%)	46	36

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	170	THR
1	A	170	THR
1	A	278	GLU
1	B	285	CYS

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	273/309 (88%)	268 (98%)	5 (2%)	66	62
1	B	276/309 (89%)	268 (97%)	8 (3%)	50	42
2	C	182/193 (94%)	180 (99%)	2 (1%)	80	79
2	E	186/193 (96%)	183 (98%)	3 (2%)	70	67
3	D	191/192 (100%)	188 (98%)	3 (2%)	70	67
3	F	191/192 (100%)	191 (100%)	0	100	100
4	G	22/36 (61%)	22 (100%)	0	100	100
All	All	1321/1424 (93%)	1300 (98%)	21 (2%)	72	67

5 of 21 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	193	ILE
1	B	228	GLN
3	D	202	GLN
1	B	149	ILE
2	E	32	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	61	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 7 ligands modelled in this entry, 3 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	ADP	A	401[B]	6	22,29,29	4.08	7 (31%)	27,45,45	2.77	4 (14%)
8	ATP	A	404[A]	6	24,33,33	2.29	9 (37%)	31,52,52	2.23	9 (29%)
5	ADP	B	401[B]	6	22,29,29	4.10	6 (27%)	27,45,45	2.80	4 (14%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
8	ATP	B	403[A]	6	24,33,33	2.27	8 (33%)	31,52,52	2.24	7 (22%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	ADP	A	401[B]	6	-	0/12/32/32	0/3/3/3
8	ATP	A	404[A]	6	-	0/18/38/38	0/3/3/3
5	ADP	B	401[B]	6	-	0/12/32/32	0/3/3/3
8	ATP	B	403[A]	6	-	0/18/38/38	0/3/3/3

The worst 5 of 30 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	B	401[B]	ADP	O4'-C4'	-5.53	1.32	1.45
5	A	401[B]	ADP	O4'-C4'	-5.51	1.32	1.45
8	B	403[A]	ATP	C2-N1	-5.11	1.24	1.33
8	A	404[A]	ATP	C2-N1	-5.09	1.24	1.33
8	A	404[A]	ATP	C6-N1	-2.50	1.25	1.37

The worst 5 of 24 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	B	401[B]	ADP	N3-C2-N1	-11.84	119.83	128.89
5	A	401[B]	ADP	N3-C2-N1	-11.71	119.92	128.89
8	B	403[A]	ATP	N3-C2-N1	-8.15	122.66	128.89
8	A	404[A]	ATP	N3-C2-N1	-8.05	122.73	128.89
5	A	401[B]	ADP	N6-C6-N1	-6.90	104.39	119.20

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å ²)	Q<0.9
1	A	305/354 (86%)	0.64	25 (8%)	14	16	32, 50, 122, 138	0
1	B	311/354 (87%)	0.51	14 (4%)	37	42	25, 40, 97, 134	0
2	C	217/230 (94%)	0.59	18 (8%)	14	16	24, 42, 97, 131	0
2	E	222/230 (96%)	0.19	0	100	100	22, 35, 58, 102	0
3	D	216/217 (99%)	0.56	21 (9%)	10	11	26, 46, 90, 135	0
3	F	216/217 (99%)	0.28	4 (1%)	70	75	28, 49, 71, 111	0
4	G	22/37 (59%)	4.16	18 (81%)	0	0	90, 112, 124, 143	0
All	All	1509/1639 (92%)	0.53	100 (6%)	22	24	22, 44, 103, 143	0

The worst 5 of 100 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	212	ILE	10.7
4	G	275	LEU	9.0
1	A	126	LEU	8.5
1	A	190	PHE	8.1
4	G	270	TYR	7.8

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
8	ATP	A	404[A]	31/31	0.96	0.14	0.33	30,36,45,48	42
5	ADP	A	401[B]	27/27	0.97	0.14	0.17	30,36,46,48	38
8	ATP	B	403[A]	31/31	0.96	0.15	0.07	25,34,45,52	43
5	ADP	B	401[B]	27/27	0.95	0.15	-0.10	30,36,45,50	39
7	ZN	A	403	1/1	0.94	0.09	-2.01	68,68,68,68	0
6	MG	A	402	1/1	0.94	0.05	-3.54	41,41,41,41	0
6	MG	B	402	1/1	0.94	0.08	-4.22	35,35,35,35	0

6.5 Other polymers [i](#)

There are no such residues in this entry.