



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 07:07 AM GMT

PDB ID : 2ZKF  
Title : Crystal structure of the SRA domain of mouse Np95 in complex with hemi-methylated CpG DNA  
Authors : Arita, K.; Ariyoshi, M.; Tochio, H.; Nakamura, Y.; Shirakawa, M.  
Deposited on : 2008-03-19  
Resolution : 2.55 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.55 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	4549 (2.58-2.50)
Clashscore	102246	5292 (2.58-2.50)
Ramachandran outliers	100387	5194 (2.58-2.50)
Sidechain outliers	100360	5196 (2.58-2.50)
RSRZ outliers	91569	4561 (2.58-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	210	<div> <div></div> <div> <div></div> <div>78%</div> <div>18%</div> <div>.</div> </div> </div>
2	C	11	<div> <div>9%</div> <div>64%</div> <div>36%</div> </div>
3	D	11	<div> <div>9%</div> <div>27%</div> <div>36%</div> <div>36%</div> </div>

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 2137 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called E3 ubiquitin-protein ligase UHRF1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	209	Total	C	N	O	S	0	0	0
			1641	1015	312	307	7			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	404	SER	LYS	ENGINEERED	UNP Q8VDF2

- Molecule 2 is a DNA chain called DNA (5'-D(P\*DCP\*DAP\*DCP\*DCP\*DGP\*DGP\*DAP\*DTP\*DAP\*DGP\*DA)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	11	Total	C	N	O	P	0	0	0
			228	107	46	64	11			

- Molecule 3 is a DNA chain called DNA (5'-D(\*DCP\*DTP\*DAP\*DTP\*DCP\*(5CM)P\*DG P\*DGP\*DTP\*DGP\*DA)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	D	11	Total	C	N	O	P	0	0	0
			223	108	40	65	10			


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	41	Total	O	0	0
			41	41		
4	C	1	Total	O	0	0
			1	1		
4	D	3	Total	O	0	0
			3	3		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: E3 ubiquitin-protein ligase UHRF1



- Molecule 2: DNA (5'-D(P\*DCP\*DAP\*DCP\*DCP\*DGP\*DGP\*DAP\*DTP\*DAP\*DGP\*DA)-3')



- Molecule 3: DNA (5'-D(\*DCP\*DTP\*DAP\*DTP\*DCP\*(5CM)P\*DGP\*DGP\*DTP\*DGP\*DA)-3')



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	58.88Å 58.88Å 182.31Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.46 – 2.55 49.46 – 2.55	Depositor EDS
% Data completeness (in resolution range)	99.8 (49.46-2.55) 99.8 (49.46-2.55)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	10.95 (at 2.54Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, $R_{free}$	0.214 , 0.275 0.222 , 0.281	Depositor DCC
$R_{free}$ test set	532 reflections (5.02%)	DCC
Wilson B-factor (Å <sup>2</sup> )	36.7	Xtriage
Anisotropy	0.063	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 38.6	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Outliers	0 of 11126 reflections	Xtriage
$F_o, F_c$ correlation	0.91	EDS
Total number of atoms	2137	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	25.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.51% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality ⓘ

### 5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: 5CM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.85	0/1682	0.92	5/2273 (0.2%)
2	C	1.55	3/256 (1.2%)	2.24	12/391 (3.1%)
3	D	1.37	0/226	2.46	18/345 (5.2%)
All	All	1.02	3/2164 (0.1%)	1.41	35/3009 (1.2%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	2	DC	OP3-P	-10.12	1.49	1.61
2	C	11	DG	C3'-O3'	-5.29	1.37	1.44
2	C	5	DC	C3'-O3'	-5.06	1.37	1.44

All (35) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	596	ARG	NE-CZ-NH1	10.04	125.32	120.30
3	D	11	DG	O4'-C1'-N9	-9.71	101.21	108.00
2	C	2	DC	OP1-P-OP2	-9.56	105.26	119.60
2	C	3	DA	O4'-C1'-N9	-9.26	101.52	108.00
2	C	10	DA	O4'-C1'-N9	-8.73	101.89	108.00
3	D	8	DG	O4'-C1'-N9	-8.52	102.03	108.00
3	D	3	DT	O4'-C1'-N1	-7.76	102.57	108.00
3	D	8	DG	N3-C2-N2	7.73	125.31	119.90
2	C	5	DC	O4'-C1'-N1	-7.68	102.62	108.00
2	C	9	DT	O4'-C1'-N1	-7.40	102.82	108.00
3	D	10	DT	P-O5'-C5'	-7.12	109.51	120.90
3	D	5	DT	C4-C5-C7	7.00	123.20	119.00
1	A	596	ARG	NE-CZ-NH2	-6.95	116.83	120.30
3	D	5	DT	N3-C2-O2	-6.89	118.16	122.30
2	C	7	DG	O4'-C1'-N9	-6.77	103.26	108.00

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	D	10	DT	C4-C5-C7	6.54	122.92	119.00
1	A	448	ARG	NE-CZ-NH2	-6.26	117.17	120.30
3	D	11	DG	P-O3'-C3'	6.22	127.17	119.70
3	D	5	DT	C6-C5-C7	-6.16	119.20	122.90
1	A	448	ARG	NE-CZ-NH1	5.87	123.23	120.30
2	C	4	DC	C5-C4-N4	5.87	124.31	120.20
2	C	10	DA	N1-C6-N6	-5.68	115.19	118.60
2	C	7	DG	N3-C2-N2	5.63	123.84	119.90
3	D	8	DG	N1-C2-N2	-5.62	111.15	116.20
1	A	469	GLY	N-CA-C	-5.55	99.23	113.10
3	D	10	DT	C4'-C3'-C2'	5.52	108.07	103.10
3	D	8	DG	N1-C6-O6	-5.42	116.65	119.90
2	C	5	DC	C5-C4-N4	5.36	123.95	120.20
3	D	2	DC	P-O3'-C3'	5.24	125.98	119.70
3	D	12	DA	O4'-C1'-N9	5.21	111.65	108.00
3	D	11	DG	C4-N9-C1'	-5.18	119.77	126.50
3	D	10	DT	C6-C5-C7	-5.17	119.80	122.90
3	D	3	DT	N1-C1'-C2'	5.12	122.33	112.60
2	C	8	DA	O4'-C1'-N9	-5.10	104.43	108.00
2	C	12	DA	O4'-C1'-N9	5.00	111.50	108.00

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1641	0	1578	37	0
2	C	228	0	123	3	0
3	D	223	0	128	5	0
4	A	41	0	0	3	0
4	C	1	0	0	0	0
4	D	3	0	0	0	0
All	All	2137	0	1829	42	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 11.

All (42) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:595:THR:HG22	1:A:598:GLY:H	1.26	0.99
1:A:530:ASP:CG	1:A:533:GLN:HE21	1.87	0.78
1:A:595:THR:CG2	1:A:598:GLY:H	1.96	0.77
1:A:529:GLU:OE2	1:A:530:ASP:N	2.25	0.70
1:A:417:ILE:HD12	1:A:417:ILE:H	1.59	0.68
1:A:468:ALA:HB3	3:D:7:5CM:H1'	1.79	0.65
1:A:530:ASP:CB	1:A:533:GLN:HE21	2.09	0.65
1:A:578:LEU:HD23	1:A:579:VAL:N	2.12	0.65
1:A:595:THR:HG23	1:A:597:GLU:H	1.66	0.60
1:A:416:THR:H	1:A:417:ILE:HD12	1.68	0.59
3:D:11:DG:H1'	3:D:12:DA:O5'	2.03	0.58
1:A:494:ASN:ND2	1:A:496:ARG:HH12	2.04	0.56
1:A:529:GLU:CD	1:A:530:ASP:H	2.12	0.52
1:A:595:THR:HG23	1:A:597:GLU:N	2.25	0.51
1:A:469:GLY:O	1:A:471:TYR:N	2.39	0.51
1:A:465:LEU:C	1:A:465:LEU:HD12	2.31	0.51
1:A:450:HIS:NE2	2:C:6:DG:N2	2.59	0.51
1:A:416:THR:H	1:A:417:ILE:CD1	2.24	0.50
1:A:415:CYS:HA	1:A:417:ILE:HD13	1.94	0.49
1:A:595:THR:HG22	1:A:598:GLY:N	2.10	0.48
1:A:539:VAL:HG21	1:A:563:TYR:CD2	2.48	0.47
1:A:465:LEU:HD13	1:A:483:TYR:CE1	2.50	0.47
1:A:529:GLU:N	1:A:529:GLU:CD	2.68	0.47
1:A:530:ASP:CB	1:A:533:GLN:NE2	2.79	0.46
3:D:2:DC:H2''	3:D:3:DT:OP2	2.15	0.46
1:A:592:GLU:O	1:A:595:THR:HB	2.16	0.45
1:A:438:ARG:NH2	1:A:449:PRO:O	2.45	0.45
1:A:555:GLU:O	4:A:37:HOH:O	2.21	0.45
1:A:523:GLU:HG3	4:A:46:HOH:O	2.17	0.44
2:C:9:DT:H2''	2:C:10:DA:C8	2.53	0.44
1:A:592:GLU:HG2	1:A:594:TRP:CZ2	2.53	0.44
1:A:418:VAL:HB	1:A:419:PRO:CD	2.48	0.44
1:A:602:THR:HG23	1:A:607:LEU:HB2	2.00	0.43
1:A:490:ASP:OD1	1:A:492:SER:OG	2.33	0.42
1:A:417:ILE:CD1	1:A:417:ILE:H	2.29	0.42
1:A:474:ASP:OD2	3:D:7:5CM:N3	2.53	0.41
2:C:11:DG:H2''	2:C:12:DA:OP2	2.20	0.41
3:D:11:DG:CI'	3:D:12:DA:O5'	2.68	0.41

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:434:MET:HE3	1:A:608:THR:C	2.41	0.41
1:A:530:ASP:HB3	1:A:533:GLN:HE21	1.84	0.40
1:A:609:MET:HE1	4:A:37:HOH:O	2.21	0.40
1:A:527:GLU:HB3	1:A:567:LYS:HD3	2.02	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	207/210 (99%)	193 (93%)	12 (6%)	2 (1%)	19	33

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	417	ILE
1	A	470	GLY

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	170/171 (99%)	159 (94%)	11 (6%)	21	37

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	417	ILE
1	A	450	HIS
1	A	492	SER
1	A	494	ASN
1	A	529	GLU
1	A	547	LYS
1	A	584	LEU
1	A	585	ARG
1	A	595	THR
1	A	596	ARG
1	A	604	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	494	ASN
1	A	509	ASN
1	A	533	GLN
1	A	604	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	5CM	D	7	3	13,21,22	2.45	5 (38%)	17,30,33	2.62	3 (17%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical

component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	5CM	D	7	3	-	0/3/21/22	0/2/2/2

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	D	7	5CM	C5A-C5	-4.95	1.41	1.51
3	D	7	5CM	C5-C4	-3.24	1.36	1.41
3	D	7	5CM	O4'-C1'	3.15	1.49	1.42
3	D	7	5CM	C6-N1	3.27	1.39	1.35
3	D	7	5CM	C4-N3	4.30	1.41	1.35

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	D	7	5CM	O4'-C1'-N1	-8.21	93.51	107.72
3	D	7	5CM	N4-C4-N3	3.20	121.59	116.95
3	D	7	5CM	C2'-C1'-N1	5.06	126.47	114.16

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	D	7	5CM	2	0

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	209/210 (99%)	-0.04	2 (0%) 84 87	15, 22, 30, 36	0
2	C	11/11 (100%)	0.24	1 (9%) 11 13	26, 29, 37, 49	0
3	D	10/11 (90%)	0.34	1 (10%) 9 10	24, 33, 55, 59	0
All	All	230/232 (99%)	-0.01	4 (1%) 73 77	15, 22, 33, 59	0

All (4) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	409	VAL	2.6
1	A	613	GLU	2.4
2	C	12	DA	2.4
3	D	12	DA	2.3

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
3	5CM	D	7	20/21	0.98	0.15	-	14,18,24,27	0

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.