



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 02:30 PM GMT

PDB ID : 3ZKD
Title : CRYSTAL STRUCTURE OF THE ATPASE REGION OF Mycobacterium tuberculosis GyrB WITH AMPPNP
Authors : Agrawal, A.; Roue, M.; Spitzfaden, C.; Petrella, S.; Aubry, A.; Volker, C.; Mossakowska, D.; Hann, M.; Bax, B.; Mayer, C.
Deposited on : 2013-01-22
Resolution : 2.95 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	2184 (3.00-2.92)
Clashscore	102246	2552 (3.00-2.92)
Ramachandran outliers	100387	2468 (3.00-2.92)
Sidechain outliers	100360	2471 (3.00-2.92)
RSRZ outliers	91569	2201 (3.00-2.92)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	432	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 1%, green 65%, yellow 21%, grey 13%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> % 65% 21% • 13% </div> </div>
1	B	432	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, green 66%, yellow 20%, grey 13%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> % 66% 20% • 13% </div> </div>
1	C	432	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 1%, green 69%, yellow 20%, grey 10%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> % 69% 20% • 10% </div> </div>
1	D	432	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 1%, green 69%, yellow 17%, grey 12%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> % 69% 17% • 12% </div> </div>
1	E	432	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 1%, green 72%, yellow 16%, grey 12%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> % 72% 16% 12% </div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	432	<div><div></div><div>63%22%13%</div></div>
1	G	432	<div>%<div><div></div><div>68%18%13%</div></div></div>
1	H	432	<div>%<div><div></div><div>68%19%13%</div></div></div>

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 23192 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA GYRASE SUBUNIT B.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	377	Total	C	N	O	S	0	0	0
			2855	1789	503	557	6			
1	B	375	Total	C	N	O	S	0	0	0
			2843	1783	498	556	6			
1	C	390	Total	C	N	O	S	0	0	0
			2927	1835	519	567	6			
1	D	379	Total	C	N	O	S	0	0	0
			2817	1766	489	556	6			
1	E	382	Total	C	N	O	S	0	0	0
			2878	1803	506	563	6			
1	F	376	Total	C	N	O	S	0	0	0
			2821	1768	497	550	6			
1	G	378	Total	C	N	O	S	0	0	0
			2845	1786	497	556	6			
1	H	378	Total	C	N	O	S	0	0	0
			2835	1776	494	559	6			

There are 40 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-4	GLY	-	EXPRESSION TAG	UNP I6WX66
A	-3	PRO	-	EXPRESSION TAG	UNP I6WX66
A	-2	LEU	-	EXPRESSION TAG	UNP I6WX66
A	-1	GLY	-	EXPRESSION TAG	UNP I6WX66
A	0	SER	-	EXPRESSION TAG	UNP I6WX66
B	-4	GLY	-	EXPRESSION TAG	UNP I6WX66
B	-3	PRO	-	EXPRESSION TAG	UNP I6WX66
B	-2	LEU	-	EXPRESSION TAG	UNP I6WX66
B	-1	GLY	-	EXPRESSION TAG	UNP I6WX66
B	0	SER	-	EXPRESSION TAG	UNP I6WX66
C	-4	GLY	-	EXPRESSION TAG	UNP I6WX66
C	-3	PRO	-	EXPRESSION TAG	UNP I6WX66
C	-2	LEU	-	EXPRESSION TAG	UNP I6WX66

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	-1	GLY	-	EXPRESSION TAG	UNP I6WX66
C	0	SER	-	EXPRESSION TAG	UNP I6WX66
D	-4	GLY	-	EXPRESSION TAG	UNP I6WX66
D	-3	PRO	-	EXPRESSION TAG	UNP I6WX66
D	-2	LEU	-	EXPRESSION TAG	UNP I6WX66
D	-1	GLY	-	EXPRESSION TAG	UNP I6WX66
D	0	SER	-	EXPRESSION TAG	UNP I6WX66
E	-4	GLY	-	EXPRESSION TAG	UNP I6WX66
E	-3	PRO	-	EXPRESSION TAG	UNP I6WX66
E	-2	LEU	-	EXPRESSION TAG	UNP I6WX66
E	-1	GLY	-	EXPRESSION TAG	UNP I6WX66
E	0	SER	-	EXPRESSION TAG	UNP I6WX66
F	-4	GLY	-	EXPRESSION TAG	UNP I6WX66
F	-3	PRO	-	EXPRESSION TAG	UNP I6WX66
F	-2	LEU	-	EXPRESSION TAG	UNP I6WX66
F	-1	GLY	-	EXPRESSION TAG	UNP I6WX66
F	0	SER	-	EXPRESSION TAG	UNP I6WX66
G	-4	GLY	-	EXPRESSION TAG	UNP I6WX66
G	-3	PRO	-	EXPRESSION TAG	UNP I6WX66
G	-2	LEU	-	EXPRESSION TAG	UNP I6WX66
G	-1	GLY	-	EXPRESSION TAG	UNP I6WX66
G	0	SER	-	EXPRESSION TAG	UNP I6WX66
H	-4	GLY	-	EXPRESSION TAG	UNP I6WX66
H	-3	PRO	-	EXPRESSION TAG	UNP I6WX66
H	-2	LEU	-	EXPRESSION TAG	UNP I6WX66
H	-1	GLY	-	EXPRESSION TAG	UNP I6WX66
H	0	SER	-	EXPRESSION TAG	UNP I6WX66

- Molecule 2 is PHOSPHOAMINOPHOSPHONIC ACID-ADENYLATE ESTER (three-letter code: ANP) (formula: $C_{10}H_{17}N_6O_{12}P_3$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			31	10	6	12	3		
2	B	1	Total	C	N	O	P	0	0
			31	10	6	12	3		
2	C	1	Total	C	N	O	P	0	0
			31	10	6	12	3		
2	D	1	Total	C	N	O	P	0	0
			31	10	6	12	3		
2	E	1	Total	C	N	O	P	0	0
			31	10	6	12	3		
2	F	1	Total	C	N	O	P	0	0
			31	10	6	12	3		
2	G	1	Total	C	N	O	P	0	0
			31	10	6	12	3		
2	H	1	Total	C	N	O	P	0	0
			31	10	6	12	3		

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	G	1	Total	Mg	0	0
			1	1		
3	D	1	Total	Mg	0	0
			1	1		
3	E	1	Total	Mg	0	0
			1	1		
3	H	1	Total	Mg	0	0
			1	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total 1	Mg 1	0	0
3	C	1	Total 1	Mg 1	0	0
3	A	1	Total 1	Mg 1	0	0
3	F	1	Total 1	Mg 1	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	20	Total 20	O 20	0	0
4	B	11	Total 11	O 11	0	0
4	C	20	Total 20	O 20	0	0
4	D	8	Total 8	O 8	0	0
4	E	15	Total 15	O 15	0	0
4	F	12	Total 12	O 12	0	0
4	G	22	Total 22	O 22	0	0
4	H	7	Total 7	O 7	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: DNA GYRASE SUBUNIT B




• Molecule 1: DNA GYRASE SUBUNIT B




• Molecule 1: DNA GYRASE SUBUNIT B







• Molecule 1: DNA GYRASE SUBUNIT B




• Molecule 1: DNA GYRASE SUBUNIT B




• Molecule 1: DNA GYRASE SUBUNIT B



- Molecule 1: DNA GYRASE SUBUNIT B



● Molecule 1: DNA GYRASE SUBUNIT B



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	71.95Å 97.85Å 138.07Å 108.21° 105.18° 91.04°	Depositor
Resolution (Å)	24.81 – 2.95 24.82 – 2.95	Depositor EDS
% Data completeness (in resolution range)	84.6 (24.81-2.95) 75.8 (24.82-2.95)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.13 (at 2.94Å)	Xtriage
Refinement program	BUSTER 2.11.4	Depositor
R, R_{free}	0.182 , 0.257 0.196 , 0.281	Depositor DCC
R_{free} test set	2459 reflections (4.20%)	DCC
Wilson B-factor (Å ²)	55.9	Xtriage
Anisotropy	0.705	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.28 , 30.6	EDS
Estimated twinning fraction	0.168 for h,-k,-h-l 0.015 for -h,k,-k-l 0.009 for -h,-k,h+k+l	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 61071 reflections	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	23192	wwPDB-VP
Average B, all atoms (Å ²)	80.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.45% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MG, ANP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.44	0/2907	0.69	0/3940
1	B	0.44	0/2896	0.67	0/3928
1	C	0.46	0/2981	0.71	0/4044
1	D	0.43	0/2869	0.68	0/3901
1	E	0.44	0/2932	0.69	0/3979
1	F	0.45	0/2873	0.70	0/3900
1	G	0.45	0/2898	0.70	0/3932
1	H	0.44	0/2886	0.68	0/3919
All	All	0.44	0/23242	0.69	0/31543

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2
1	C	0	1
1	H	0	1
All	All	0	4

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	342	ASP	Mainchain,Peptide
1	C	342	ASP	Mainchain

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Group
1	H	342	ASP	Mainchain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2855	0	2774	51	0
1	B	2843	0	2764	57	0
1	C	2927	0	2837	46	0
1	D	2817	0	2690	42	0
1	E	2878	0	2780	37	0
1	F	2821	0	2726	54	0
1	G	2845	0	2756	44	0
1	H	2835	0	2723	43	0
2	A	31	0	13	3	0
2	B	31	0	13	1	0
2	C	31	0	13	1	0
2	D	31	0	13	1	0
2	E	31	0	13	2	0
2	F	31	0	13	1	0
2	G	31	0	13	2	0
2	H	31	0	13	1	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
3	C	1	0	0	0	0
3	D	1	0	0	0	0
3	E	1	0	0	0	0
3	F	1	0	0	0	0
3	G	1	0	0	0	0
3	H	1	0	0	0	0
4	A	20	0	0	0	0
4	B	11	0	0	0	0
4	C	20	0	0	0	0
4	D	8	0	0	0	0
4	E	15	0	0	0	0
4	F	12	0	0	0	0
4	G	22	0	0	0	0
4	H	7	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	23192	0	22154	351	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

The worst 5 of 351 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:93:ILE:HG13	1:B:94:PRO:HD2	1.53	0.90
1:C:308:ILE:HD13	1:C:372:LYS:HD3	1.61	0.82
1:D:312:GLU:CG	1:D:373:THR:HB	2.13	0.78
1:D:312:GLU:HG3	1:D:373:THR:HB	1.67	0.76
2:A:601:ANP:H8	2:A:601:ANP:H5'2	1.68	0.75

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	371/432 (86%)	342 (92%)	22 (6%)	7 (2%)	10	40
1	B	369/432 (85%)	346 (94%)	22 (6%)	1 (0%)	46	81
1	C	384/432 (89%)	359 (94%)	21 (6%)	4 (1%)	19	58
1	D	373/432 (86%)	351 (94%)	15 (4%)	7 (2%)	10	40
1	E	376/432 (87%)	355 (94%)	20 (5%)	1 (0%)	46	81
1	F	370/432 (86%)	349 (94%)	19 (5%)	2 (0%)	34	74
1	G	372/432 (86%)	349 (94%)	18 (5%)	5 (1%)	15	51
1	H	372/432 (86%)	349 (94%)	19 (5%)	4 (1%)	17	56
All	All	2987/3456 (86%)	2800 (94%)	156 (5%)	31 (1%)	19	58

5 of 31 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	343	PRO
1	C	343	PRO
1	H	343	PRO
1	A	87	ALA
1	A	124	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	297/350 (85%)	280 (94%)	17 (6%)	25	62
1	B	298/350 (85%)	273 (92%)	25 (8%)	14	42
1	C	301/350 (86%)	280 (93%)	21 (7%)	19	53
1	D	289/350 (83%)	270 (93%)	19 (7%)	21	55
1	E	299/350 (85%)	283 (95%)	16 (5%)	27	64
1	F	292/350 (83%)	261 (89%)	31 (11%)	8	29
1	G	296/350 (85%)	272 (92%)	24 (8%)	15	44
1	H	293/350 (84%)	273 (93%)	20 (7%)	20	54
All	All	2365/2800 (84%)	2192 (93%)	173 (7%)	17	50

5 of 173 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	335	ARG
1	E	402	ASP
1	H	165	LYS
1	D	388	CYS
1	E	247	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 21 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	302	HIS
1	D	102	GLN
1	H	195	GLN
1	C	89	HIS
1	H	316	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 16 ligands modelled in this entry, 8 are monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	ANP	A	601	3	27,33,33	4.13	12 (44%)	30,52,52	2.46	9 (30%)
2	ANP	B	601	3	27,33,33	3.99	12 (44%)	30,52,52	2.32	8 (26%)
2	ANP	C	601	3	27,33,33	4.19	10 (37%)	30,52,52	2.48	6 (20%)
2	ANP	D	601	3	27,33,33	4.14	11 (40%)	30,52,52	2.38	6 (20%)
2	ANP	E	601	3	27,33,33	4.09	10 (37%)	30,52,52	2.49	5 (16%)
2	ANP	F	601	3	27,33,33	4.13	10 (37%)	30,52,52	2.70	7 (23%)
2	ANP	G	601	3	27,33,33	4.11	11 (40%)	30,52,52	2.36	7 (23%)
2	ANP	H	601	3	27,33,33	4.03	11 (40%)	30,52,52	2.54	5 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	ANP	A	601	3	-	0/12/38/38	0/3/3/3
2	ANP	B	601	3	-	0/12/38/38	0/3/3/3
2	ANP	C	601	3	-	0/12/38/38	0/3/3/3
2	ANP	D	601	3	-	0/12/38/38	0/3/3/3
2	ANP	E	601	3	-	0/12/38/38	0/3/3/3
2	ANP	F	601	3	-	1/12/38/38	0/3/3/3
2	ANP	G	601	3	-	0/12/38/38	0/3/3/3
2	ANP	H	601	3	-	0/12/38/38	0/3/3/3

The worst 5 of 87 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	F	601	ANP	PG-N3B	-4.74	1.50	1.63
2	B	601	ANP	PG-N3B	-4.38	1.51	1.63
2	C	601	ANP	PG-N3B	-4.31	1.51	1.63
2	G	601	ANP	PG-N3B	-4.28	1.52	1.63
2	H	601	ANP	PG-N3B	-4.19	1.52	1.63

The worst 5 of 53 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	F	601	ANP	N3-C2-N1	-11.98	119.72	128.89
2	E	601	ANP	N3-C2-N1	-11.61	120.00	128.89
2	C	601	ANP	N3-C2-N1	-11.41	120.16	128.89
2	H	601	ANP	N3-C2-N1	-11.19	120.32	128.89
2	D	601	ANP	N3-C2-N1	-10.97	120.49	128.89

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	F	601	ANP	O1G-PG-N3B-PB

There are no ring outliers.

8 monomers are involved in 12 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	601	ANP	3	0
2	B	601	ANP	1	0
2	C	601	ANP	1	0
2	D	601	ANP	1	0
2	E	601	ANP	2	0
2	F	601	ANP	1	0
2	G	601	ANP	2	0
2	H	601	ANP	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	377/432 (87%)	-0.20	4 (1%) 82 65	47, 74, 130, 156	0
1	B	375/432 (86%)	-0.27	1 (0%) 94 87	46, 75, 110, 129	0
1	C	390/432 (90%)	-0.30	4 (1%) 84 67	39, 71, 129, 179	0
1	D	379/432 (87%)	-0.28	3 (0%) 87 73	51, 83, 124, 160	0
1	E	382/432 (88%)	-0.23	6 (1%) 74 55	41, 70, 135, 171	0
1	F	376/432 (87%)	-0.30	2 (0%) 91 81	44, 72, 123, 160	0
1	G	378/432 (87%)	-0.30	4 (1%) 82 65	41, 70, 132, 155	0
1	H	378/432 (87%)	-0.21	6 (1%) 74 55	49, 82, 130, 149	0
All	All	3035/3456 (87%)	-0.26	30 (0%) 84 67	39, 75, 128, 179	0

The worst 5 of 30 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	340	ASP	5.4
1	E	339	LYS	5.1
1	E	338	LEU	4.8
1	C	339	LYS	4.5
1	H	338	LEU	3.9

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	MG	D	602	1/1	0.97	0.20	1.91	60,60,60,60	0
3	MG	A	602	1/1	0.99	0.19	1.67	38,38,38,38	0
3	MG	E	602	1/1	0.98	0.17	1.07	51,51,51,51	0
3	MG	H	602	1/1	0.99	0.17	0.80	37,37,37,37	0
2	ANP	E	601	31/31	0.98	0.16	0.20	42,51,57,58	0
2	ANP	D	601	31/31	0.98	0.17	0.12	49,76,86,87	0
2	ANP	G	601	31/31	0.99	0.16	0.12	44,49,55,58	0
2	ANP	C	601	31/31	0.99	0.16	0.09	48,59,66,69	0
3	MG	G	602	1/1	0.98	0.16	-0.10	35,35,35,35	0
2	ANP	H	601	31/31	0.98	0.17	-0.11	51,80,88,92	0
2	ANP	F	601	31/31	0.98	0.15	-0.25	38,54,60,61	0
3	MG	C	602	1/1	0.98	0.15	-0.29	35,35,35,35	0
2	ANP	B	601	31/31	0.98	0.15	-0.33	52,64,69,71	0
2	ANP	A	601	31/31	0.98	0.15	-0.33	49,58,62,63	0
3	MG	B	602	1/1	0.98	0.13	-0.80	37,37,37,37	0
3	MG	F	602	1/1	0.96	0.20	-	48,48,48,48	0

6.5 Other polymers [i](#)

There are no such residues in this entry.