



Full wwPDB X-ray Structure Validation Report ⓘ

May 23, 2016 – 10:49 AM EDT

PDB ID : 4ZS4
Title : Crystal Structure of the Inactive Alpha-kinase Domain of Myosin-II Heavy Chain Kinase A (D756A) Complexed with ATP
Authors : Ye, Q.; Jia, Z.
Deposited on : 2015-05-13
Resolution : 2.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.1 (RC1), CSD as537be (2016)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20027674
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20027674


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	2919 (2.40-2.40)
Clashscore	102246	3407 (2.40-2.40)
Ramachandran outliers	100387	3351 (2.40-2.40)
Sidechain outliers	100360	3352 (2.40-2.40)
RSRZ outliers	91569	2928 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	307	<div> <div>4%</div> <div> <div></div> <div>75%</div> <div>7%</div> <div>18%</div> </div> </div>
1	B	307	<div> <div>3%</div> <div> <div></div> <div>78%</div> <div>6%</div> <div>16%</div> </div> </div>

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 4428 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Myosin heavy chain kinase A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	253	Total	C	N	O	S	0	0	0
			2004	1284	336	372	12			
1	B	257	Total	C	N	O	S	0	0	0
			2028	1300	341	375	12			

There are 36 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	535	MET	-	initiating methionine	UNP P42527
A	536	GLY	-	expression tag	UNP P42527
A	537	GLY	-	expression tag	UNP P42527
A	538	HIS	-	expression tag	UNP P42527
A	539	HIS	-	expression tag	UNP P42527
A	540	HIS	-	expression tag	UNP P42527
A	541	HIS	-	expression tag	UNP P42527
A	542	HIS	-	expression tag	UNP P42527
A	543	HIS	-	expression tag	UNP P42527
A	544	GLY	-	expression tag	UNP P42527
A	545	GLU	-	expression tag	UNP P42527
A	546	ASN	-	expression tag	UNP P42527
A	547	LEU	-	expression tag	UNP P42527
A	548	TYR	-	expression tag	UNP P42527
A	549	PHE	-	expression tag	UNP P42527
A	550	GLN	-	expression tag	UNP P42527
A	551	GLY	-	expression tag	UNP P42527
A	756	ALA	ASP	engineered mutation	UNP P42527
B	535	MET	-	initiating methionine	UNP P42527
B	536	GLY	-	expression tag	UNP P42527
B	537	GLY	-	expression tag	UNP P42527
B	538	HIS	-	expression tag	UNP P42527
B	539	HIS	-	expression tag	UNP P42527
B	540	HIS	-	expression tag	UNP P42527
B	541	HIS	-	expression tag	UNP P42527

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	542	HIS	-	expression tag	UNP P42527
B	543	HIS	-	expression tag	UNP P42527
B	544	GLY	-	expression tag	UNP P42527
B	545	GLU	-	expression tag	UNP P42527
B	546	ASN	-	expression tag	UNP P42527
B	547	LEU	-	expression tag	UNP P42527
B	548	TYR	-	expression tag	UNP P42527
B	549	PHE	-	expression tag	UNP P42527
B	550	GLN	-	expression tag	UNP P42527
B	551	GLY	-	expression tag	UNP P42527
B	756	ALA	ASP	engineered mutation	UNP P42527

- Molecule 2 is PHOSPHATE ION (three-letter code: PO4) (formula: O₄P).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total O P 5 4 1	0	0
2	B	1	Total O P 5 4 1	0	0

- Molecule 3 is ZINC ION (three-letter code: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	B	1	Total Zn 1 1	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Zn	0	0
			1	1		

- Molecule 4 is ADENOSINE-5'-TRIPHOSPHATE (three-letter code: ATP) (formula: $C_{10}H_{16}N_5O_{13}P_3$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O	P	0	0
			31	10	5	13	3		
4	B	1	Total	C	N	O	P	0	0
			31	10	5	13	3		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	151	Total	O	0	0
			151	151		
5	B	171	Total	O	0	0
			171	171		

- Molecule 1: Myosin heavy chain kinase A



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	83.21Å 109.74Å 77.84Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 2.40 44.85 – 2.39	Depositor EDS
% Data completeness (in resolution range)	97.0 (50.00-2.40) 97.0 (44.85-2.39)	Depositor EDS
R_{merge}	0.14	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.82 (at 2.39Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.178 , 0.223 0.184 , 0.224	Depositor DCC
R_{free} test set	1409 reflections (5.31%)	DCC
Wilson B-factor (Å ²)	32.8	Xtriage
Anisotropy	0.117	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 38.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	4428	wwPDB-VP
Average B, all atoms (Å ²)	36.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 13.73% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, PO4, ATP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.54	0/2046	0.73	2/2760 (0.1%)
1	B	0.56	0/2071	0.71	3/2795 (0.1%)
All	All	0.55	0/4117	0.72	5/5555 (0.1%)

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	747	LEU	CA-CB-CG	-5.92	101.70	115.30
1	B	671	ARG	NE-CZ-NH2	-5.84	117.38	120.30
1	A	773	ASP	CB-CG-OD1	5.18	122.96	118.30
1	B	747	LEU	CA-CB-CG	-5.13	103.50	115.30
1	B	773	ASP	CB-CG-OD1	5.07	122.86	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2004	0	2022	20	0
1	B	2028	0	2053	12	0
2	A	5	0	0	0	0
2	B	5	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	31	0	12	2	0
4	B	31	0	12	0	0
5	A	151	0	0	5	0
5	B	171	0	0	1	0
All	All	4428	0	4099	32	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (32) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:671:ARG:NH1	5:A:1001:HOH:O	2.09	0.79
1:A:611:GLY:HA2	1:A:616:LEU:HD11	1.71	0.72
1:A:571:LYS:NZ	1:B:751:GLN:OE1	2.22	0.72
1:B:655:ALA:HA	1:B:659:LEU:HD12	1.72	0.72
1:A:668:MET:HE3	1:A:691:SER:H	1.62	0.64
1:B:753:ILE:HD13	1:B:787:ILE:HG12	1.81	0.62
1:A:610:LEU:HG	1:A:616:LEU:HD21	1.82	0.60
1:A:611:GLY:HA2	1:A:616:LEU:CD1	2.32	0.58
4:A:903:ATP:O2G	4:A:903:ATP:O2B	2.23	0.57
1:A:589:GLY:HA3	4:A:903:ATP:O2B	2.07	0.54
1:A:671:ARG:NE	5:A:1002:HOH:O	2.21	0.53
1:A:743:PHE:CE1	1:A:747:LEU:HD13	2.43	0.53
1:B:743:PHE:CE1	1:B:747:LEU:HD13	2.44	0.52
1:A:614:THR:HG22	5:A:1087:HOH:O	2.09	0.52
1:A:614:THR:O	1:A:615:LYS:HB2	2.11	0.51
1:B:656:SER:H	1:B:659:LEU:HD12	1.75	0.50
1:A:555:GLU:HB3	1:A:556:THR:HB	1.96	0.48
1:A:743:PHE:CD1	1:A:747:LEU:HD13	2.49	0.47
1:A:671:ARG:NH2	5:A:1002:HOH:O	2.41	0.46
1:B:675:ASN:HB3	5:B:1149:HOH:O	2.15	0.46
1:A:563:TRP:CH2	1:A:574:ARG:HD2	2.51	0.46
1:B:743:PHE:CD1	1:B:747:LEU:HD13	2.52	0.45
1:A:671:ARG:NH1	5:A:1003:HOH:O	2.27	0.44
1:B:563:TRP:CH2	1:B:574:ARG:HD2	2.53	0.44
1:B:771:THR:O	1:B:783:GLY:HA2	2.17	0.44
1:A:652:GLU:HG2	1:A:653:GLN:N	2.33	0.44
1:A:771:THR:O	1:A:783:GLY:HA2	2.18	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:656:SER:N	1:B:659:LEU:HD12	2.33	0.43
1:A:555:GLU:HB3	1:A:556:THR:CA	2.49	0.42
1:A:554:SER:HA	1:A:555:GLU:HA	1.94	0.41
1:B:677:PHE:CE2	1:B:682:PRO:HG3	2.56	0.41
1:B:655:ALA:HA	1:B:659:LEU:CD1	2.46	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	247/307 (80%)	242 (98%)	5 (2%)	0	100	100
1	B	253/307 (82%)	246 (97%)	7 (3%)	0	100	100
All	All	500/614 (81%)	488 (98%)	12 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	224/269 (83%)	224 (100%)	0	100	100
1	B	226/269 (84%)	225 (100%)	1 (0%)	93	98
All	All	450/538 (84%)	449 (100%)	1 (0%)	95	99

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	571	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 2 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	PO4	A	901	-	4,4,4	0.89	0	6,6,6	0.30	0
4	ATP	A	903	-	26,33,33	1.07	2 (7%)	26,52,52	1.82	3 (11%)
2	PO4	B	901	-	4,4,4	1.15	0	6,6,6	0.28	0
4	ATP	B	903	-	26,33,33	0.89	1 (3%)	26,52,52	1.97	3 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means

no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PO4	A	901	-	-	0/0/0/0	0/0/0/0
4	ATP	A	903	-	-	0/18/38/38	0/3/3/3
2	PO4	B	901	-	-	0/0/0/0	0/0/0/0
4	ATP	B	903	-	-	0/18/38/38	0/3/3/3

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	903	ATP	C5-C4	2.44	1.46	1.40
4	A	903	ATP	O4'-C1'	2.45	1.44	1.41
4	A	903	ATP	C5-C4	3.19	1.47	1.40

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	903	ATP	N3-C2-N1	-8.02	122.57	128.87
4	A	903	ATP	N3-C2-N1	-6.95	123.41	128.87
4	B	903	ATP	C4'-O4'-C1'	2.04	111.81	109.64
4	A	903	ATP	N6-C6-N1	2.54	122.78	118.52
4	A	903	ATP	O2A-PA-O3A	2.67	116.69	105.27
4	B	903	ATP	O3G-PG-O2G	2.98	118.39	107.44

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	903	ATP	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	253/307 (82%)	-0.21	11 (4%) 39 40	17, 31, 73, 107	0
1	B	257/307 (83%)	-0.25	8 (3%) 52 52	18, 31, 70, 107	0
All	All	510/614 (83%)	-0.23	19 (3%) 45 46	17, 31, 73, 107	0

All (19) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	703	SER	6.4
1	B	613	THR	5.3
1	A	554	SER	4.4
1	B	554	SER	4.1
1	A	612	THR	4.0
1	A	653	GLN	3.6
1	A	555	GLU	3.6
1	B	614	THR	3.4
1	A	702	SER	3.4
1	B	704	ASN	3.4
1	A	703	SER	3.3
1	B	702	SER	3.3
1	A	553	SER	3.2
1	A	654	GLN	3.1
1	A	556	THR	2.8
1	B	612	THR	2.5
1	A	706	GLN	2.1
1	B	706	GLN	2.1
1	A	704	ASN	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	PO4	A	901	5/5	0.97	0.12	-0.26	37,42,48,50	0
2	PO4	B	901	5/5	0.99	0.10	-1.01	27,28,29,32	0
4	ATP	B	903	31/31	0.96	0.09	-1.54	27,35,48,58	0
3	ZN	A	902	1/1	0.98	0.07	-1.56	32,32,32,32	0
4	ATP	A	903	31/31	0.96	0.09	-1.56	27,32,64,67	0
3	ZN	B	902	1/1	1.00	0.06	-1.62	23,23,23,23	0

6.5 Other polymers [i](#)

There are no such residues in this entry.