



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 19, 2016 – 09:11 PM GMT

PDB ID : 4ZSZ
Title : Structure of a fusion protein with a helix linker, 2ARH-3-3KAW-3.0
Authors : Lai, Y.-T.; Yeates, T.O.
Deposited on : 2015-05-14
Resolution : 4.25 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : unknown
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026982
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20026982


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 4.25 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1043 (4.92-3.60)
Clashscore	102246	1145 (4.92-3.60)
Ramachandran outliers	100387	1088 (4.92-3.60)
Sidechain outliers	100360	1072 (4.92-3.60)
RSRZ outliers	91569	1047 (4.92-3.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	296	 78% 18% . .
1	B	296	 75% 21% . .

2 Entry composition

There is only 1 type of molecule in this entry. The entry contains 4794 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Uncharacterized Fusion Protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	294	Total	C	N	O	S	0	0	0
			2397	1528	421	434	14			
1	B	294	Total	C	N	O	S	0	0	0
			2397	1528	421	434	14			

There are 26 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	138	PHE	TYR	mutation'	UNP O67778
A	158	ALA	GLU	mutation'	UNP O67778
A	162	ALA	LYS	mutation'	UNP O67778
A	166	ALA	GLU	mutation'	UNP O67778
A	197	GLU	-	linker	UNP O67778
A	198	GLU	-	linker	UNP O67778
A	199	ILE	-	linker	UNP O67778
A	200	TYR	-	linker	UNP O67778
A	208	HIS	GLN	engineered mutation	UNP Q9I208
A	209	HIS	ALA	engineered mutation	UNP Q9I208
A	212	HIS	ARG	engineered mutation	UNP Q9I208
A	290	ALA	GLU	mutation'	UNP Q9I208
A	293	HIS	ARG	mutation'	UNP Q9I208
B	138	PHE	TYR	mutation'	UNP O67778
B	158	ALA	GLU	mutation'	UNP O67778
B	162	ALA	LYS	mutation'	UNP O67778
B	166	ALA	GLU	mutation'	UNP O67778
B	197	GLU	-	linker	UNP O67778
B	198	GLU	-	linker	UNP O67778
B	199	ILE	-	linker	UNP O67778
B	200	TYR	-	linker	UNP O67778
B	208	HIS	GLN	engineered mutation	UNP Q9I208
B	209	HIS	ALA	engineered mutation	UNP Q9I208
B	212	HIS	ARG	engineered mutation	UNP Q9I208
B	290	ALA	GLU	mutation'	UNP Q9I208

Continued on next page...


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	293	HIS	ARG	mutation'	UNP Q9I208


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Uncharacterized Fusion Protein



• Molecule 1: Uncharacterized Fusion Protein



4 Data and refinement statistics

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, α , β , γ	112.93Å 112.93Å 150.01Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	46.49 – 4.25 81.93 – 4.25	Depositor EDS
% Data completeness (in resolution range)	98.9 (46.49-4.25) 98.9 (81.93-4.25)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.23 (at 4.30Å)	Xtriage
Refinement program	PHENIX	Depositor
R, R_{free}	0.265 , 0.286 0.265 , 0.286	Depositor DCC
R_{free} test set	401 reflections (4.98%)	DCC
Wilson B-factor (Å ²)	163.8	Xtriage
Anisotropy	0.010	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 12.3	EDS
Estimated twinning fraction	0.045 for -h,-k,l	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.31$	Xtriage
Outliers	0 of 8061 reflections	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	4794	wwPDB-VP
Average B, all atoms (Å ²)	49.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.35% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.31	0/2453	0.47	0/3299
1	B	0.30	0/2453	0.45	0/3299
All	All	0.30	0/4906	0.46	0/6598

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2397	0	2358	33	0
1	B	2397	0	2358	43	0
All	All	4794	0	4716	73	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (73) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:225:SER:HB3	1:A:228:GLN:HG3	1.65	0.78
1:A:215:GLU:OE2	1:A:240:ARG:NH2	2.15	0.77

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:201:GLU:HA	1:B:204:ARG:HD3	1.72	0.71
1:B:213:ALA:HB1	1:B:281:CYS:HA	1.77	0.66
1:B:56:ARG:HB3	1:B:60:TRP:HD1	1.63	0.63
1:B:23:ARG:NH1	1:B:43:GLY:O	2.38	0.57
1:B:19:GLU:CD	1:B:80:SER:HB2	2.28	0.54
1:A:48:LEU:HD13	1:A:66:VAL:HG23	1.90	0.52
1:B:215:GLU:OE2	1:B:240:ARG:NH2	2.42	0.52
1:B:52:VAL:HG12	1:B:64:VAL:HG23	1.92	0.51
1:A:274:ASP:HA	1:A:279:ARG:NH1	2.26	0.50
1:A:65:GLU:OE2	1:A:149:LYS:NZ	2.39	0.50
1:B:181:ASP:HB3	1:B:184:LEU:HB2	1.94	0.50
1:A:115:VAL:HG12	1:A:175:PHE:CZ	2.47	0.49
1:B:204:ARG:HD2	1:B:249:ARG:HH12	1.76	0.49
1:A:138:PHE:O	1:A:140:PRO:HD3	2.13	0.49
1:B:64:VAL:HG13	1:B:97:ILE:HG13	1.94	0.49
1:B:140:PRO:O	1:B:141:GLU:HG3	2.13	0.48
1:A:225:SER:OG	1:A:226:ARG:N	2.47	0.48
1:A:208:HIS:HD1	1:A:211:ILE:HD12	1.78	0.48
1:A:26:ARG:HH11	1:A:28:SER:HB3	1.79	0.48
1:B:53:PHE:CE2	1:B:55:GLY:HA2	2.49	0.48
1:B:38:PHE:HB2	1:B:50:LEU:HB2	1.96	0.48
1:A:259:LEU:HA	1:A:259:LEU:HD23	1.72	0.48
1:A:5:GLU:O	1:A:9:LYS:HB3	2.14	0.48
1:A:53:PHE:CE2	1:A:55:GLY:HA2	2.49	0.47
1:B:225:SER:HB3	1:B:228:GLN:HG3	1.95	0.47
1:B:48:LEU:HD13	1:B:66:VAL:HG23	1.95	0.47
1:B:51:LYS:HB3	1:B:65:GLU:HB3	1.97	0.47
1:B:274:ASP:HA	1:B:279:ARG:NH1	2.30	0.47
1:A:192:TYR:O	1:A:195:LEU:HB3	2.15	0.46
1:B:56:ARG:HB3	1:B:60:TRP:CD1	2.46	0.46
1:B:110:GLU:HB3	1:B:115:VAL:HG21	1.98	0.46
1:B:223:GLU:CD	1:B:223:GLU:H	2.19	0.46
1:A:257:CYS:HB3	1:A:292:CYS:SG	2.56	0.45
1:A:19:GLU:CD	1:A:80:SER:HB2	2.37	0.45
1:B:55:GLY:HA3	1:B:62:PRO:HG3	1.99	0.45
1:B:48:LEU:HA	1:B:68:GLY:O	2.16	0.45
1:B:165:LEU:HD12	1:B:294:PRO:HB3	1.98	0.45
1:A:271:CYS:O	1:A:278:GLU:HB3	2.16	0.45
1:B:224:SER:O	1:B:229:ARG:NH2	2.50	0.45
1:B:259:LEU:HD23	1:B:259:LEU:HA	1.74	0.45
1:A:56:ARG:HH11	1:B:32:PHE:HE2	1.63	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:248:ARG:HG2	1:A:248:ARG:H	1.63	0.43
1:A:91:SER:HA	1:A:131:TYR:OH	2.19	0.43
1:B:259:LEU:HD23	1:B:262:ARG:HD3	2.01	0.43
1:A:133:TYR:O	1:A:152:ALA:HA	2.18	0.43
1:B:34:GLU:HG2	1:B:34:GLU:H	1.70	0.43
1:B:78:PHE:CE1	1:B:104:ASP:HB2	2.52	0.43
1:A:268:ALA:O	1:A:272:ASP:HB2	2.19	0.43
1:B:116:PRO:HA	1:B:117:PRO:HD3	1.88	0.43
1:B:271:CYS:O	1:B:278:GLU:HB3	2.19	0.43
1:A:53:PHE:CE2	1:B:60:TRP:HH2	2.35	0.42
1:A:117:PRO:HB2	1:A:134:PHE:HB3	2.01	0.42
1:A:211:ILE:HD13	1:A:240:ARG:HG3	2.02	0.42
1:B:91:SER:HB3	1:B:129:LEU:HD23	2.01	0.42
1:A:292:CYS:HA	1:A:295:LEU:HD12	2.01	0.42
1:A:58:PRO:HD3	1:B:144:MET:HB2	2.02	0.42
1:A:101:TYR:HH	1:A:138:PHE:HD2	1.67	0.42
1:A:116:PRO:HA	1:A:117:PRO:HD3	1.90	0.42
1:B:176:ILE:HD13	1:B:176:ILE:HA	1.87	0.41
1:B:193:ASN:O	1:B:197:GLU:HG2	2.21	0.41
1:B:237:ASP:HA	1:B:240:ARG:HB2	2.02	0.41
1:A:111:LEU:HD11	1:A:150:ILE:HD11	2.02	0.41
1:B:57:LYS:HA	1:B:58:PRO:HA	1.82	0.41
1:B:34:GLU:HB2	1:B:53:PHE:CE1	2.55	0.41
1:B:270:ARG:HB3	1:B:270:ARG:NH1	2.36	0.41
1:A:110:GLU:HB3	1:A:115:VAL:HG21	2.03	0.41
1:B:204:ARG:HD2	1:B:249:ARG:NH1	2.36	0.40
1:B:125:GLU:O	1:B:129:LEU:HD13	2.21	0.40
1:A:139:ILE:HG22	1:A:141:GLU:HG2	2.03	0.40
1:B:53:PHE:CE2	1:B:60:TRP:HB3	2.56	0.40
1:A:259:LEU:HD22	1:A:263:TYR:HE2	1.86	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	292/296 (99%)	283 (97%)	8 (3%)	1 (0%)	46	83
1	B	292/296 (99%)	286 (98%)	6 (2%)	0	100	100
All	All	584/592 (99%)	569 (97%)	14 (2%)	1 (0%)	52	86

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	140	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	248/250 (99%)	231 (93%)	17 (7%)	19	59
1	B	248/250 (99%)	228 (92%)	20 (8%)	15	52
All	All	496/500 (99%)	459 (92%)	37 (8%)	17	56

All (37) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	9	LYS
1	A	11	LEU
1	A	23	ARG
1	A	26	ARG
1	A	34	GLU
1	A	59	TYR
1	A	85	LYS
1	A	105	LYS
1	A	139	ILE
1	A	184	LEU
1	A	204	ARG
1	A	223	GLU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	226	ARG
1	A	246	LEU
1	A	248	ARG
1	A	259	LEU
1	A	274	ASP
1	B	11	LEU
1	B	23	ARG
1	B	26	ARG
1	B	29	GLN
1	B	34	GLU
1	B	59	TYR
1	B	66	VAL
1	B	85	LYS
1	B	105	LYS
1	B	141	GLU
1	B	184	LEU
1	B	196	GLU
1	B	223	GLU
1	B	226	ARG
1	B	229	ARG
1	B	247	GLU
1	B	259	LEU
1	B	270	ARG
1	B	274	ASP
1	B	293	HIS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	294/296 (99%)	-0.09	1 (0%) 94 92	31, 50, 50, 50	0
1	B	294/296 (99%)	-0.18	1 (0%) 94 92	31, 50, 50, 50	0
All	All	588/592 (99%)	-0.13	2 (0%) 94 92	31, 50, 50, 50	0

All (2) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	146	GLY	3.1
1	B	119	LEU	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.