



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 13, 2017 – 08:49 pm GMT

PDB ID : 4BF2
Title : Crystal Structures of Ask1-inhibitor Complexes
Authors : Singh, O.; Shillings, A.; Craggs, P.; Wall, I.; Rowland, P.; Skarzynski, T.;
Hobbs, C.I.; Hardwick, P.; Tanner, R.; Blunt, M.; Witty, D.R.; Smith, K.J.
Deposited on : 2013-03-13
Resolution : 2.11 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : trunk28620
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.11 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	4988 (2.14-2.10)
Clashscore	112137	5557 (2.14-2.10)
Ramachandran outliers	110173	5504 (2.14-2.10)
Sidechain outliers	110143	5505 (2.14-2.10)
RSRZ outliers	101464	5021 (2.14-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	334	<div> <div>4%</div> <div>72%</div> <div>24%</div> </div>
1	B	334	<div> <div>9%</div> <div>71%</div> <div>5%</div> <div>25%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	ACT	B	1942	-	-	-	X

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 4234 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called MITOGEN-ACTIVATED PROTEIN KINASE KINASE KINASE 5.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	253	Total	C	N	O	S	0	0	0
			1973	1270	325	369	9			
1	B	252	Total	C	N	O	S	0	0	0
			1961	1262	325	365	9			

There are 34 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	644	MET	-	EXPRESSION TAG	UNP Q99683
A	645	ASP	-	EXPRESSION TAG	UNP Q99683
A	646	TYR	-	EXPRESSION TAG	UNP Q99683
A	647	LYS	-	EXPRESSION TAG	UNP Q99683
A	648	ASP	-	EXPRESSION TAG	UNP Q99683
A	649	ASP	-	EXPRESSION TAG	UNP Q99683
A	650	ASP	-	EXPRESSION TAG	UNP Q99683
A	651	ASP	-	EXPRESSION TAG	UNP Q99683
A	652	LYS	-	EXPRESSION TAG	UNP Q99683
A	653	GLU	-	EXPRESSION TAG	UNP Q99683
A	654	ASN	-	EXPRESSION TAG	UNP Q99683
A	655	LEU	-	EXPRESSION TAG	UNP Q99683
A	656	TYR	-	EXPRESSION TAG	UNP Q99683
A	657	PHE	-	EXPRESSION TAG	UNP Q99683
A	658	GLN	-	EXPRESSION TAG	UNP Q99683
A	659	GLY	-	EXPRESSION TAG	UNP Q99683
A	838	GLU	THR	ENGINEERED MUTATION	UNP Q99683
B	644	MET	-	EXPRESSION TAG	UNP Q99683
B	645	ASP	-	EXPRESSION TAG	UNP Q99683
B	646	TYR	-	EXPRESSION TAG	UNP Q99683
B	647	LYS	-	EXPRESSION TAG	UNP Q99683
B	648	ASP	-	EXPRESSION TAG	UNP Q99683
B	649	ASP	-	EXPRESSION TAG	UNP Q99683
B	650	ASP	-	EXPRESSION TAG	UNP Q99683

Continued on next page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	651	ASP	-	EXPRESSION TAG	UNP Q99683
B	652	LYS	-	EXPRESSION TAG	UNP Q99683
B	653	GLU	-	EXPRESSION TAG	UNP Q99683
B	654	ASN	-	EXPRESSION TAG	UNP Q99683
B	655	LEU	-	EXPRESSION TAG	UNP Q99683
B	656	TYR	-	EXPRESSION TAG	UNP Q99683
B	657	PHE	-	EXPRESSION TAG	UNP Q99683
B	658	GLN	-	EXPRESSION TAG	UNP Q99683
B	659	GLY	-	EXPRESSION TAG	UNP Q99683
B	838	GLU	THR	ENGINEERED MUTATION	UNP Q99683

- # STU
-
- The diagram shows the chemical structure of STU, a complex organic molecule. The structure features a central benzene ring (C6-C19) substituted with two phenyl groups (C1-C5 and C10-C14) and two pyrrole rings (C9-C10 and C20-C21). The molecule also contains two nitrogen atoms (N1, N2, N3, N4) and two oxygen atoms (O5, O6). The structure is shown with displacement ellipsoids at the 50% probability level, and the atoms are labeled with their respective symbols and numbers.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total 35	C 28	N 4	O 3	0	0
2	B	1	Total 35	C 28	N 4	O 3	0	0

- 



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		

- Molecule 4 is ACETATE ION (three-letter code: ACT) (formula: $C_2H_3O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	C	O	0	0
			4	2	2		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	131	Total 131	O 131	0	0
5	B	89	Total 89	O 89	0	0

4 Data and refinement statistics

Property	Value	Source
Space group	P 65 2 2	Depositor
Cell constants a, b, c, α , β , γ	78.25Å 78.25Å 421.38Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	13.36 – 2.11 33.45 – 2.11	Depositor EDS
% Data completeness (in resolution range)	93.4 (13.36-2.11) 93.0 (33.45-2.11)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.11 (at 2.12Å)	Xtriage
Refinement program	BUSTER 2.11.1	Depositor
R, R_{free}	0.206 , 0.247 0.219 , 0.264	Depositor DCC
R_{free} test set	1728 reflections (4.25%)	DCC
Wilson B-factor (Å ²)	43.4	Xtriage
Anisotropy	0.152	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 61.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	4234	wwPDB-VP
Average B, all atoms (Å ²)	53.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.00% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, STU, ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.51	0/2015	0.63	0/2717
1	B	0.46	0/2002	0.63	0/2699
All	All	0.49	0/4017	0.63	0/5416

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1973	0	1941	6	0
1	B	1961	0	1945	6	0
2	A	35	0	26	2	0
2	B	35	0	26	1	0
3	A	6	0	8	0	0
4	B	4	0	3	0	0
5	A	131	0	0	0	0
5	B	89	0	0	1	0
All	All	4234	0	3949	15	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

The worst 5 of 15 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:1941:STU:H261	2:B:1941:STU:H16	1.71	0.73
1:B:869:CYS:HB3	1:B:880:PRO:HG3	1.77	0.66
1:B:679:GLU:H	1:B:679:GLU:CD	2.06	0.57
1:A:821:SER:OG	1:A:822:ASP:N	2.36	0.56
1:B:846:MET:HG2	1:B:850:ILE:HB	1.87	0.56

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	247/334 (74%)	237 (96%)	9 (4%)	1 (0%)	38	35
1	B	246/334 (74%)	237 (96%)	9 (4%)	0	100	100
All	All	493/668 (74%)	474 (96%)	18 (4%)	1 (0%)	51	51

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	822	ASP

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	206/289 (71%)	202 (98%)	4 (2%)	62	67
1	B	206/289 (71%)	200 (97%)	6 (3%)	48	50
All	All	412/578 (71%)	402 (98%)	10 (2%)	54	58

5 of 10 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	728	LEU
1	B	733	LYS
1	B	905	GLU
1	A	936	GLU
1	B	737	ILE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	786	GLN
1	B	756	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length

(or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	STU	A	1940	-	29,42,42	2.03	11 (37%)	27,68,68	1.59	7 (25%)
3	GOL	A	1941	-	5,5,5	0.54	0	5,5,5	1.35	1 (20%)
2	STU	B	1941	-	29,42,42	2.10	11 (37%)	27,68,68	1.23	3 (11%)
4	ACT	B	1942	-	1,3,3	3.47	1 (100%)	0,3,3	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	STU	A	1940	-	-	0/4/42/42	0/0/8/8
3	GOL	A	1941	-	-	0/4/4/4	0/0/0/0
2	STU	B	1941	-	-	0/4/42/42	0/0/8/8
4	ACT	B	1942	-	-	0/0/0/0	0/0/0/0

The worst 5 of 23 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	1941	STU	C6-C19	-3.92	1.37	1.42
2	A	1940	STU	C11-C18	-3.23	1.38	1.42
2	B	1941	STU	C11-C18	-2.01	1.40	1.42
2	A	1940	STU	C3-C2	2.01	1.42	1.38
2	A	1940	STU	C3-C4	2.03	1.41	1.36

The worst 5 of 11 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1940	STU	C9-N1-C8	-4.07	109.39	113.78
2	B	1941	STU	C14-C13-C12	-3.31	115.23	120.81
2	A	1940	STU	C3-C4-C5	-2.38	116.79	120.81
3	A	1941	GOL	C3-C2-C1	-2.35	102.18	111.52
2	A	1940	STU	C9-C10-C7	-2.30	107.56	109.61

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1940	STU	2	0
2	B	1941	STU	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	253/334 (75%)	0.11	12 (4%) 32 38	31, 45, 76, 114	0
1	B	252/334 (75%)	0.59	31 (12%) 5 6	35, 57, 87, 104	0
All	All	505/668 (75%)	0.35	43 (8%) 11 15	31, 50, 82, 114	0

The worst 5 of 43 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	854	GLY	6.0
1	B	855	PRO	5.7
1	B	714	ARG	5.5
1	B	851	ILE	5.4
1	B	892	PHE	5.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
4	ACT	B	1942	4/4	0.71	0.24	9.65	64,64,65,66	0
3	GOL	A	1941	6/6	0.96	0.14	0.63	50,51,51,51	0
2	STU	A	1940	35/35	0.98	0.13	-0.26	32,33,34,35	0
2	STU	B	1941	35/35	0.97	0.09	-1.14	37,38,39,41	0

6.5 Other polymers [i](#)

There are no such residues in this entry.