



# wwPDB X-ray Structure Validation Summary Report ⓘ

Oct 29, 2017 – 01:08 PM EDT

PDB ID : 3KL5  
Title : Structure Analysis of a Xylanase From Glycosyl Hydrolase Family Thirty: Carbohydrate Ligand Complexes Reveal this Family of Enzymes Unique Mechanism of Substrate Specificity and Recognition  
Authors : St John, F.J.; Hurlbert, J.C.; Pozharski, E.  
Deposited on : unknown  
Resolution : 2.59 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.2 (RC1), CSD as538be (2017)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20030345  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20030345


# 1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.59 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	2542 (2.60-2.60)
Clashscore	112137	2895 (2.60-2.60)
Ramachandran outliers	110173	2848 (2.60-2.60)
Sidechain outliers	110143	2848 (2.60-2.60)
RSRZ outliers	101464	2550 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	401	<div> <div>6%</div> <div>85% 12% .</div> </div>
1	B	401	<div> <div>%</div> <div>86% 10% ..</div> </div>
1	C	401	<div> <div>2%</div> <div>84% 13% .</div> </div>
1	D	401	<div> <div>53%</div> <div>70% 20% . 8%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit crite-

ria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	XYP	A	404	-	-	-	X
3	XYP	B	404	-	-	-	X
3	XYP	C	403	-	-	-	X
3	XYP	C	404	-	-	-	X

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 12477 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Glucuronoxylanase xynC.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	389	Total	C	N	O	S	0	1	0
			3109	1966	552	582	9			
1	B	390	Total	C	N	O	S	0	3	0
			3136	1984	557	586	9			
1	C	389	Total	C	N	O	S	0	1	0
			3110	1967	552	582	9			
1	D	367	Total	C	N	O	S	0	0	0
			2948	1878	516	545	9			

There are 44 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MET	-	EXPRESSION TAG	UNP Q45070
A	392	LEU	-	EXPRESSION TAG	UNP Q45070
A	393	GLU	-	EXPRESSION TAG	UNP Q45070
A	394	HIS	-	EXPRESSION TAG	UNP Q45070
A	395	HIS	-	EXPRESSION TAG	UNP Q45070
A	396	HIS	-	EXPRESSION TAG	UNP Q45070
A	397	HIS	-	EXPRESSION TAG	UNP Q45070
A	398	HIS	-	EXPRESSION TAG	UNP Q45070
A	399	HIS	-	EXPRESSION TAG	UNP Q45070
A	400	HIS	-	EXPRESSION TAG	UNP Q45070
A	401	HIS	-	EXPRESSION TAG	UNP Q45070
B	1	MET	-	EXPRESSION TAG	UNP Q45070
B	392	LEU	-	EXPRESSION TAG	UNP Q45070
B	393	GLU	-	EXPRESSION TAG	UNP Q45070
B	394	HIS	-	EXPRESSION TAG	UNP Q45070
B	395	HIS	-	EXPRESSION TAG	UNP Q45070
B	396	HIS	-	EXPRESSION TAG	UNP Q45070
B	397	HIS	-	EXPRESSION TAG	UNP Q45070
B	398	HIS	-	EXPRESSION TAG	UNP Q45070
B	399	HIS	-	EXPRESSION TAG	UNP Q45070
B	400	HIS	-	EXPRESSION TAG	UNP Q45070

*Continued on next page...*

*Continued from previous page...*


Chain	Residue	Modelled	Actual	Comment	Reference
B	401	HIS	-	EXPRESSION TAG	UNP Q45070
C	1	MET	-	EXPRESSION TAG	UNP Q45070
C	392	LEU	-	EXPRESSION TAG	UNP Q45070
C	393	GLU	-	EXPRESSION TAG	UNP Q45070
C	394	HIS	-	EXPRESSION TAG	UNP Q45070
C	395	HIS	-	EXPRESSION TAG	UNP Q45070
C	396	HIS	-	EXPRESSION TAG	UNP Q45070
C	397	HIS	-	EXPRESSION TAG	UNP Q45070
C	398	HIS	-	EXPRESSION TAG	UNP Q45070
C	399	HIS	-	EXPRESSION TAG	UNP Q45070
C	400	HIS	-	EXPRESSION TAG	UNP Q45070
C	401	HIS	-	EXPRESSION TAG	UNP Q45070
D	1	MET	-	EXPRESSION TAG	UNP Q45070
D	392	LEU	-	EXPRESSION TAG	UNP Q45070
D	393	GLU	-	EXPRESSION TAG	UNP Q45070
D	394	HIS	-	EXPRESSION TAG	UNP Q45070
D	395	HIS	-	EXPRESSION TAG	UNP Q45070
D	396	HIS	-	EXPRESSION TAG	UNP Q45070
D	397	HIS	-	EXPRESSION TAG	UNP Q45070
D	398	HIS	-	EXPRESSION TAG	UNP Q45070
D	399	HIS	-	EXPRESSION TAG	UNP Q45070
D	400	HIS	-	EXPRESSION TAG	UNP Q45070
D	401	HIS	-	EXPRESSION TAG	UNP Q45070

- Molecule 2 is 4-O-METHYL-ALPHA-D-GLUCURONIC ACID (three-letter code: GCV) (formula: C<sub>7</sub>H<sub>12</sub>O<sub>7</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			14	7	7		
2	B	1	Total	C	O	0	0
			14	7	7		
2	B	1	Total	C	O	0	0
			14	7	7		
2	C	1	Total	C	O	0	0
			14	7	7		

- Molecule 3 is BETA-D-XYLOPYRANOSE (three-letter code: XYP) (formula: C<sub>5</sub>H<sub>10</sub>O<sub>5</sub>).





Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			9	5	4		
3	A	1	Total	C	O	0	0
			9	5	4		
3	B	1	Total	C	O	0	0
			9	5	4		
3	B	1	Total	C	O	0	0
			9	5	4		
3	B	1	Total	C	O	0	0
			9	5	4		
3	B	1	Total	C	O	0	0
			8	5	3		
3	C	1	Total	C	O	0	0
			9	5	4		
3	C	1	Total	C	O	0	0
			9	5	4		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	10	Total 10	O 10	0	0
4	B	17	Total 17	O 17	0	0
4	C	19	Total 19	O 19	0	0
4	D	1	Total 1	O 1	0	0



- Molecule 1: Glucuronoxylanase xynC







## 4 Data and refinement statistics

Property	Value	Source
Space group	P 2 <sub>1</sub> 2 <sub>1</sub> 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	137.72Å 194.01Å 65.71Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 2.59 44.67 – 2.59	Depositor EDS
% Data completeness (in resolution range)	97.5 (50.00-2.59) 97.6 (44.67-2.59)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.32 (at 2.58Å)	Xtriage
Refinement program	REFMAC 5.5.0102	Depositor
R, $R_{free}$	0.241 , 0.290 0.237 , 0.284	Depositor DCC
$R_{free}$ test set	2775 reflections (5.36%)	DCC
Wilson B-factor (Å <sup>2</sup> )	65.9	Xtriage
Anisotropy	0.068	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.31 , 43.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	12477	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	73.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.68% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: XYP, GCV

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.59	0/3196	0.63	4/4353 (0.1%)
1	B	0.58	0/3224	0.70	5/4390 (0.1%)
1	C	0.59	2/3197 (0.1%)	0.70	5/4354 (0.1%)
1	D	0.43	0/3031	0.51	0/4123
All	All	0.55	2/12648 (0.0%)	0.64	14/17220 (0.1%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	162	ASN	CG-OD1	-5.33	1.12	1.24
1	C	162	ASN	CG-ND2	-5.22	1.19	1.32

The worst 5 of 14 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	169	ARG	NE-CZ-NH1	-14.19	113.21	120.30
1	C	60	ARG	NE-CZ-NH2	12.89	126.75	120.30
1	C	60	ARG	NE-CZ-NH1	-12.73	113.93	120.30
1	B	169	ARG	NE-CZ-NH2	12.08	126.34	120.30
1	C	169	ARG	NE-CZ-NH2	-8.86	115.87	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within

the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3109	0	2963	25	0
1	B	3136	0	2993	26	0
1	C	3110	0	2965	34	0
1	D	2948	0	2810	56	0
2	A	14	0	10	1	0
2	B	28	0	20	1	0
2	C	14	0	10	0	0
3	A	18	0	15	2	0
3	B	35	0	28	3	0
3	C	18	0	15	1	0
4	A	10	0	0	2	0
4	B	17	0	0	0	0
4	C	19	0	0	4	0
4	D	1	0	0	0	0
All	All	12477	0	11829	128	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 128 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:254:HIS:HD2	4:C:411:HOH:O	1.57	0.87
1:B:244[B]:PRO:O	1:B:245[B]:GLU:HB2	1.76	0.82
1:B:244[B]:PRO:O	1:B:245[B]:GLU:CB	2.30	0.76
1:D:144:ALA:HB3	1:D:149:TRP:HD1	1.53	0.74
1:B:315:TYR:OH	1:C:222:GLY:HA2	1.86	0.74

There are no symmetry-related clashes.

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	388/401 (97%)	373 (96%)	15 (4%)	0	100	100
1	B	391/401 (98%)	373 (95%)	15 (4%)	3 (1%)	22	44
1	C	388/401 (97%)	373 (96%)	14 (4%)	1 (0%)	44	70
1	D	357/401 (89%)	316 (88%)	36 (10%)	5 (1%)	13	26
All	All	1524/1604 (95%)	1435 (94%)	80 (5%)	9 (1%)	32	53

5 of 9 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	268	TRP
1	B	245[A]	GLU
1	B	245[B]	GLU
1	B	4	ASP
1	D	58	GLU

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	331/341 (97%)	321 (97%)	10 (3%)	46	74
1	B	334/341 (98%)	321 (96%)	13 (4%)	37	65
1	C	331/341 (97%)	320 (97%)	11 (3%)	43	70
1	D	312/341 (92%)	293 (94%)	19 (6%)	22	43
All	All	1308/1364 (96%)	1255 (96%)	53 (4%)	36	63

5 of 53 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	58	GLU
1	C	230	VAL
1	D	261	ASP
1	C	60	ARG
1	C	178	TYR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 12 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	76	HIS
1	D	145	HIS
1	D	313	ASN
1	D	61	ASN
1	D	309	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

### 5.6 Ligand geometry ⓘ

12 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	GCV	A	402	3	11,14,14	1.70	1 (9%)	15,20,20	1.38	1 (6%)
3	XYP	A	403	3,2	9,9,10	1.88	4 (44%)	10,12,14	1.46	2 (20%)
3	XYP	A	404	3	9,9,10	1.11	1 (11%)	11,12,14	0.85	1 (9%)
2	GCV	B	402	3	11,14,14	0.83	0	15,20,20	1.88	2 (13%)
3	XYP	B	403	3,2	9,9,10	1.08	0	10,12,14	1.39	2 (20%)
3	XYP	B	404	3	9,9,10	1.23	2 (22%)	11,12,14	1.51	4 (36%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	GCV	B	405	3	11,14,14	1.83	2 (18%)	15,20,20	1.96	3 (20%)
3	XYP	B	406	3,2	9,9,10	2.51	5 (55%)	10,12,14	1.29	2 (20%)
3	XYP	B	407	3	8,8,10	1.23	0	7,10,14	1.44	1 (14%)
2	GCV	C	402	3	11,14,14	1.81	3 (27%)	15,20,20	2.58	4 (26%)
3	XYP	C	403	3,2	9,9,10	2.53	4 (44%)	10,12,14	2.09	2 (20%)
3	XYP	C	404	3	9,9,10	1.76	2 (22%)	11,12,14	1.54	1 (9%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GCV	A	402	3	-	0/2/26/26	0/1/1/1
3	XYP	A	403	3,2	-	0/0/13/17	0/1/1/1
3	XYP	A	404	3	-	0/0/14/17	0/1/1/1
2	GCV	B	402	3	-	0/2/26/26	0/1/1/1
3	XYP	B	403	3,2	-	0/0/13/17	0/1/1/1
3	XYP	B	404	3	-	0/0/14/17	0/1/1/1
2	GCV	B	405	3	-	0/2/26/26	0/1/1/1
3	XYP	B	406	3,2	-	0/0/13/17	0/1/1/1
3	XYP	B	407	3	-	0/0/11/17	0/1/1/1
2	GCV	C	402	3	-	0/2/26/26	0/1/1/1
3	XYP	C	403	3,2	-	0/0/13/17	0/1/1/1
3	XYP	C	404	3	-	0/0/14/17	0/1/1/1

The worst 5 of 24 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	406	XYP	C4B-C3B	2.04	1.54	1.52
3	C	403	XYP	C4B-C3B	2.10	1.55	1.52
3	A	403	XYP	O5B-C5B	2.12	1.47	1.43
2	C	402	GCV	O1-C1	2.14	1.46	1.39
3	A	404	XYP	C4B-C3B	2.17	1.55	1.52

The worst 5 of 25 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	403	XYP	C1B-C2B-C3B	-5.90	101.02	111.13
2	B	402	GCV	C3-C4-C5	-5.57	103.05	110.14
3	B	403	XYP	O4A-C1B-C2B	-3.20	94.81	111.47

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	405	GCV	O4-C4-C3	-3.08	102.87	110.25
3	A	403	XYP	O4A-C1B-C2B	-2.80	96.91	111.47

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	402	GCV	1	0
3	A	404	XYP	2	0
2	B	402	GCV	1	0
3	B	404	XYP	3	0
3	C	404	XYP	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	389/401 (97%)	0.41	25 (6%) 20 15	42, 58, 72, 80	0
1	B	390/401 (97%)	0.42	6 (1%) 74 69	42, 58, 73, 79	1 (0%)
1	C	389/401 (97%)	0.50	7 (1%) 69 63	42, 58, 72, 79	0
1	D	367/401 (91%)	2.75	213 (58%) 0 0	74, 120, 155, 179	364 (99%)
All	All	1535/1604 (95%)	1.00	251 (16%) 2 1	42, 62, 140, 179	365 (23%)

The worst 5 of 251 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	102	SER	10.2
1	D	343	LEU	9.7
1	D	103	ALA	9.3
1	D	100	ASP	9.0
1	D	319	TYR	7.9

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
3	XYP	C	404	9/10	0.44	0.38	9.60	118,118,118,119	0
3	XYP	C	403	9/10	0.46	0.43	5.44	115,116,117,118	0
3	XYP	A	404	9/10	0.89	0.31	2.82	92,94,94,95	0
3	XYP	B	404	9/10	0.82	0.21	2.01	81,82,83,83	0
2	GCV	C	402	14/14	0.82	0.28	1.40	112,114,115,116	0
2	GCV	A	402	14/14	0.88	0.28	0.87	91,94,95,95	0
2	GCV	B	402	14/14	0.94	0.16	0.06	73,76,78,79	0
2	GCV	B	405	14/14	0.86	0.17	-1.83	86,88,90,90	0
3	XYP	B	403	9/10	0.92	0.12	-2.08	77,78,78,80	0
3	XYP	A	403	9/10	0.80	0.29	-	92,93,94,95	0
3	XYP	B	407	8/10	0.91	0.19	-	91,92,92,93	0
3	XYP	B	406	9/10	0.78	0.17	-	88,90,91,91	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.