



wwPDB X-ray Structure Validation Summary Report ⓘ

Nov 15, 2017 – 06:34 AM EST

PDB ID : 4PC4
Title : Bombyx mori lipoprotein 6
Authors : Pietrzyk, A.J.; Bujacz, A.; Jaskolski, M.; Bujacz, G.
Deposited on : unknown
Resolution : 1.80 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix)	:	1.9-1692
EDS	:	rb-20030345
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	rb-20030345


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	4827 (1.80-1.80)
Clashscore	112137	5742 (1.80-1.80)
Ramachandran outliers	110173	5676 (1.80-1.80)
Sidechain outliers	110143	5675 (1.80-1.80)
RSRZ outliers	101464	4906 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	245	
1	B	245	
1	C	245	
1	D	245	
1	E	245	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	MES	A	302	-	-	X	X
2	MES	B	301	-	-	-	X
2	MES	C	301	-	-	-	X
3	EDO	B	302	-	-	-	X
3	EDO	C	302	-	-	X	-
4	PEG	B	304	-	-	-	X
5	IPA	C	304	-	-	-	X
5	IPA	C	306[A]	-	-	X	-

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 11107 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 30K lipoprotein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	238	Total	C	N	O	S	0	3	0
			1973	1254	341	372	6			
1	B	238	Total	C	N	O	S	0	3	0
			1972	1254	338	374	6			
1	C	240	Total	C	N	O	S	0	7	0
			2007	1279	346	376	6			
1	D	238	Total	C	N	O	S	0	2	0
			1966	1250	338	372	6			
1	E	239	Total	C	N	O	S	0	2	0
			1967	1249	338	374	6			

There are 5 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	217	ASN	TYR	conflict	UNP A7LIK7
B	217	ASN	TYR	conflict	UNP A7LIK7
C	217	ASN	TYR	conflict	UNP A7LIK7
D	217	ASN	TYR	conflict	UNP A7LIK7
E	217	ASN	TYR	conflict	UNP A7LIK7

- Molecule 2 is 2-(N-MORPHOLINO)-ETHANESULFONIC ACID (three-letter code: MES) (formula: C₆H₁₃NO₄S).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	S	0	0
			12	6	1	4	1		
2	A	1	Total	C	N	O	S	0	0
			12	6	1	4	1		
2	B	1	Total	C	N	O	S	0	0
			12	6	1	4	1		
2	C	1	Total	C	N	O	S	0	0
			12	6	1	4	1		

- Molecule 3 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			4	2	2		
3	B	1	Total	C	O	0	0
			4	2	2		
3	B	1	Total	C	O	0	0
			4	2	2		
3	C	1	Total	C	O	0	0
			4	2	2		
3	C	1	Total	C	O	0	0
			4	2	2		

- Molecule 4 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula: C₄H₁₀O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	C	O	0	0
			7	4	3		

- Molecule 5 is ISOPROPYL ALCOHOL (three-letter code: IPA) (formula: C₃H₈O).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	C	1	Total	C	O	0	0
			4	3	1		
5	C	1	Total	C	O	0	0
			4	3	1		
5	C	1	Total	C	O	0	1
			8	6	2		

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	268	Total	O	0	0
			268	268		
6	B	274	Total	O	0	0
			274	274		
6	C	302	Total	O	0	0
			302	302		
6	D	171	Total	O	0	0
			171	171		
6	E	116	Total	O	0	0
			116	116		

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: 30K lipoprotein

Chain A: 




• Molecule 1: 30K lipoprotein

Chain B: 





• Molecule 1: 30K lipoprotein

Chain C: 




• Molecule 1: 30K lipoprotein

Chain D: 



• Molecule 1: 30K lipoprotein

Chain E: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	83.21Å 85.75Å 104.47Å 90.00° 104.84° 90.00°	Depositor
Resolution (Å)	47.00 – 1.80 47.05 – 1.80	Depositor EDS
% Data completeness (in resolution range)	94.9 (47.00-1.80) 95.0 (47.05-1.80)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.48 (at 1.79Å)	Xtriage
Refinement program	REFMAC 5.7.0032	Depositor
R, R_{free}	0.161 , 0.183 0.171 , 0.197	Depositor DCC
R_{free} test set	1265 reflections (1.02%)	DCC
Wilson B-factor (Å ²)	26.7	Xtriage
Anisotropy	0.091	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 42.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	11107	wwPDB-VP
Average B, all atoms (Å ²)	36.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.50% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PEG, IPA, EDO, MES

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.69	0/2027	0.77	1/2743 (0.0%)
1	B	0.71	0/2026	0.78	1/2742 (0.0%)
1	C	0.70	0/2075	0.80	2/2809 (0.1%)
1	D	0.63	0/2017	0.80	2/2730 (0.1%)
1	E	0.57	0/2018	0.77	2/2734 (0.1%)
All	All	0.66	0/10163	0.78	8/13758 (0.1%)

There are no bond length outliers.

The worst 5 of 8 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	E	143	ARG	NE-CZ-NH2	-5.88	117.36	120.30
1	E	143	ARG	NE-CZ-NH1	5.50	123.05	120.30
1	D	143	ARG	NE-CZ-NH2	-5.49	117.56	120.30
1	A	134	LYS	CD-CE-NZ	5.19	123.64	111.70
1	C	241[A]	PHE	CB-CA-C	-5.12	100.15	110.40

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1973	0	1913	29	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	1972	0	1908	22	0
1	C	2007	0	1951	24	0
1	D	1966	0	1902	20	0
1	E	1967	0	1895	14	0
2	A	24	0	26	15	0
2	B	12	0	13	0	0
2	C	12	0	13	0	0
3	A	4	0	6	0	0
3	B	8	0	12	3	0
3	C	8	0	12	6	0
4	B	7	0	10	3	0
5	C	16	0	32	8	0
6	A	268	0	0	5	0
6	B	274	0	0	3	0
6	C	302	0	0	10	0
6	D	171	0	0	7	0
6	E	116	0	0	3	0
All	All	11107	0	9693	113	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 113 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:302:MES:H22	1:B:142:ASN:HD22	1.09	1.13
1:A:15:GLU:HB2	2:A:302:MES:O2S	1.59	1.03
2:A:302:MES:H22	1:B:142:ASN:ND2	1.87	0.88
1:A:15:GLU:CB	2:A:302:MES:O2S	2.22	0.88
1:A:12:GLN:HG2	2:A:302:MES:H82	1.59	0.85

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	239/245 (98%)	236 (99%)	3 (1%)	0	100	100
1	B	239/245 (98%)	236 (99%)	3 (1%)	0	100	100
1	C	244/245 (100%)	240 (98%)	4 (2%)	0	100	100
1	D	238/245 (97%)	235 (99%)	3 (1%)	0	100	100
1	E	239/245 (98%)	235 (98%)	4 (2%)	0	100	100
All	All	1199/1225 (98%)	1182 (99%)	17 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	215/217 (99%)	214 (100%)	1 (0%)	91	90
1	B	215/217 (99%)	215 (100%)	0	100	100
1	C	220/217 (101%)	218 (99%)	2 (1%)	82	78
1	D	214/217 (99%)	210 (98%)	4 (2%)	62	50
1	E	214/217 (99%)	207 (97%)	7 (3%)	43	26
All	All	1078/1085 (99%)	1064 (99%)	14 (1%)	75	66

5 of 14 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	234	LEU
1	E	48[A]	ASN
1	E	207	ASN
1	D	230	GLU
1	E	167	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 33 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	141	ASN
1	C	227	HIS
1	E	177	ASN
1	C	149	HIS
1	C	154	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

14 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	MES	A	301	-	12,12,12	1.91	1 (8%)	14,16,16	1.74	4 (28%)
2	MES	A	302	-	12,12,12	2.10	1 (8%)	14,16,16	6.13	8 (57%)
3	EDO	A	303	-	3,3,3	0.43	0	2,2,2	0.39	0
2	MES	B	301	-	12,12,12	2.00	1 (8%)	14,16,16	2.12	6 (42%)
3	EDO	B	302	-	3,3,3	0.29	0	2,2,2	0.46	0
3	EDO	B	303	-	3,3,3	0.39	0	2,2,2	0.42	0
4	PEG	B	304	-	6,6,6	0.49	0	5,5,5	0.38	0
2	MES	C	301	-	12,12,12	1.82	1 (8%)	14,16,16	2.02	2 (14%)
3	EDO	C	302	-	3,3,3	0.60	0	2,2,2	0.08	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	EDO	C	303	-	3,3,3	0.52	0	2,2,2	0.07	0
5	IPA	C	304	-	3,3,3	0.37	0	3,3,3	0.42	0
5	IPA	C	305	-	3,3,3	0.66	0	3,3,3	0.39	0
5	IPA	C	306[A]	-	3,3,3	0.44	0	3,3,3	0.72	0
5	IPA	C	306[B]	-	3,3,3	0.38	0	3,3,3	0.39	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	MES	A	301	-	-	0/6/14/14	0/1/1/1
2	MES	A	302	-	-	0/6/14/14	0/1/1/1
3	EDO	A	303	-	-	0/1/1/1	0/0/0/0
2	MES	B	301	-	-	0/6/14/14	0/1/1/1
3	EDO	B	302	-	-	0/1/1/1	0/0/0/0
3	EDO	B	303	-	-	0/1/1/1	0/0/0/0
4	PEG	B	304	-	-	0/4/4/4	0/0/0/0
2	MES	C	301	-	-	0/6/14/14	0/1/1/1
3	EDO	C	302	-	-	0/1/1/1	0/0/0/0
3	EDO	C	303	-	-	0/1/1/1	0/0/0/0
5	IPA	C	304	-	-	0/0/0/0	0/0/0/0
5	IPA	C	305	-	-	0/0/0/0	0/0/0/0
5	IPA	C	306[A]	-	-	0/0/0/0	0/0/0/0
5	IPA	C	306[B]	-	-	0/0/0/0	0/0/0/0

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	302	MES	C8-S	-6.88	1.67	1.77
2	A	301	MES	C8-S	-5.96	1.68	1.77
2	B	301	MES	C8-S	-5.76	1.68	1.77
2	C	301	MES	C8-S	-5.69	1.69	1.77

The worst 5 of 20 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	302	MES	O1S-S-C8	-10.75	97.55	106.79
2	A	302	MES	O2S-S-C8	-10.16	98.07	106.79
2	A	302	MES	C6-C5-N4	-4.27	104.13	110.11
2	A	302	MES	C2-C3-N4	-4.24	104.16	110.11

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	302	MES	O3S-S-O2S	-3.98	102.25	111.37

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

8 monomers are involved in 30 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	302	MES	15	0
3	B	303	EDO	3	0
4	B	304	PEG	3	0
3	C	302	EDO	5	0
3	C	303	EDO	1	0
5	C	304	IPA	1	0
5	C	306[A]	IPA	5	0
5	C	306[B]	IPA	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	238/245 (97%)	-0.73	1 (0%) 92 90	18, 28, 49, 89	0
1	B	238/245 (97%)	-0.63	0 100 100	17, 26, 43, 69	0
1	C	240/245 (97%)	-0.57	1 (0%) 92 90	19, 27, 44, 81	0
1	D	238/245 (97%)	-0.34	1 (0%) 92 90	23, 40, 61, 91	0
1	E	239/245 (97%)	0.50	28 (11%) 5 4	22, 51, 88, 106	0
All	All	1193/1225 (97%)	-0.36	31 (2%) 56 52	17, 32, 66, 106	0

The worst 5 of 31 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	7	ALA	7.0
1	E	7	ALA	6.3
1	E	193	TYR	4.3
1	E	83	LEU	3.9
1	E	105	LEU	3.8

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron

density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
4	PEG	B	304	7/7	0.82	0.34	29.09	48,50,54,59	0
2	MES	A	302	12/12	0.67	0.23	6.99	44,50,65,67	12
2	MES	B	301	12/12	0.91	0.14	5.19	35,43,45,48	12
5	IPA	C	304	4/4	0.92	0.18	4.77	52,53,55,57	0
3	EDO	B	302	4/4	0.96	0.17	4.76	34,35,41,43	0
2	MES	C	301	12/12	0.84	0.17	3.97	51,58,62,69	12
2	MES	A	301	12/12	0.93	0.13	-	49,55,68,71	0
3	EDO	B	303	4/4	0.93	0.26	-	37,43,45,47	0
3	EDO	C	303	4/4	0.78	0.21	-	67,70,70,71	0
5	IPA	C	306[A]	4/4	0.96	0.19	-	19,20,20,23	4
3	EDO	C	302	4/4	0.73	0.17	-	64,72,74,75	0
5	IPA	C	305	4/4	0.84	0.15	-	38,42,48,50	0
5	IPA	C	306[B]	4/4	0.96	0.19	-	29,31,31,32	4
3	EDO	A	303	4/4	0.95	0.12	-	37,52,57,64	0

6.5 Other polymers [i](#)

There are no such residues in this entry.