



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 12, 2017 – 07:56 pm GMT

PDB ID : 2A1I
Title : Crystal Structure of the Central Domain of Human ERCC1
Authors : Tsodikov, O.V.; Enzlin, J.H.; Scharer, O.D.; Ellenberger, T.
Deposited on : 2005-06-20
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.9-1692
EDS	:	trunk28620
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	5047 (1.90-1.90)
Clashscore	112137	5731 (1.90-1.90)
Ramachandran outliers	110173	5669 (1.90-1.90)
Sidechain outliers	110143	5670 (1.90-1.90)
RSRZ outliers	101464	5100 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	146	

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 1170 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA excision repair protein ERCC-1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	129	Total	C	N	O	S	0	0	0
			1039	672	178	184	5			

There are 14 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	82	MET	-	CLONING ARTIFACT	UNP P07992
A	83	GLY	-	CLONING ARTIFACT	UNP P07992
A	84	SER	-	CLONING ARTIFACT	UNP P07992
A	85	SER	-	CLONING ARTIFACT	UNP P07992
A	86	HIS	-	EXPRESSION TAG	UNP P07992
A	87	HIS	-	EXPRESSION TAG	UNP P07992
A	88	HIS	-	EXPRESSION TAG	UNP P07992
A	89	HIS	-	EXPRESSION TAG	UNP P07992
A	90	HIS	-	EXPRESSION TAG	UNP P07992
A	91	HIS	-	EXPRESSION TAG	UNP P07992
A	92	SER	-	CLONING ARTIFACT	UNP P07992
A	93	GLN	-	CLONING ARTIFACT	UNP P07992
A	94	ASP	-	CLONING ARTIFACT	UNP P07992
A	95	PRO	-	CLONING ARTIFACT	UNP P07992

- Molecule 2 is MERCURY (II) ION (three-letter code: HG) (formula: Hg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Hg	0	0
			1	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	130	Total	O	0	0
			130	130		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: DNA excision repair protein ERCC-1



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	40.07Å 50.29Å 72.22Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.00 – 1.90 29.33 – 1.90	Depositor EDS
% Data completeness (in resolution range)	99.6 (30.00-1.90) 99.6 (29.33-1.90)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.93 (at 1.91Å)	Xtriage
Refinement program	REFMAC 5.1.24	Depositor
R, R_{free}	0.204 , 0.248 0.248 , 0.281	Depositor DCC
R_{free} test set	586 reflections (4.93%)	DCC
Wilson B-factor (Å ²)	35.5	Xtriage
Anisotropy	0.542	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 55.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	1170	wwPDB-VP
Average B, all atoms (Å ²)	33.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 11.06% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: HG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.83	3/1063 (0.3%)	0.92	2/1443 (0.1%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	220	ALA	C-O	9.49	1.41	1.23
1	A	215	TYR	CG-CD1	6.73	1.48	1.39
1	A	215	TYR	CE2-CZ	6.53	1.47	1.38

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	125	ASP	CB-CG-OD2	5.29	123.06	118.30
1	A	169	LEU	CB-CG-CD2	5.20	119.84	111.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1039	0	1059	31	0
2	A	1	0	0	0	0
3	A	130	0	0	19	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	1170	0	1059	31	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 15.

All (31) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:147:ASN:HB2	3:A:327:HOH:O	1.56	1.05
1:A:181:ALA:O	1:A:184:GLU:HG2	1.78	0.84
1:A:99:ASN:N	3:A:348:HOH:O	2.19	0.74
1:A:216:GLU:HG2	3:A:343:HOH:O	1.88	0.71
1:A:225:GLU:HB2	3:A:334:HOH:O	1.91	0.70
1:A:198:LEU:HG	3:A:357:HOH:O	1.91	0.69
1:A:216:GLU:OE2	1:A:219:PRO:HA	1.92	0.69
1:A:128:PRO:HG3	1:A:164:PHE:CE1	2.29	0.67
1:A:218:LYS:O	3:A:351:HOH:O	2.13	0.66
1:A:204:GLU:OE2	3:A:330:HOH:O	2.14	0.65
1:A:188:MET:HE2	3:A:313:HOH:O	2.00	0.61
1:A:220:ALA:O	3:A:233:HOH:O	2.17	0.60
1:A:154:HIS:HB2	3:A:313:HOH:O	2.03	0.59
1:A:162:LYS:NZ	3:A:320:HOH:O	2.35	0.58
1:A:183:LYS:HD3	3:A:309:HOH:O	2.05	0.55
1:A:154:HIS:O	1:A:158:GLN:HG2	2.07	0.55
1:A:118:ASN:HB2	3:A:317:HOH:O	2.05	0.55
1:A:167:ARG:HD2	3:A:266:HOH:O	2.08	0.53
1:A:216:GLU:O	1:A:219:PRO:HD3	2.08	0.53
1:A:176:LYS:HE2	3:A:303:HOH:O	2.10	0.51
1:A:179:GLN:H	1:A:179:GLN:CD	2.18	0.47
1:A:136:THR:HG23	1:A:166:LEU:HB3	1.96	0.47
1:A:171:VAL:HB	3:A:357:HOH:O	2.14	0.47
1:A:196:LEU:CD2	1:A:198:LEU:HD12	2.45	0.47
1:A:207:ARG:HH11	1:A:207:ARG:CG	2.32	0.42
1:A:222:LEU:O	1:A:225:GLU:HB3	2.20	0.42
1:A:188:MET:CE	3:A:313:HOH:O	2.63	0.41
1:A:103:VAL:O	1:A:123:PHE:HA	2.20	0.41
1:A:149:HIS:HD2	3:A:323:HOH:O	2.04	0.41
1:A:137:CYS:SG	1:A:164:PHE:CD1	3.14	0.41
1:A:99:ASN:HB3	3:A:338:HOH:O	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	127/146 (87%)	124 (98%)	3 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	114/129 (88%)	99 (87%)	15 (13%)	5	1

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	132	LEU
1	A	135	SER
1	A	139	LEU
1	A	142	SER
1	A	148	LEU
1	A	166	LEU
1	A	169	LEU
1	A	175	VAL
1	A	176	LYS
1	A	180	GLN
1	A	207	ARG
1	A	209	LEU
1	A	218	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	224	MET
1	A	227	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	179	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	129/146 (88%)	1.07	27 (20%) 1 1	19, 27, 44, 58	0

All (27) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	163	ASN	5.5
1	A	134	GLN	4.4
1	A	162	LYS	3.6
1	A	99	ASN	3.4
1	A	220	ALA	3.3
1	A	161	GLY	3.3
1	A	197	ILE	3.2
1	A	133	GLY	3.1
1	A	127	ILE	3.1
1	A	218	LYS	3.1
1	A	165	ALA	3.0
1	A	135	SER	3.0
1	A	224	MET	2.8
1	A	209	LEU	2.8
1	A	140	PHE	2.7
1	A	168	VAL	2.6
1	A	141	LEU	2.5
1	A	208	TYR	2.5
1	A	171	VAL	2.4
1	A	158	GLN	2.4
1	A	170	LEU	2.4
1	A	176	LYS	2.4
1	A	130	TYR	2.3
1	A	164	PHE	2.3
1	A	196	LEU	2.3
1	A	151	ASP	2.1
1	A	169	LEU	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	HG	A	1	1/1	0.98	0.16	0.43	76,76,76,76	1

6.5 Other polymers [i](#)

There are no such residues in this entry.