



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 13, 2017 – 03:59 pm GMT

PDB ID : 3A1D
Title : Crystal structure of the P- and N-domains of CopA, a copper-transporting P-type ATPase, bound with ADP-Mg
Authors : Tsuda, T.; Toyoshima, C.
Deposited on : 2009-03-31
Resolution : 1.85 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : trunk28620
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	1923 (1.86-1.86)
Clashscore	112137	2083 (1.86-1.86)
Ramachandran outliers	110173	2060 (1.86-1.86)
Sidechain outliers	110143	2060 (1.86-1.86)
RSRZ outliers	101464	1932 (1.86-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	287	<div> <div style="width: 100%; height: 10px; position: relative;"> <div style="position: absolute; top: -10px; left: 0;">%</div> <div style="position: absolute; top: 0; left: 0; width: 84%; height: 10px; background-color: green;"></div> <div style="position: absolute; top: 0; left: 84%; width: 9%; height: 10px; background-color: yellow;"></div> <div style="position: absolute; top: 0; left: 93%; width: 7%; height: 10px; background-color: grey;"></div> </div> <div>84% 9% 7%</div> </div>
1	B	287	<div> <div style="width: 100%; height: 10px; position: relative;"> <div style="position: absolute; top: 0; left: 0; width: 8%; height: 10px; background-color: red;"></div> <div style="position: absolute; top: 0; left: 8%; width: 84%; height: 10px; background-color: green;"></div> <div style="position: absolute; top: 0; left: 92%; width: 10%; height: 10px; background-color: yellow;"></div> <div style="position: absolute; top: 0; left: 99%; width: 5%; height: 10px; background-color: grey;"></div> </div> <div>8% 84% 10% 5%</div> </div>

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 4644 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Probable copper-exporting P-type ATPase A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	268	Total	C	N	O	S	0	0	0
			2004	1255	350	396	3			
1	B	272	Total	C	N	O	S	0	0	0
			2029	1269	354	402	4			

There are 22 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	387	MET	-	EXPRESSION TAG	UNP O29777
A	388	GLY	-	EXPRESSION TAG	UNP O29777
A	389	HIS	-	EXPRESSION TAG	UNP O29777
A	390	HIS	-	EXPRESSION TAG	UNP O29777
A	391	HIS	-	EXPRESSION TAG	UNP O29777
A	392	HIS	-	EXPRESSION TAG	UNP O29777
A	393	HIS	-	EXPRESSION TAG	UNP O29777
A	394	HIS	-	EXPRESSION TAG	UNP O29777
A	395	GLY	-	EXPRESSION TAG	UNP O29777
A	396	SER	-	EXPRESSION TAG	UNP O29777
A	397	ARG	-	EXPRESSION TAG	UNP O29777
B	387	MET	-	EXPRESSION TAG	UNP O29777
B	388	GLY	-	EXPRESSION TAG	UNP O29777
B	389	HIS	-	EXPRESSION TAG	UNP O29777
B	390	HIS	-	EXPRESSION TAG	UNP O29777
B	391	HIS	-	EXPRESSION TAG	UNP O29777
B	392	HIS	-	EXPRESSION TAG	UNP O29777
B	393	HIS	-	EXPRESSION TAG	UNP O29777
B	394	HIS	-	EXPRESSION TAG	UNP O29777
B	395	GLY	-	EXPRESSION TAG	UNP O29777
B	396	SER	-	EXPRESSION TAG	UNP O29777
B	397	ARG	-	EXPRESSION TAG	UNP O29777

- Molecule 2 is ADENOSINE-5'-DIPHOSPHATE (three-letter code: ADP) (formula: $C_{10}H_{15}N_5O_{10}P_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			27	10	5	10	2		
2	B	1	Total	C	N	O	P	0	0
			27	10	5	10	2		

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total	Mg	0	0
			1	1		
3	A	1	Total	Mg	0	0
			1	1		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	291	Total	O	0	0
			291	291		
4	B	264	Total	O	0	0
			264	264		

- Molecule 1: Probable copper-exporting P-type ATPase A



4 Data and refinement statistics

Property	Value	Source
Space group	P 43 2 2	Depositor
Cell constants a, b, c, α , β , γ	89.96Å 89.96Å 190.38Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 1.85 81.33 – 1.85	Depositor EDS
% Data completeness (in resolution range)	99.8 (20.00-1.85) 99.7 (81.33-1.85)	Depositor EDS
R_{merge}	0.13	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.41 (at 1.84Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.192 , 0.226 0.192 , 0.224	Depositor DCC
R_{free} test set	3415 reflections (5.34%)	DCC
Wilson B-factor (Å ²)	35.2	Xtriage
Anisotropy	0.045	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 53.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	4644	wwPDB-VP
Average B, all atoms (Å ²)	41.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.27% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MG, ADP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.54	0/2018	0.65	2/2725 (0.1%)
1	B	0.59	2/2043 (0.1%)	0.66	3/2758 (0.1%)
All	All	0.57	2/4061 (0.0%)	0.65	5/5483 (0.1%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	447	ARG	CZ-NH1	15.36	1.53	1.33
1	B	447	ARG	NE-CZ	6.76	1.41	1.33

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	447	ARG	NE-CZ-NH1	14.25	127.42	120.30
1	B	447	ARG	NE-CZ-NH2	-9.94	115.33	120.30
1	A	526	ARG	NE-CZ-NH1	6.40	123.50	120.30
1	A	526	ARG	NE-CZ-NH2	-5.72	117.44	120.30
1	B	654	ASP	CB-CG-OD1	5.17	122.96	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2004	0	2091	21	0
1	B	2029	0	2115	20	0
2	A	27	0	12	0	0
2	B	27	0	12	1	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	291	0	0	8	0
4	B	264	0	0	4	0
All	All	4644	0	4230	40	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (40) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:426:THR:O	4:A:1106:HOH:O	1.87	0.91
1:A:618:ASP:O	1:A:622:ASP:OD1	1.95	0.83
1:B:424:ASP:CG	4:B:1440:HOH:O	2.25	0.73
1:A:424:ASP:CG	4:A:1141:HOH:O	2.25	0.73
1:B:450:LEU:HD11	1:B:471:LYS:HG3	1.72	0.72
1:A:602:GLU:HG2	4:A:1227:HOH:O	1.88	0.72
1:A:583:SER:HB2	1:A:591:VAL:HG21	1.79	0.64
1:B:619:GLY:HA2	1:B:626:LEU:CD1	2.30	0.62
1:B:605:LYS:HA	1:B:608:GLN:HE21	1.64	0.61
1:A:560:GLU:HG2	1:A:563:ARG:HH22	1.65	0.60
1:A:605:LYS:HA	1:A:608:GLN:HE21	1.66	0.60
1:A:407:LYS:HD3	4:A:1075:HOH:O	2.03	0.58
1:B:424:ASP:OD2	4:B:1440:HOH:O	2.16	0.58
1:B:422:ILE:CD1	1:B:613:VAL:HG13	2.35	0.57
1:B:422:ILE:HD13	1:B:613:VAL:HG13	1.89	0.55
1:B:619:GLY:HA2	1:B:626:LEU:HD11	1.90	0.54
1:A:483:GLU:HG2	1:A:484:LYS:HG3	1.90	0.53
1:A:580:GLU:HG3	1:B:576:TRP:CZ3	2.44	0.52
1:A:556:PRO:O	1:A:560:GLU:HG3	2.10	0.51
1:A:428:THR:N	4:A:1085:HOH:O	2.42	0.51
1:A:618:ASP:OD2	1:A:618:ASP:O	2.29	0.50
1:A:504:ARG:NH2	4:A:1231:HOH:O	2.42	0.49
1:A:618:ASP:OD2	1:A:618:ASP:C	2.53	0.47
1:B:452:LEU:HD21	1:B:536:ARG:HH11	1.79	0.47
1:B:620:ILE:CD1	1:B:643:VAL:HG23	2.46	0.45
1:B:422:ILE:HD13	1:B:613:VAL:CG1	2.47	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:614:ALA:HA	1:A:631:LEU:O	2.17	0.44
1:B:503:LYS:HD2	1:B:518:GLU:OE1	2.18	0.43
1:B:620:ILE:HD11	1:B:643:VAL:HG23	1.99	0.43
1:A:424:ASP:O	1:A:428:THR:HB	2.19	0.43
1:B:517:VAL:HG13	1:B:543:ILE:HD11	2.01	0.43
1:B:457:GLU:OE1	2:B:997:ADP:N1	2.51	0.43
1:A:575:ASN:HB3	4:A:1230:HOH:O	2.20	0.41
1:B:614:ALA:HA	1:B:631:LEU:O	2.20	0.41
1:B:511:VAL:HG22	1:B:538:GLY:HA2	2.02	0.41
1:B:663:GLN:NE2	4:B:1349:HOH:O	2.52	0.41
1:A:539:ARG:HG3	4:A:1282:HOH:O	2.22	0.40
1:B:635:VAL:HB	4:B:1434:HOH:O	2.21	0.40
1:A:421:VAL:HG23	1:A:566:ILE:HG21	2.04	0.40
1:A:618:ASP:C	1:A:620:ILE:H	2.25	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	266/287 (93%)	263 (99%)	3 (1%)	0	100	100
1	B	270/287 (94%)	266 (98%)	3 (1%)	1 (0%)	38	22
All	All	536/574 (93%)	529 (99%)	6 (1%)	1 (0%)	51	35

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	496	ASP

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	212/228 (93%)	210 (99%)	2 (1%)	82	77
1	B	215/228 (94%)	212 (99%)	3 (1%)	71	61
All	All	427/456 (94%)	422 (99%)	5 (1%)	75	67

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	428	THR
1	A	667	LYS
1	B	450	LEU
1	B	459	ARG
1	B	499	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	559	GLN
1	A	587	ASN
1	A	599	GLN
1	A	608	GLN
1	A	628	GLN
1	B	537	ASN
1	B	608	GLN
1	B	621	ASN
1	B	628	GLN
1	B	663	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 2 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	ADP	A	997	3	25,29,29	1.06	1 (4%)	24,45,45	2.06	3 (12%)
2	ADP	B	997	3	25,29,29	1.06	1 (4%)	24,45,45	1.81	3 (12%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	ADP	A	997	3	-	0/12/32/32	0/3/3/3
2	ADP	B	997	3	-	0/12/32/32	0/3/3/3

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	997	ADP	C5-C4	3.08	1.47	1.40
2	B	997	ADP	C5-C4	3.30	1.47	1.40

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	997	ADP	N3-C2-N1	-8.64	121.33	128.86
2	B	997	ADP	N3-C2-N1	-7.30	122.50	128.86
2	B	997	ADP	C4-C5-N7	-2.46	107.03	109.41
2	A	997	ADP	C4-C5-N7	-2.07	107.41	109.41
2	B	997	ADP	C2-N1-C6	2.30	122.79	118.77
2	A	997	ADP	C2-N1-C6	2.63	123.36	118.77

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	997	ADP	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	268/287 (93%)	0.52	2 (0%) 87 88	24, 34, 44, 57	0
1	B	272/287 (94%)	0.71	24 (8%) 11 10	27, 40, 62, 71	0
All	All	540/574 (94%)	0.62	26 (4%) 31 30	24, 35, 60, 71	0

All (26) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	481	GLU	5.3
1	B	477	ILE	5.1
1	B	399	GLY	5.0
1	B	459	ARG	4.3
1	B	455	ILE	4.2
1	B	480	GLY	4.0
1	B	442	LEU	3.7
1	B	479	LEU	3.7
1	B	443	ASN	3.6
1	A	667	LYS	3.5
1	B	485	VAL	3.2
1	B	496	ASP	3.2
1	B	494	VAL	3.2
1	B	441	PRO	3.0
1	B	482	PRO	2.8
1	B	478	GLU	2.8
1	B	483	GLU	2.7
1	B	440	VAL	2.7
1	B	495	ALA	2.6
1	B	476	GLY	2.6
1	B	473	LEU	2.5
1	B	474	GLU	2.3
1	B	487	VAL	2.2
1	A	400	ALA	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	422	ILE	2.1
1	B	458	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	ADP	B	997	27/27	0.95	0.14	0.58	41,48,49,50	0
2	ADP	A	997	27/27	0.97	0.08	-3.61	29,32,36,39	0
3	MG	A	998	1/1	0.99	0.04	-	25,25,25,25	0
3	MG	B	998	1/1	0.89	0.06	-	39,39,39,39	0

6.5 Other polymers [i](#)

There are no such residues in this entry.