



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 13, 2017 – 09:37 pm GMT

PDB ID : 5A1K
Title : Crystal structure of calcium-free human adseverin domains A1-A3
Authors : Chumnarnsilpa, S.; Robinson, R.C.; Grimes, J.M.; Leyrat, C.
Deposited on : 2015-04-30
Resolution : 2.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.9-1692
EDS	:	trunk28620
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	1586 (2.90-2.90)
Clashscore	112137	1807 (2.90-2.90)
Ramachandran outliers	110173	1768 (2.90-2.90)
Sidechain outliers	110143	1770 (2.90-2.90)
RSRZ outliers	101464	1596 (2.90-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	344	<div> <div>3%</div> <div> <div></div> <div>72%</div> <div>24%</div> <div>• •</div> </div> </div>
1	B	344	<div> <div>4%</div> <div> <div></div> <div>72%</div> <div>22%</div> <div>• •</div> </div> </div>

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 5521 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ADSEVERIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	341	Total	C	N	O	S	0	0	0
			2713	1718	467	515	13			
1	B	340	Total	C	N	O	S	0	0	0
			2701	1709	466	513	13			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	61	ARG	HIS	VARIANT	UNP Q9Y6U3
B	61	ARG	HIS	VARIANT	UNP Q9Y6U3


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	63	Total	O	0	0
			63	63		
2	B	44	Total	O	0	0
			44	44		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($\text{RSRZ} > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: ADSEVERIN



• Molecule 1: ADSEVERIN



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	135.41Å 65.25Å 105.19Å 90.00° 118.27° 90.00°	Depositor
Resolution (Å)	59.63 – 2.90 59.63 – 2.90	Depositor EDS
% Data completeness (in resolution range)	97.8 (59.63-2.90) 97.6 (59.63-2.90)	Depositor EDS
R_{merge}	0.15	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.58 (at 2.91Å)	Xtriage
Refinement program	BUSTER 2.10.2	Depositor
R, R_{free}	0.240 , 0.248 0.251 , 0.260	Depositor DCC
R_{free} test set	904 reflections (5.37%)	DCC
Wilson B-factor (Å ²)	57.0	Xtriage
Anisotropy	0.140	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 27.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.90	EDS
Total number of atoms	5521	wwPDB-VP
Average B, all atoms (Å ²)	55.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 9.92% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.39	0/2770	0.69	0/3733
1	B	0.38	0/2757	0.69	0/3715
All	All	0.38	0/5527	0.69	0/7448

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2713	0	2655	41	1
1	B	2701	0	2646	60	1
2	A	63	0	0	0	0
2	B	44	0	0	0	0
All	All	5521	0	5301	99	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (99) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:250:LYS:HZ1	1:B:268:GLU:HA	0.98	1.09

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:130:LEU:H	1:A:130:LEU:HD12	1.30	0.96
1:B:250:LYS:NZ	1:B:268:GLU:HA	1.80	0.96
1:B:250:LYS:HZ1	1:B:268:GLU:CA	1.78	0.95
1:A:146:ARG:HH11	1:A:146:ARG:HA	1.32	0.94
1:B:134:LEU:HD13	1:B:134:LEU:H	1.34	0.93
1:B:205:ARG:HB2	1:B:205:ARG:HH11	1.34	0.93
1:B:134:LEU:CD2	1:B:135:THR:HG23	2.05	0.87
1:B:135:THR:HA	1:B:171:GLY:HA3	1.54	0.86
1:B:146:ARG:HH11	1:B:146:ARG:HG3	1.42	0.85
1:B:134:LEU:HD23	1:B:135:THR:CG2	2.08	0.83
1:B:256:ASP:HB3	1:B:261:MET:HA	1.60	0.83
1:B:12:ARG:HH11	1:B:12:ARG:HG2	1.43	0.81
1:B:134:LEU:HD23	1:B:135:THR:HG23	1.68	0.74
1:A:127:ASN:HB3	1:A:129:VAL:HG22	1.71	0.71
1:B:127:ASN:HB3	1:B:129:VAL:HG22	1.71	0.70
1:B:135:THR:HA	1:B:171:GLY:CA	2.21	0.70
1:B:7:HIS:HA	1:B:10:PHE:CD1	2.28	0.69
1:A:97:ARG:HB2	1:A:310:MET:HE1	1.75	0.69
1:B:205:ARG:CB	1:B:205:ARG:HH11	2.05	0.68
1:B:134:LEU:O	1:B:171:GLY:HA2	1.94	0.68
1:B:134:LEU:HD23	1:B:135:THR:HG22	1.75	0.66
1:B:97:ARG:HB2	1:B:310:MET:HE1	1.76	0.66
1:A:130:LEU:HD12	1:A:130:LEU:N	2.07	0.65
1:A:227:LYS:HD2	1:A:230:LEU:HD11	1.80	0.64
1:B:7:HIS:HA	1:B:10:PHE:HD1	1.62	0.63
1:B:14:GLY:H	1:B:20:GLN:HE22	1.45	0.63
1:A:14:GLY:H	1:A:20:GLN:HE22	1.44	0.63
1:A:56:ARG:CA	1:A:56:ARG:HE	2.11	0.63
1:B:237:ASP:O	1:B:240:ILE:HD11	1.98	0.62
1:B:134:LEU:HD22	1:B:135:THR:H	1.65	0.61
1:B:130:LEU:H	1:B:130:LEU:HD12	1.65	0.60
1:B:134:LEU:CD1	1:B:134:LEU:H	2.08	0.60
1:B:146:ARG:HH11	1:B:146:ARG:CG	2.13	0.60
1:A:97:ARG:HB2	1:A:310:MET:CE	2.32	0.60
1:B:61:ARG:HH12	1:B:96:ASN:ND2	2.01	0.59
1:A:56:ARG:HE	1:A:56:ARG:N	2.00	0.59
1:A:61:ARG:HH12	1:A:96:ASN:ND2	2.00	0.59
1:B:97:ARG:HB2	1:B:310:MET:CE	2.33	0.58
1:B:60:TYR:HB2	1:B:93:PRO:HB3	1.86	0.57
1:A:42:VAL:HG13	1:A:69:GLU:HG3	1.88	0.55
1:B:12:ARG:NH1	1:B:12:ARG:HG2	2.17	0.55

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:196:GLY:O	1:B:200:ASN:HB2	2.07	0.54
1:B:143:LYS:HE2	1:B:236:ASP:HB2	1.90	0.54
1:A:183:ASN:ND2	1:A:185:TYR:HD2	2.06	0.54
1:B:205:ARG:HB2	1:B:205:ARG:NH1	2.13	0.53
1:B:137:LYS:HG3	1:B:137:LYS:O	2.09	0.53
1:B:252:TYR:HB3	1:B:263:VAL:HG13	1.91	0.52
1:B:24:ILE:HD11	1:B:47:LEU:HD13	1.91	0.52
1:A:24:ILE:HD11	1:A:47:LEU:HD13	1.91	0.52
1:A:56:ARG:N	1:A:56:ARG:NE	2.58	0.52
1:B:183:ASN:O	1:B:187:ARG:HG3	2.11	0.51
1:A:108:VAL:HG11	1:A:147:VAL:CG1	2.40	0.51
1:A:60:TYR:CE2	1:A:89:LEU:HD22	2.46	0.51
1:A:196:GLY:O	1:A:200:ASN:HB2	2.11	0.50
1:A:183:ASN:O	1:A:187:ARG:HG3	2.11	0.50
1:A:238:ASP:O	1:A:240:ILE:HD13	2.12	0.50
1:B:247:LYS:HB2	1:B:269:GLU:HG2	1.95	0.49
1:A:167:ILE:HD11	1:A:190:ALA:O	2.13	0.49
1:B:205:ARG:CG	1:B:205:ARG:HH11	2.26	0.48
1:A:6:TYR:CZ	1:A:7:HIS:CE1	3.01	0.48
1:B:42:VAL:HG13	1:B:69:GLU:HG3	1.95	0.48
1:A:108:VAL:HG11	1:A:147:VAL:HG11	1.94	0.48
1:A:60:TYR:HB2	1:A:93:PRO:HB3	1.95	0.48
1:B:167:ILE:HD11	1:B:190:ALA:O	2.15	0.47
1:B:250:LYS:NZ	1:B:268:GLU:C	2.68	0.47
1:A:239:ILE:HG21	1:B:319:GLN:HA	1.97	0.46
1:A:92:LYS:N	1:A:93:PRO:HD2	2.31	0.46
1:B:137:LYS:HA	1:B:169:ASP:O	2.16	0.46
1:A:239:ILE:HG23	1:B:319:GLN:NE2	2.31	0.46
1:B:92:LYS:N	1:B:93:PRO:HD2	2.31	0.45
1:A:247:LYS:HB2	1:A:269:GLU:HG2	1.98	0.45
1:B:134:LEU:O	1:B:171:GLY:CA	2.64	0.45
1:B:145:ARG:HG3	1:B:185:TYR:HE2	1.81	0.45
1:B:163:GLY:HA2	1:B:181:SER:OG	2.17	0.45
1:A:54:THR:HG23	1:A:57:GLY:C	2.37	0.45
1:B:40:PHE:HE2	1:B:115:LEU:HD22	1.82	0.45
1:A:40:PHE:HE2	1:A:115:LEU:HD22	1.82	0.45
1:A:146:ARG:HH11	1:A:146:ARG:CA	2.17	0.45
1:B:227:LYS:HD3	1:B:230:LEU:HD11	1.99	0.45
1:B:23:ARG:HD3	1:B:25:GLU:HG2	1.99	0.44
1:A:23:ARG:HD3	1:A:25:GLU:HG2	2.00	0.44
1:B:99:LEU:HD23	1:B:325:ASN:HD22	1.83	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:138:ARG:HG3	1:A:169:ASP:HB3	2.01	0.43
1:B:295:VAL:HG21	1:B:313:ALA:HB2	2.00	0.43
1:B:26:LYS:HD3	1:B:26:LYS:HA	1.84	0.43
1:A:262:ARG:HA	1:A:262:ARG:HD2	1.48	0.42
1:A:99:LEU:HD23	1:A:325:ASN:HD22	1.83	0.42
1:B:134:LEU:CD2	1:B:135:THR:H	2.32	0.42
1:A:295:VAL:HG21	1:A:313:ALA:HB2	2.01	0.42
1:B:145:ARG:HG3	1:B:185:TYR:CE2	2.55	0.42
1:A:11:ALA:O	1:A:15:LYS:HE3	2.19	0.42
1:A:56:ARG:CA	1:A:56:ARG:NE	2.83	0.41
1:B:250:LYS:HZ1	1:B:268:GLU:C	2.17	0.41
1:B:250:LYS:NZ	1:B:268:GLU:CA	2.58	0.41
1:B:169:ASP:OD1	1:B:174:ILE:HG12	2.20	0.41
1:A:202:ARG:HA	1:A:202:ARG:HD3	1.93	0.41
1:A:7:HIS:HB2	1:A:10:PHE:CE1	2.56	0.40
1:A:60:TYR:CB	1:A:93:PRO:HB3	2.51	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:137:LYS:NZ	1:B:348:ARG:CD[3_555]	2.10	0.10

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	337/344 (98%)	325 (96%)	8 (2%)	4 (1%)	15	46
1	B	336/344 (98%)	326 (97%)	10 (3%)	0	100	100
All	All	673/688 (98%)	651 (97%)	18 (3%)	4 (1%)	28	64

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	57	GLY
1	A	56	ARG
1	A	133	ASP
1	A	228	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	288/289 (100%)	243 (84%)	45 (16%)	3	9
1	B	287/289 (99%)	243 (85%)	44 (15%)	3	10
All	All	575/578 (100%)	486 (84%)	89 (16%)	3	9

All (89) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	27	LEU
1	A	42	VAL
1	A	47	LEU
1	A	53	LYS
1	A	54	THR
1	A	55	SER
1	A	56	ARG
1	A	58	PHE
1	A	59	THR
1	A	108	VAL
1	A	116	LYS
1	A	130	LEU
1	A	131	THR
1	A	134	LEU
1	A	135	THR
1	A	137	LYS
1	A	143	LYS
1	A	146	ARG
1	A	151	THR

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	162	LYS
1	A	172	THR
1	A	173	GLU
1	A	202	ARG
1	A	205	ARG
1	A	206	SER
1	A	207	GLU
1	A	208	LEU
1	A	229	GLU
1	A	235	ASP
1	A	236	ASP
1	A	237	ASP
1	A	248	MET
1	A	260	SER
1	A	261	MET
1	A	262	ARG
1	A	273	SER
1	A	284	ILE
1	A	297	LYS
1	A	304	GLN
1	A	318	GLN
1	A	324	LYS
1	A	331	LEU
1	A	339	ILE
1	A	348	ARG
1	A	349	ASP
1	B	12	ARG
1	B	26	LYS
1	B	27	LEU
1	B	42	VAL
1	B	47	LEU
1	B	53	LYS
1	B	58	PHE
1	B	92	LYS
1	B	108	VAL
1	B	116	LYS
1	B	130	LEU
1	B	132	ASN
1	B	134	LEU
1	B	135	THR
1	B	146	ARG
1	B	151	THR

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	162	LYS
1	B	169	ASP
1	B	172	THR
1	B	173	GLU
1	B	182	CYS
1	B	203	LYS
1	B	205	ARG
1	B	207	GLU
1	B	208	LEU
1	B	221	ILE
1	B	224	LEU
1	B	227	LYS
1	B	229	GLU
1	B	239	ILE
1	B	243	ILE
1	B	250	LYS
1	B	260	SER
1	B	261	MET
1	B	273	SER
1	B	284	ILE
1	B	297	LYS
1	B	304	GLN
1	B	318	GLN
1	B	324	LYS
1	B	331	LEU
1	B	339	ILE
1	B	348	ARG
1	B	349	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	7	HIS
1	A	20	GLN
1	A	96	ASN
1	A	319	GLN
1	A	325	ASN
1	B	20	GLN
1	B	84	GLN
1	B	96	ASN
1	B	319	GLN
1	B	325	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	341/344 (99%)	0.32	11 (3%) 48 42	40, 53, 65, 73	0
1	B	340/344 (98%)	0.38	14 (4%) 38 32	43, 57, 77, 89	0
All	All	681/688 (98%)	0.35	25 (3%) 42 37	40, 55, 70, 89	0

All (25) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	134	LEU	4.5
1	B	238	ASP	4.0
1	B	130	LEU	3.6
1	B	132	ASN	3.1
1	B	7	HIS	3.0
1	A	238	ASP	2.7
1	A	132	ASN	2.6
1	A	12	ARG	2.6
1	A	261	MET	2.5
1	A	237	ASP	2.4
1	A	234	GLY	2.4
1	A	53	LYS	2.2
1	B	260	SER	2.2
1	B	223	VAL	2.2
1	B	157	TRP	2.2
1	B	222	LYS	2.1
1	B	128	HIS	2.1
1	B	137	LYS	2.1
1	B	234	GLY	2.1
1	A	56	ARG	2.1
1	A	6	TYR	2.1
1	B	205	ARG	2.1
1	A	27	LEU	2.0
1	A	229	GLU	2.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	170	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.