



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 6, 2018 – 10:56 AM EST

PDB ID : 3A37
Title : Structural insight into the membrane insertion of tail-anchored proteins by Get3
Authors : Yamagata, A.; Mimura, H.; Sato, Y.; Yamashita, M.; Yoshikawa, A.; Fukai, S.
Deposited on : 2009-06-10
Resolution : 3.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix)	:	1.9-1692
EDS	:	rb-20030736
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	rb-20030736


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	1692 (3.00-3.00)
Clashscore	112137	2037 (3.00-3.00)
Ramachandran outliers	110173	1973 (3.00-3.00)
Sidechain outliers	110143	1976 (3.00-3.00)
RSRZ outliers	101464	1716 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	362	<div> <div>2%</div> <div> <div></div> <div>47%</div> <div>35%</div> <div>•</div> <div>14%</div> </div> </div>
1	B	362	<div> <div>3%</div> <div> <div></div> <div>53%</div> <div>32%</div> <div>•</div> <div>12%</div> </div> </div>

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 4925 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ATPase GET3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	313	Total	C	N	O	S	0	0	0
			2398	1514	401	467	16			
1	B	319	Total	C	N	O	S	0	0	0
			2472	1560	414	481	17			

There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	155	ASP	GLY	CONFLICT	UNP Q12154
A	355	LEU	-	EXPRESSION TAG	UNP Q12154
A	356	GLU	-	EXPRESSION TAG	UNP Q12154
A	357	HIS	-	EXPRESSION TAG	UNP Q12154
A	358	HIS	-	EXPRESSION TAG	UNP Q12154
A	359	HIS	-	EXPRESSION TAG	UNP Q12154
A	360	HIS	-	EXPRESSION TAG	UNP Q12154
A	361	HIS	-	EXPRESSION TAG	UNP Q12154
A	362	HIS	-	EXPRESSION TAG	UNP Q12154
B	155	ASP	GLY	CONFLICT	UNP Q12154
B	355	LEU	-	EXPRESSION TAG	UNP Q12154
B	356	GLU	-	EXPRESSION TAG	UNP Q12154
B	357	HIS	-	EXPRESSION TAG	UNP Q12154
B	358	HIS	-	EXPRESSION TAG	UNP Q12154
B	359	HIS	-	EXPRESSION TAG	UNP Q12154
B	360	HIS	-	EXPRESSION TAG	UNP Q12154
B	361	HIS	-	EXPRESSION TAG	UNP Q12154
B	362	HIS	-	EXPRESSION TAG	UNP Q12154

- Molecule 2 is ADENOSINE-5'-DIPHOSPHATE (three-letter code: ADP) (formula: $C_{10}H_{15}N_5O_{10}P_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total 27	C 10	N 5	O 10	P 2	0	0
2	B	1	Total 27	C 10	N 5	O 10	P 2	0	0

- Molecule 3 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Zn	0	0
			1	1		

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

[illegible]

Chain B:

3% 53% 32% 12%

Label	Value
E349	1
L380	2
GLU	3
ASP	4
LNS	5
GLU	6
LEU	7
GLU	8
HIS	9
HIS	10
HIS	11
HIS	12
HIS	13
HIS	14
HIS	15
HIS	16
HIS	17
HIS	18
HIS	19
HIS	20
HIS	21
HIS	22
HIS	23
HIS	24
HIS	25
HIS	26
HIS	27
HIS	28
HIS	29
HIS	30
HIS	31
HIS	32
HIS	33
HIS	34
HIS	35
HIS	36
HIS	37
HIS	38
HIS	39
HIS	40
HIS	41
HIS	42
HIS	43
HIS	44
HIS	45
HIS	46
HIS	47
HIS	48
HIS	49
HIS	50
HIS	51
HIS	52
HIS	53
HIS	54
HIS	55
HIS	56
HIS	57
HIS	58
HIS	59
HIS	60
HIS	61
HIS	62
HIS	63
HIS	64
HIS	65
HIS	66
HIS	67
HIS	68
HIS	69
HIS	70
HIS	71
HIS	72
HIS	73
HIS	74
HIS	75
HIS	76
HIS	77
HIS	78
HIS	79
HIS	80
HIS	81
HIS	82
HIS	83
HIS	84
HIS	85
HIS	86
HIS	87
HIS	88
HIS	89
HIS	90
HIS	91
HIS	92
HIS	93
HIS	94
HIS	95
HIS	96
HIS	97
HIS	98
HIS	99
HIS	100
HIS	101
HIS	102
HIS	103
HIS	104
HIS	105
HIS	106
HIS	107
HIS	108
HIS	109
HIS	110
HIS	111
HIS	112
HIS	113
HIS	114
HIS	115
HIS	116
HIS	117
HIS	118
HIS	119
HIS	120
HIS	121
HIS	122
HIS	123
HIS	124
HIS	125
HIS	126
HIS	127
HIS	128
HIS	129
HIS	130
HIS	131
HIS	132
HIS	133
HIS	134
HIS	135
HIS	136
HIS	137
HIS	138
HIS	139
HIS	140
HIS	141
HIS	142
HIS	143
HIS	144
HIS	145
HIS	146
HIS	147
HIS	148
HIS	149
HIS	150
HIS	151
HIS	152
HIS	153
HIS	154
HIS	155
HIS	156
HIS	157
HIS	158
HIS	159
HIS	160
HIS	161
HIS	162
HIS	163
HIS	164
HIS	165
HIS	166
HIS	167
HIS	168
HIS	169
HIS	170
HIS	171
HIS	172
HIS	173
HIS	174
HIS	175
HIS	176
HIS	177
HIS	178
HIS	179
HIS	180
HIS	181
HIS	182
HIS	183
HIS	184
HIS	185
HIS	186
HIS	187
HIS	188
HIS	189
HIS	190
HIS	191
HIS	192
HIS	193
HIS	194
HIS	195
HIS	196
HIS	197
HIS	198
HIS	199
HIS	200
HIS	201
HIS	202
HIS	203
HIS	204
HIS	205
HIS	206
HIS	207
HIS	208
HIS	209
HIS	210
HIS	211
HIS	212
HIS	213
HIS	214
HIS	215
HIS	216
HIS	217
HIS	218
HIS	219
HIS	220
HIS	221
HIS	222
HIS	223
HIS	224
HIS	225
HIS	226
HIS	

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	115.23Å 222.55Å 49.29Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.29 – 3.00 49.29 – 3.00	Depositor EDS
% Data completeness (in resolution range)	97.2 (49.29-3.00) 97.3 (49.29-3.00)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.06	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.77 (at 3.01Å)	Xtriage
Refinement program	CNS 1.2	Depositor
R, R_{free}	0.253 , 0.282 0.253 , 0.281	Depositor DCC
R_{free} test set	1233 reflections (4.82%)	DCC
Wilson B-factor (Å ²)	71.0	Xtriage
Anisotropy	0.482	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 44.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.42$, $\langle L^2 \rangle = 0.25$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	4925	wwPDB-VP
Average B, all atoms (Å ²)	93.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.64% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, ADP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.43	0/2438	0.68	2/3297 (0.1%)
1	B	0.45	0/2510	0.66	0/3389
All	All	0.44	0/4948	0.67	2/6686 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	275	LEU	CA-CB-CG	5.47	127.88	115.30
1	A	52	LEU	CA-CB-CG	5.29	127.47	115.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	250	TYR	Sidechain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2398	0	2330	134	0
1	B	2472	0	2442	90	0
2	A	27	0	12	0	0
2	B	27	0	12	1	0
3	A	1	0	0	0	0
All	All	4925	0	4796	221	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 23.

The worst 5 of 221 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:242:CYS:HB3	1:A:252:THR:HG21	1.37	1.04
1:B:20:TRP:HB2	1:B:236:THR:HB	1.50	0.94
1:B:52:LEU:HD21	1:B:86:MET:HG2	1.50	0.93
1:A:8:ASN:HA	1:A:312:VAL:HG22	1.55	0.88
1:B:243:ILE:HD12	1:B:244:SER:H	1.36	0.88

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	309/362 (85%)	278 (90%)	26 (8%)	5 (2%)	11	46
1	B	313/362 (86%)	283 (90%)	24 (8%)	6 (2%)	9	41
All	All	622/724 (86%)	561 (90%)	50 (8%)	11 (2%)	10	43

5 of 11 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	173	THR
1	B	173	THR
1	B	283	HIS
1	A	280	ASP
1	A	285	CYS

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	260/318 (82%)	242 (93%)	18 (7%)	18	53
1	B	274/318 (86%)	254 (93%)	20 (7%)	16	50
All	All	534/636 (84%)	496 (93%)	38 (7%)	17	52

5 of 38 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	329	LYS
1	B	64	ASP
1	B	313	LYS
1	B	14	THR
1	B	84	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 16 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	46	GLN
1	B	50	GLN
1	B	229	GLN
1	A	310	HIS
1	B	267	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 1 is monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	ADP	A	400	-	25,29,29	2.01	5 (20%)	24,45,45	2.74	4 (16%)
2	ADP	B	401	-	25,29,29	1.86	6 (24%)	24,45,45	2.67	2 (8%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	ADP	A	400	-	-	0/12/32/32	0/3/3/3
2	ADP	B	401	-	-	0/12/32/32	0/3/3/3

The worst 5 of 11 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	400	ADP	PB-O3A	-4.71	1.52	1.60
2	B	401	ADP	PB-O3A	-3.47	1.54	1.60
2	A	400	ADP	C5-N7	-3.23	1.28	1.39
2	B	401	ADP	C5-N7	-3.15	1.28	1.39
2	B	401	ADP	PA-O2A	-2.07	1.44	1.55

The worst 5 of 6 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
2	B	401	ADP	N3-C2-N1	-12.09	118.33	128.86
2	A	400	ADP	N3-C2-N1	-11.90	118.50	128.86
2	A	400	ADP	C4-C5-N7	-2.11	107.37	109.41
2	B	401	ADP	C4'-O4'-C1'	2.32	112.24	109.77
2	A	400	ADP	C2'-C3'-C4'	2.67	107.83	102.62

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	401	ADP	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	313/362 (86%)	0.06	7 (2%) 62 33	58, 91, 139, 155	0
1	B	319/362 (88%)	0.01	12 (3%) 41 17	54, 84, 143, 159	0
All	All	632/724 (87%)	0.03	19 (3%) 51 23	54, 88, 140, 159	0

The worst 5 of 19 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	212	ILE	5.0
1	B	122	GLN	4.3
1	B	119	SER	3.5
1	B	211	ASP	3.3
1	B	121	LEU	3.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	ADP	B	401	27/27	0.89	0.25	0.03	93,100,103,104	0
3	ZN	A	363	1/1	0.99	0.12	-1.02	97,97,97,97	0
2	ADP	A	400	27/27	0.95	0.18	-1.27	97,105,107,107	0

6.5 Other polymers [i](#)

There are no such residues in this entry.