



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 14, 2017 – 02:55 pm GMT

PDB ID : 4A3Y  
Title : Crystal structure of Raucaffricine glucosidase from ajmaline biosynthesis pathway  
Authors : Xia, L.; Ruppert, M.; Wang, M.; Panjekar, S.; Barleben, L.; Rajendran, C.; Lin, H.; Stoeckigt, J.  
Deposited on : 2011-10-06  
Resolution : 2.15 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix)	:	1.9-1692
EDS	:	trunk28620
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	recalc28949


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.15 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	1170 (2.16-2.16)
Clashscore	112137	1278 (2.16-2.16)
Ramachandran outliers	110173	1256 (2.16-2.16)
Sidechain outliers	110143	1255 (2.16-2.16)
RSRZ outliers	101464	1175 (2.16-2.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	540	<div> <div>3%</div> <div> <div></div> <div>78%</div> <div>8%</div> <div>13%</div> </div> </div>
1	B	540	<div> <div>2%</div> <div> <div></div> <div>75%</div> <div>10%</div> <div>13%</div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	SO4	A	1514	-	-	-	X
2	SO4	B	1514	-	-	-	X
3	GOL	A	1516	-	-	-	X
3	GOL	B	1515	-	-	-	X

## 2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 8016 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called RAUCAFFRICINE-O-BETA-D-GLUCOSIDASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	469	Total	C	N	O	S	0	3	0
			3775	2416	641	705	13			
1	B	468	Total	C	N	O	S	0	4	0
			3771	2416	637	705	13			

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	230	Total	O	0	0
			230	230		
4	B	213	Total	O	0	0
			213	213		


### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: RAUCAFFRICINE-O-BETA-D-GLUCOSIDASE



#### • Molecule 1: RAUCAFFRICINE-O-BETA-D-GLUCOSIDASE



## 4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	102.77Å 127.33Å 215.84Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	19.99 – 2.15 19.99 – 2.15	Depositor EDS
% Data completeness (in resolution range)	99.3 (19.99-2.15) 99.3 (19.99-2.15)	Depositor EDS
$R_{merge}$	0.12	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	4.22 (at 2.15Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, $R_{free}$	0.165 , 0.204 0.178 , 0.215	Depositor DCC
$R_{free}$ test set	1054 reflections (1.40%)	DCC
Wilson B-factor (Å <sup>2</sup> )	28.2	Xtriage
Anisotropy	0.112	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 29.8	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	8016	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	36.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.60% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	1.09	3/3895 (0.1%)	0.88	11/5288 (0.2%)
1	B	1.06	4/3894 (0.1%)	0.86	15/5287 (0.3%)
All	All	1.07	7/7789 (0.1%)	0.87	26/10575 (0.2%)

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	80	GLU	CD-OE1	8.73	1.35	1.25
1	B	435	GLU	CG-CD	7.13	1.62	1.51
1	B	165	GLU	CG-CD	-6.20	1.42	1.51
1	B	398	GLU	CG-CD	5.86	1.60	1.51
1	A	252	ALA	CA-CB	5.23	1.63	1.52
1	B	404	LEU	CG-CD2	-5.04	1.33	1.51
1	A	511	PHE	CG-CD1	-5.04	1.31	1.38

All (26) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	94	ARG	NE-CZ-NH1	7.98	124.29	120.30
1	B	42	ARG	NE-CZ-NH2	-7.24	116.68	120.30
1	B	404	LEU	CB-CG-CD1	6.92	122.77	111.00
1	B	404	LEU	CA-CB-CG	6.91	131.18	115.30
1	A	60	ASP	CB-CG-OD1	6.56	124.20	118.30
1	B	206	ARG	NE-CZ-NH2	-6.53	117.03	120.30
1	B	60	ASP	CB-CG-OD1	6.40	124.06	118.30
1	A	94	ARG	NE-CZ-NH2	6.26	123.43	120.30
1	A	115	GLU	OE1-CD-OE2	-5.92	116.20	123.30
1	A	415	LEU	CA-CB-CG	5.64	128.26	115.30
1	A	149	ASP	CB-CG-OD2	5.59	123.33	118.30
1	A	80	GLU	OE1-CD-OE2	5.51	129.91	123.30

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	435	GLU	OE1-CD-OE2	-5.43	116.78	123.30
1	B	91	ASP	CB-CA-C	-5.38	99.64	110.40
1	B	162	ASP	CB-CG-OD2	5.38	123.14	118.30
1	B	415	LEU	CA-CB-CG	5.35	127.61	115.30
1	B	378	ASP	CB-CG-OD1	5.27	123.04	118.30
1	A	454	ARG	NE-CZ-NH1	5.25	122.93	120.30
1	B	294	ARG	NE-CZ-NH1	5.24	122.92	120.30
1	B	206	ARG	NE-CZ-NH1	5.21	122.91	120.30
1	B	425	ASP	CB-CG-OD1	5.18	122.96	118.30
1	A	323	ARG	NE-CZ-NH1	5.17	122.89	120.30
1	B	282	ASP	CB-CG-OD2	5.12	122.91	118.30
1	A	323	ARG	NE-CZ-NH2	-5.07	117.77	120.30
1	A	162	ASP	CB-CG-OD2	5.04	122.83	118.30
1	A	142	ASP	CB-CG-OD1	5.00	122.80	118.30

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3775	0	3577	29	0
1	B	3771	0	3570	39	0
2	A	10	0	0	0	0
2	B	5	0	0	1	0
3	A	6	0	8	0	0
3	B	6	0	8	0	0
4	A	230	0	0	7	0
4	B	213	0	0	9	0
All	All	8016	0	7163	69	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (69) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:1514:SO4:O1	4:B:2180:HOH:O	1.64	1.11
1:B:80[A]:GLU:OE2	4:B:2057:HOH:O	1.81	0.96
1:A:165:GLU:CD	4:A:2110:HOH:O	2.11	0.89
1:A:16:ARG:HH11	1:A:16:ARG:HG2	1.38	0.89
1:B:283:GLU:OE2	4:B:2134:HOH:O	1.88	0.89
1:A:165:GLU:HG3	4:A:2110:HOH:O	1.81	0.79
1:A:165:GLU:CG	4:A:2110:HOH:O	2.31	0.75
1:A:428:ASN:OD1	1:A:430:ASN:HB2	1.88	0.74
1:B:453:VAL:HG12	1:B:457:MET:HE3	1.70	0.73
1:A:185:ASN:HD21	1:A:345:ASN:HD21	1.36	0.72
1:B:183:THR:OG1	1:B:250:HIS:HD2	1.74	0.71
1:B:94:ARG:HG2	4:B:2065:HOH:O	1.90	0.69
1:B:428:ASN:OD1	1:B:430:ASN:HB2	1.93	0.69
1:A:16:ARG:HH11	1:A:16:ARG:CG	2.05	0.69
1:B:73:ASP:OD1	1:B:76:HIS:HD2	1.76	0.69
1:B:185:ASN:HD21	1:B:345:ASN:HD21	1.42	0.67
1:A:183:THR:OG1	1:A:250:HIS:HD2	1.77	0.67
1:B:283:GLU:HG2	4:B:2134:HOH:O	1.93	0.67
1:B:16:ARG:HH11	1:B:16:ARG:HB3	1.64	0.63
1:A:445:TYR:O	1:A:449:HIS:HD2	1.83	0.62
1:B:76:HIS:HE1	4:B:2053:HOH:O	1.82	0.61
1:A:244:HIS:HD2	1:A:312:TYR:OH	1.86	0.58
1:A:16:ARG:NH1	1:A:16:ARG:HG2	2.08	0.57
1:B:457:MET:HE2	1:B:463:VAL:HB	1.87	0.56
1:B:283:GLU:CG	4:B:2134:HOH:O	2.52	0.55
1:B:26:PHE:CE1	1:B:457:MET:HE1	2.43	0.54
1:B:71:ALA:HA	1:B:481:TYR:OH	2.08	0.53
1:B:370:ASP:O	1:B:372:HIS:HD2	1.91	0.52
1:B:121:ASN:HD21	1:B:177:ARG:HH22	1.57	0.52
1:A:154:PHE:CE1	1:A:246:LEU:HD23	2.45	0.52
1:A:397:PRO:HA	1:A:449:HIS:CE1	2.44	0.51
1:B:445:TYR:O	1:B:449:HIS:HD2	1.93	0.51
1:B:185:ASN:HD21	1:B:345:ASN:ND2	2.07	0.49
1:A:185:ASN:ND2	1:A:345:ASN:HD21	2.08	0.49
1:A:185:ASN:HD21	1:A:345:ASN:ND2	2.07	0.48
1:B:397:PRO:HA	1:B:449:HIS:CE1	2.48	0.48
1:B:449:HIS:HE1	4:B:2148:HOH:O	1.97	0.48
1:A:370:ASP:O	1:A:372:HIS:HD2	1.97	0.47
1:A:94:ARG:HG2	4:A:2067:HOH:O	2.13	0.47
1:B:346:TYR:OH	1:B:399:GLY:HA3	2.14	0.47
1:A:346:TYR:OH	1:A:399:GLY:HA3	2.15	0.46
1:B:121:ASN:HD22	1:B:177:ARG:HH12	1.64	0.46

Continued on next page...

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:445:TYR:O	1:A:449:HIS:CD2	2.66	0.46
1:A:71:ALA:HA	1:A:481:TYR:OH	2.16	0.45
1:B:154:PHE:CE1	1:B:246:LEU:HD23	2.50	0.45
1:B:235:PRO:HB2	1:B:366:SER:HB2	1.99	0.45
1:B:91:ASP:OD1	1:B:513:LYS:HE3	2.17	0.45
1:B:16:ARG:HH11	1:B:16:ARG:CG	2.30	0.45
1:B:16:ARG:CB	1:B:16:ARG:HH11	2.28	0.44
1:B:121:ASN:ND2	1:B:177:ARG:HH22	2.16	0.43
1:B:94:ARG:NH2	1:B:420:GLU:OE2	2.51	0.43
1:A:251:ALA:HB2	1:A:335:LEU:HD23	2.01	0.42
1:B:401:ARG:O	1:B:405:VAL:HG23	2.19	0.42
1:B:396:TYR:OH	1:B:402:LYS:HE3	2.19	0.42
1:A:244:HIS:HE1	4:A:2163:HOH:O	2.03	0.42
1:A:422:GLY:HA3	1:A:469:TRP:O	2.19	0.42
1:B:183:THR:HG1	1:B:250:HIS:HD2	1.68	0.42
1:B:469:TRP:HA	1:B:470:SER:HA	1.73	0.42
1:A:243:THR:HG21	1:A:302:TRP:NE1	2.35	0.42
1:B:433:LEU:N	4:B:2180:HOH:O	2.52	0.41
1:A:508:MET:O	1:A:512:HIS:HB2	2.19	0.41
1:A:89:GLY:O	1:A:512:HIS:HD2	2.03	0.41
1:A:250:HIS:CE1	1:A:338:SER:O	2.74	0.41
1:A:414:PRO:HD2	4:A:2167:HOH:O	2.18	0.41
1:B:243:THR:HG23	1:B:303:PHE:CE1	2.56	0.41
1:B:445:TYR:O	1:B:449:HIS:CD2	2.73	0.41
1:A:243:THR:HG21	1:A:302:TRP:CE2	2.56	0.41
1:B:244:HIS:CE1	1:B:327:PHE:CE1	3.08	0.41
4:A:2047:HOH:O	1:B:493:ASN:ND2	2.27	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	466/540 (86%)	450 (97%)	15 (3%)	1 (0%)	51	50
1	B	465/540 (86%)	451 (97%)	14 (3%)	0	100	100
All	All	931/1080 (86%)	901 (97%)	29 (3%)	1 (0%)	55	56

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	388	SER

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	397/461 (86%)	394 (99%)	3 (1%)	85	89
1	B	396/461 (86%)	387 (98%)	9 (2%)	56	59
All	All	793/922 (86%)	781 (98%)	12 (2%)	73	75

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	16	ARG
1	A	326	LYS
1	A	513	LYS
1	B	16	ARG
1	B	20	SER
1	B	80[A]	GLU
1	B	80[B]	GLU
1	B	206	ARG
1	B	326	LYS
1	B	364[A]	ASN
1	B	364[B]	ASN
1	B	404	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (19) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	140	HIS
1	A	185	ASN
1	A	244	HIS
1	A	250	HIS
1	A	354	ASN
1	A	421	ASN
1	A	449	HIS
1	A	512	HIS
1	B	36	GLN
1	B	76	HIS
1	B	121	ASN
1	B	185	ASN
1	B	250	HIS
1	B	354	ASN
1	B	372	HIS
1	B	421	ASN
1	B	449	HIS
1	B	452	ASN
1	B	512	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected

value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	SO4	A	1514	-	4,4,4	0.44	0	6,6,6	0.38	0
2	SO4	A	1515	-	4,4,4	0.54	0	6,6,6	0.59	0
3	GOL	A	1516	-	5,5,5	1.06	0	5,5,5	0.70	0
2	SO4	B	1514	-	4,4,4	0.30	0	6,6,6	0.57	0
3	GOL	B	1515	-	5,5,5	1.17	0	5,5,5	0.61	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	SO4	A	1514	-	-	0/0/0/0	0/0/0/0
2	SO4	A	1515	-	-	0/0/0/0	0/0/0/0
3	GOL	A	1516	-	-	0/4/4/4	0/0/0/0
2	SO4	B	1514	-	-	0/0/0/0	0/0/0/0
3	GOL	B	1515	-	-	0/4/4/4	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	1514	SO4	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	469/540 (86%)	-0.36	14 (2%) 51 59	26, 34, 50, 66	0
1	B	468/540 (86%)	-0.39	9 (1%) 67 74	27, 36, 51, 62	0
All	All	937/1080 (86%)	-0.38	23 (2%) 58 65	26, 35, 51, 66	0

All (23) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	380	ASN	4.3
1	B	364[A]	ASN	3.8
1	A	284	ASN	3.3
1	B	330	GLU	3.1
1	A	206	ARG	2.9
1	B	232	THR	2.9
1	B	206	ARG	2.8
1	A	364	ASN	2.8
1	A	16	ARG	2.8
1	B	247	LEU	2.7
1	B	284	ASN	2.5
1	A	381	GLY	2.5
1	A	356	SER	2.5
1	A	330	GLU	2.4
1	A	24	ALA	2.4
1	A	283	GLU	2.3
1	A	13	ASP	2.3
1	A	453	VAL	2.2
1	B	283	GLU	2.2
1	B	16	ARG	2.1
1	A	247	LEU	2.1
1	B	450	ILE	2.1
1	A	513	LYS	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
3	GOL	B	1515	6/6	0.82	0.22	12.09	34,38,41,42	0
2	SO4	A	1514	5/5	0.92	0.21	7.88	28,32,33,36	5
3	GOL	A	1516	6/6	0.89	0.17	6.14	32,36,39,44	0
2	SO4	B	1514	5/5	0.96	0.16	3.71	29,30,31,38	5
2	SO4	A	1515	5/5	0.93	0.15	-	30,31,36,41	5

## 6.5 Other polymers [i](#)

There are no such residues in this entry.