



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 14, 2017 – 05:52 pm GMT

PDB ID : 5A59  
Title : The structure of GH101 E796Q mutant from *Streptococcus pneumoniae* TIGR4 in complex with T-antigen  
Authors : Gregg, K.J.; Suits, M.D.L.; Deng, L.; Vocadlo, D.J.; Boraston, A.B.  
Deposited on : 2015-06-16  
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.2 (RC1), CSD as538be (2017)  
Xtriage (Phenix) : 1.9-1692  
EDS : trunk28620  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : recalc28949


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	3846 (2.50-2.50)
Clashscore	112137	4554 (2.50-2.50)
Ramachandran outliers	110173	4463 (2.50-2.50)
Sidechain outliers	110143	4465 (2.50-2.50)
RSRZ outliers	101464	3876 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	1117	<div> <div></div> <div>91%</div> <div>8%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	CIT	A	2431	-	-	-	X
4	CIT	A	2432	-	-	-	X

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
6	EDO	A	2436	-	-	-	X
6	EDO	A	2437	-	-	-	X
6	EDO	A	2440	-	-	-	X
6	EDO	A	2442	-	-	-	X
6	EDO	A	2447	-	-	-	X
6	EDO	A	2448	-	-	-	X
6	EDO	A	2451	-	-	-	X
6	EDO	A	2452	-	-	-	X
6	EDO	A	2454	-	-	-	X
6	EDO	A	2455	-	-	-	X
6	EDO	A	2456	-	-	-	X
6	EDO	A	2457	-	-	-	X
6	EDO	A	2458	-	-	-	X
6	EDO	A	2459	-	-	-	X
6	EDO	A	2465	-	-	-	X

## 2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 10107 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ENDO-ALPHA-N-ACETYLGALACTOSAMINIDASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	1106	8891	5602	1521	1743	25	0	27	0

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	316	MET	-	EXPRESSION TAG	UNP Q2MGH6
A	1427	HIS	-	EXPRESSION TAG	UNP Q2MGH6
A	1428	HIS	-	EXPRESSION TAG	UNP Q2MGH6
A	1429	HIS	-	EXPRESSION TAG	UNP Q2MGH6
A	1430	HIS	-	EXPRESSION TAG	UNP Q2MGH6
A	1431	HIS	-	EXPRESSION TAG	UNP Q2MGH6
A	1432	HIS	-	EXPRESSION TAG	UNP Q2MGH6
A	796	GLN	GLU	ENGINEERED MUTATION	UNP Q2MGH6
A	461	ASP	ASN	CONFLICT	UNP Q2MGH6
A	1165	ASN	ASP	CONFLICT	UNP Q2MGH6
A	1202	GLU	GLN	CONFLICT	UNP Q2MGH6
A	1264	ASP	ASN	CONFLICT	UNP Q2MGH6

- Molecule 2 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	3	Total	Ca	0	0
			3	3		

- Molecule 3 is MANGANESE (II) ION (three-letter code: MN) (formula: Mn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Mn	0	0
			1	1		

- Molecule 4 is CITRIC ACID (three-letter code: CIT) (formula: C<sub>6</sub>H<sub>8</sub>O<sub>7</sub>).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			13	6	7		
4	A	1	Total	C	O	0	0
			13	6	7		

- Molecule 5 is a polymer of unknown type called SUGAR (2-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	2	Total	C	N	O	0	0
			26	14	1	11		

- Molecule 6 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C<sub>2</sub>H<sub>6</sub>O<sub>2</sub>).



[illegible]

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			4	2	2		
6	A	1	Total	C	O	0	0
			4	2	2		
6	A	1	Total	C	O	0	0
			4	2	2		
6	A	1	Total	C	O	0	0
			4	2	2		
6	A	1	Total	C	O	0	0
			4	2	2		
6	A	1	Total	C	O	0	0
			4	2	2		
6	A	1	Total	C	O	0	0
			4	2	2		
6	A	1	Total	C	O	0	0
			4	2	2		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	1036	Total	O	0	0
			1036	1036		

### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: ENDO-ALPHA-N-ACETYL GALACTOSAMINIDASE





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 2 <sub>1</sub> 2 <sub>1</sub> 2 <sub>1</sub>	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	87.20Å 121.89Å 139.34Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	91.74 – 2.50 27.17 – 2.50	Depositor EDS
% Data completeness (in resolution range)	99.8 (91.74-2.50) 99.9 (27.17-2.50)	Depositor EDS
$R_{merge}$	0.16	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.16 (at 2.51Å)	Xtriage
Refinement program	REFMAC 5.8.0073	Depositor
R, $R_{free}$	0.167 , 0.229 0.173 , 0.231	Depositor DCC
$R_{free}$ test set	2599 reflections (5.26%)	DCC
Wilson B-factor (Å <sup>2</sup> )	18.8	Xtriage
Anisotropy	0.655	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 36.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	10107	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	19.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.27% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CA, MN, EDO, GAL, CIT, A2G

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.61	0/9179	0.74	2/12435 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	349	ARG	NE-CZ-NH1	5.46	123.03	120.30
1	A	1403	ARG	NE-CZ-NH1	5.31	122.96	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	8891	0	8563	42	0
2	A	3	0	0	0	0
3	A	1	0	0	0	0
4	A	26	0	10	1	0
5	A	26	0	24	0	0
6	A	124	0	186	6	0
7	A	1036	0	0	8	0
All	All	10107	0	8783	46	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (46) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1026:ASP:HB3	7:A:3684:HOH:O	1.83	0.79
1:A:920[A]:ARG:NH2	1:A:922:GLU:OE2	2.29	0.66
6:A:2461:EDO:C2	7:A:4036:HOH:O	2.47	0.62
1:A:1158:TYR:CE2	1:A:1160:HIS:HB2	2.36	0.61
4:A:2432:CIT:O4	4:A:2432:CIT:O1	2.20	0.59
6:A:2437:EDO:H21	7:A:3966:HOH:O	2.02	0.58
1:A:549:TRP:CD2	1:A:568:PRO:HB3	2.39	0.57
1:A:795[A]:ILE:HD11	1:A:808:HIS:CE1	2.44	0.52
1:A:1238:ALA:HB3	6:A:2447:EDO:H11	1.91	0.51
1:A:549:TRP:CG	1:A:568:PRO:HB3	2.46	0.50
1:A:1130[B]:LYS:HB2	1:A:1198[B]:GLU:HG3	1.92	0.50
1:A:698:SER:HA	1:A:771:SER:HB3	1.94	0.50
1:A:1125:ASN:OD1	1:A:1204:THR:OG1	2.27	0.49
1:A:610:THR:HG21	1:A:643:LEU:HD13	1.92	0.49
1:A:1309[B]:GLU:OE1	1:A:1314:TYR:OH	2.31	0.48
1:A:440:ASP:HB3	1:A:443:LYS:HG3	1.94	0.48
1:A:371:PHE:HB2	1:A:449:SER:HB3	1.96	0.48
1:A:787:ASN:HA	1:A:791:TRP:O	2.13	0.47
1:A:351:LYS:HE3	7:A:3017:HOH:O	2.14	0.47
1:A:1146:THR:HB	1:A:1304:GLN:HB3	1.97	0.47
1:A:1267:THR:HB	1:A:1278:ASP:HB2	1.97	0.46
1:A:1119:VAL:HA	1:A:1210:ARG:O	2.16	0.45
1:A:1164:ARG:NH1	7:A:3836:HOH:O	2.48	0.45
1:A:1130[B]:LYS:HB2	1:A:1198[B]:GLU:CG	2.46	0.45
1:A:830:ARG:O	1:A:834:ASN:HB2	2.17	0.45
1:A:970:ASP:C	1:A:970:ASP:OD1	2.55	0.44
1:A:688:GLY:HA3	1:A:971:ALA:HB1	1.99	0.44
6:A:2461:EDO:H21	7:A:4036:HOH:O	2.17	0.43
1:A:545:HIS:CD2	1:A:545:HIS:N	2.86	0.43
1:A:965:THR:HG22	1:A:966:PRO:O	2.19	0.43
1:A:764:ASP:O	1:A:765:VAL:HB	2.18	0.42
1:A:1050:GLU:HB2	7:A:3702:HOH:O	2.19	0.42
1:A:638:LEU:HD11	1:A:1051:MET:HG3	2.01	0.42
1:A:1130[A]:LYS:HG3	1:A:1145:TYR:CD2	2.54	0.42
1:A:1357:THR:HG22	1:A:1365:LYS:HD2	2.03	0.41
1:A:1149:SER:HB3	1:A:1174:TYR:HB3	2.01	0.41
1:A:702:PRO:HA	1:A:707:PHE:CG	2.55	0.41

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:684:ALA:HB1	1:A:689:ALA:HB3	2.03	0.41
1:A:940:VAL:HA	1:A:945:TYR:CD2	2.56	0.41
1:A:1238:ALA:HB3	6:A:2447:EDO:C1	2.51	0.41
1:A:360:LEU:HA	1:A:360:LEU:HD23	1.96	0.41
1:A:359[B]:THR:CG2	7:A:3101:HOH:O	2.68	0.41
1:A:832:ILE:HG22	1:A:833:ARG:HG3	2.03	0.41
1:A:364[B]:VAL:HG22	6:A:2446:EDO:H22	2.03	0.40
1:A:808:HIS:CD2	1:A:839:ALA:HB2	2.56	0.40
1:A:1121:VAL:HG23	1:A:1209:ILE:HG13	2.03	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	1130/1117 (101%)	1086 (96%)	41 (4%)	3 (0%)	44 66

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	1227	LYS
1	A	662	LEU
1	A	765	VAL

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	957/942 (102%)	939 (98%)	18 (2%)	62 85

All (18) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	354[A]	SER
1	A	354[B]	SER
1	A	354[C]	SER
1	A	369	GLN
1	A	407	GLU
1	A	445[A]	LEU
1	A	445[B]	LEU
1	A	459	SER
1	A	532	TYR
1	A	788	LYS
1	A	846	SER
1	A	938	ASN
1	A	979	ASP
1	A	989	GLN
1	A	1166	ASN
1	A	1212	PHE
1	A	1239	GLN
1	A	1419	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	799	HIS

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

2 carbohydrates are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
5	GAL	A	2433	5	11,11,12	0.61	0	13,15,17	0.66	0
5	A2G	A	2434	5	15,15,15	0.53	0	21,21,21	1.30	4 (19%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	GAL	A	2433	5	-	0/2/19/22	0/1/1/1
5	A2G	A	2434	5	-	0/6/26/26	0/1/1/1

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	2434	A2G	C3-C4-C5	-2.34	106.09	110.22
5	A	2434	A2G	C8-C7-N2	-2.00	112.49	116.11
5	A	2434	A2G	C1-O-C5	2.14	117.25	113.39
5	A	2434	A2G	O4-C4-C5	2.46	115.49	109.28

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.6 Ligand geometry [i](#)

Of 37 ligands modelled in this entry, 4 are monoatomic - leaving 33 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
4	CIT	A	2431	-	3,12,12	2.40	2 (66%)	3,17,17	5.79	2 (66%)
4	CIT	A	2432	-	3,12,12	0.71	0	3,17,17	0.48	0
6	EDO	A	2435	-	3,3,3	0.51	0	2,2,2	0.28	0
6	EDO	A	2436	-	3,3,3	0.77	0	2,2,2	0.50	0
6	EDO	A	2437	-	3,3,3	0.66	0	2,2,2	0.33	0
6	EDO	A	2438	-	3,3,3	0.41	0	2,2,2	0.59	0
6	EDO	A	2439	-	3,3,3	0.55	0	2,2,2	0.23	0
6	EDO	A	2440	-	3,3,3	0.50	0	2,2,2	0.31	0
6	EDO	A	2441	-	3,3,3	0.45	0	2,2,2	0.34	0
6	EDO	A	2442	-	3,3,3	0.42	0	2,2,2	0.47	0
6	EDO	A	2443	-	3,3,3	0.40	0	2,2,2	0.37	0
6	EDO	A	2444	-	3,3,3	0.61	0	2,2,2	0.20	0
6	EDO	A	2445	-	3,3,3	0.44	0	2,2,2	0.53	0
6	EDO	A	2446	-	3,3,3	0.47	0	2,2,2	0.62	0
6	EDO	A	2447	-	3,3,3	0.63	0	2,2,2	0.36	0
6	EDO	A	2448	-	3,3,3	0.58	0	2,2,2	0.58	0
6	EDO	A	2449	-	3,3,3	0.82	0	2,2,2	0.40	0
6	EDO	A	2450	-	3,3,3	0.64	0	2,2,2	0.45	0
6	EDO	A	2451	-	3,3,3	0.81	0	2,2,2	0.69	0
6	EDO	A	2452	-	3,3,3	0.38	0	2,2,2	0.55	0
6	EDO	A	2453	-	3,3,3	0.38	0	2,2,2	0.61	0
6	EDO	A	2454	-	3,3,3	0.42	0	2,2,2	0.45	0
6	EDO	A	2455	-	3,3,3	0.55	0	2,2,2	0.15	0
6	EDO	A	2456	-	3,3,3	0.42	0	2,2,2	0.63	0
6	EDO	A	2457	-	3,3,3	0.52	0	2,2,2	0.18	0
6	EDO	A	2458	-	3,3,3	0.77	0	2,2,2	0.28	0
6	EDO	A	2459	-	3,3,3	0.76	0	2,2,2	0.25	0
6	EDO	A	2460	-	3,3,3	0.66	0	2,2,2	0.58	0
6	EDO	A	2461	-	3,3,3	0.46	0	2,2,2	0.27	0
6	EDO	A	2462	-	3,3,3	0.26	0	2,2,2	0.83	0
6	EDO	A	2463	-	3,3,3	0.42	0	2,2,2	0.49	0
6	EDO	A	2464	-	3,3,3	0.47	0	2,2,2	0.56	0
6	EDO	A	2465	-	3,3,3	0.60	0	2,2,2	0.27	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means

no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	CIT	A	2431	-	-	0/6/16/16	0/0/0/0
4	CIT	A	2432	-	-	0/6/16/16	0/0/0/0
6	EDO	A	2435	-	-	0/1/1/1	0/0/0/0
6	EDO	A	2436	-	-	0/1/1/1	0/0/0/0
6	EDO	A	2437	-	-	0/1/1/1	0/0/0/0
6	EDO	A	2438	-	-	0/1/1/1	0/0/0/0
6	EDO	A	2439	-	-	0/1/1/1	0/0/0/0
6	EDO	A	2440	-	-	0/1/1/1	0/0/0/0
6	EDO	A	2441	-	-	0/1/1/1	0/0/0/0
6	EDO	A	2442	-	-	0/1/1/1	0/0/0/0
6	EDO	A	2443	-	-	0/1/1/1	0/0/0/0
6	EDO	A	2444	-	-	0/1/1/1	0/0/0/0
6	EDO	A	2445	-	-	0/1/1/1	0/0/0/0
6	EDO	A	2446	-	-	0/1/1/1	0/0/0/0
6	EDO	A	2447	-	-	0/1/1/1	0/0/0/0
6	EDO	A	2448	-	-	0/1/1/1	0/0/0/0
6	EDO	A	2449	-	-	0/1/1/1	0/0/0/0
6	EDO	A	2450	-	-	0/1/1/1	0/0/0/0
6	EDO	A	2451	-	-	0/1/1/1	0/0/0/0
6	EDO	A	2452	-	-	0/1/1/1	0/0/0/0
6	EDO	A	2453	-	-	0/1/1/1	0/0/0/0
6	EDO	A	2454	-	-	0/1/1/1	0/0/0/0
6	EDO	A	2455	-	-	0/1/1/1	0/0/0/0
6	EDO	A	2456	-	-	0/1/1/1	0/0/0/0
6	EDO	A	2457	-	-	0/1/1/1	0/0/0/0
6	EDO	A	2458	-	-	0/1/1/1	0/0/0/0
6	EDO	A	2459	-	-	0/1/1/1	0/0/0/0
6	EDO	A	2460	-	-	0/1/1/1	0/0/0/0
6	EDO	A	2461	-	-	0/1/1/1	0/0/0/0
6	EDO	A	2462	-	-	0/1/1/1	0/0/0/0
6	EDO	A	2463	-	-	0/1/1/1	0/0/0/0
6	EDO	A	2464	-	-	0/1/1/1	0/0/0/0
6	EDO	A	2465	-	-	0/1/1/1	0/0/0/0

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	2431	CIT	C4-C3	-3.35	1.49	1.54
4	A	2431	CIT	C2-C3	-2.42	1.51	1.54

All (2) bond angle outliers are listed below:



Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	2431	CIT	C3-C2-C1	-3.69	109.18	114.95
4	A	2431	CIT	C4-C3-C2	9.26	132.80	109.75

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	2432	CIT	1	0
6	A	2437	EDO	1	0
6	A	2446	EDO	1	0
6	A	2447	EDO	2	0
6	A	2461	EDO	2	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	1106/1117 (99%)	-0.54	6 (0%) 90 91	10, 17, 33, 57	0

All (6) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	1345	GLN	3.9
1	A	1346	ALA	3.1
1	A	1348	ASN	2.6
1	A	431	GLN	2.3
1	A	317	GLU	2.2
1	A	1419	THR	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
5	GAL	A	2433	11/12	0.98	0.10	-0.22	16,17,18,18	0
5	A2G	A	2434	15/15	0.96	0.09	-1.40	15,20,25,27	0

## 6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
6	EDO	A	2437	4/4	0.79	0.34	21.47	50,53,55,56	0
6	EDO	A	2454	4/4	0.95	0.24	12.61	44,46,47,48	0
6	EDO	A	2455	4/4	0.91	0.22	10.51	35,37,38,41	0
6	EDO	A	2456	4/4	0.90	0.28	8.90	41,43,45,45	0
6	EDO	A	2442	4/4	0.93	0.20	6.99	43,43,44,47	0
6	EDO	A	2458	4/4	0.77	0.28	6.88	41,45,49,51	0
6	EDO	A	2457	4/4	0.89	0.27	6.45	47,47,49,49	0
6	EDO	A	2451	4/4	0.91	0.33	6.34	26,29,31,34	0
6	EDO	A	2436	4/4	0.90	0.27	6.12	36,41,44,46	0
6	EDO	A	2440	4/4	0.81	0.24	5.51	46,49,50,50	0
6	EDO	A	2452	4/4	0.93	0.18	5.12	37,40,42,44	0
6	EDO	A	2459	4/4	0.89	0.20	4.79	24,26,28,30	0
6	EDO	A	2465	4/4	0.81	0.27	4.79	39,41,41,41	0
6	EDO	A	2448	4/4	0.92	0.27	3.92	30,30,31,31	0
4	CIT	A	2432	13/13	0.85	0.30	3.73	39,53,60,63	0
6	EDO	A	2447	4/4	0.97	0.17	3.43	22,23,23,24	0
4	CIT	A	2431	13/13	0.93	0.23	2.19	22,26,29,30	0
6	EDO	A	2449	4/4	0.92	0.18	1.83	26,27,28,28	0
6	EDO	A	2462	4/4	0.96	0.16	1.75	39,40,42,47	0
6	EDO	A	2445	4/4	0.89	0.21	1.56	36,38,39,41	0
6	EDO	A	2444	4/4	0.94	0.17	1.15	29,31,31,32	0
6	EDO	A	2443	4/4	0.95	0.13	0.91	35,35,36,37	0
6	EDO	A	2435	4/4	0.87	0.14	0.63	33,34,34,35	0
6	EDO	A	2438	4/4	0.97	0.10	0.51	25,25,26,27	0
6	EDO	A	2446	4/4	0.95	0.15	0.48	27,27,28,28	0
6	EDO	A	2450	4/4	0.97	0.10	0.25	18,18,18,18	0
3	MN	A	2428	1/1	0.98	0.10	-1.52	16,16,16,16	0
2	CA	A	2427	1/1	0.99	0.04	-1.99	19,19,19,19	0
2	CA	A	2429	1/1	0.99	0.03	-2.67	13,13,13,13	0
2	CA	A	2430	1/1	0.97	0.07	-4.12	21,21,21,21	0
6	EDO	A	2461	4/4	0.81	0.27	-	48,48,49,49	0
6	EDO	A	2463	4/4	0.93	0.23	-	36,39,41,45	0
6	EDO	A	2460	4/4	0.85	0.19	-	36,38,40,44	0
6	EDO	A	2464	4/4	0.79	0.29	-	39,44,49,51	0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
6	EDO	A	2441	4/4	0.88	0.37	-	48,49,50,50	0
6	EDO	A	2453	4/4	0.86	0.24	-	38,43,45,50	0
6	EDO	A	2439	4/4	0.91	0.18	-	41,43,44,47	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.