



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 14, 2017 – 01:46 am GMT

PDB ID : 3A7Q
Title : Structural basis for specific recognition of reelin by its receptors
Authors : Yasui, N.; Nogi, T.; Takagi, J.
Deposited on : 2009-10-01
Resolution : 2.60 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix)	:	1.9-1692
EDS	:	trunk28620
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	2542 (2.60-2.60)
Clashscore	112137	2895 (2.60-2.60)
Ramachandran outliers	110173	2848 (2.60-2.60)
Sidechain outliers	110143	2848 (2.60-2.60)
RSRZ outliers	101464	2550 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	725	<div> <div>3%</div> <div> <div></div> <div>77%</div> <div>17%</div> <div>• 6%</div> </div> </div>
2	B	44	<div> <div>9%</div> <div> <div></div> <div>64%</div> <div>14%</div> <div>• 20%</div> </div> </div>

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 5803 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Reelin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	685	Total	C	N	O	S	0	2	0
			5419	3441	925	1017	36			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1946	GLY	-	EXPRESSION TAG	UNP Q60841
A	1947	ARG	-	EXPRESSION TAG	UNP Q60841
A	2101	ALA	CYS	ENGINEERED	UNP Q60841
A	2662	SER	-	EXPRESSION TAG	UNP Q60841
A	2663	ARG	-	EXPRESSION TAG	UNP Q60841
A	2664	LEU	-	EXPRESSION TAG	UNP Q60841
A	2665	GLU	-	EXPRESSION TAG	UNP Q60841
A	2666	ASN	-	EXPRESSION TAG	UNP Q60841
A	2667	LEU	-	EXPRESSION TAG	UNP Q60841
A	2668	TYR	-	EXPRESSION TAG	UNP Q60841
A	2669	PHE	-	EXPRESSION TAG	UNP Q60841
A	2670	GLN	-	EXPRESSION TAG	UNP Q60841


- Molecule 2 is a protein called Low-density lipoprotein receptor-related protein 8.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	35	Total	C	N	O	S	0	0	0
			289	166	51	66	6			

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	40	GLY	-	EXPRESSION TAG	UNP Q14114
B	41	SER	-	EXPRESSION TAG	UNP Q14114
B	46	GLU	ASP	SEE REMARK 999	UNP Q14114

- Molecule 3 is SUGAR (N-ACETYL-D-GLUCOSAMINE) (three-letter code: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	A	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 4 is a polymer of unknown type called SUGAR (2-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	2	Total	C	N	O	0	0
			28	16	2	10		

- Molecule 5 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	1	Total	Ca	0	0
			1	1		
5	A	4	Total	Ca	0	0
			4	4		

- Molecule 6 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	1	Total 1	Zn 1	0	0


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	18	Total 18	O 18	0	0
7	B	1	Total 1	O 1	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Reelin



• Molecule 2: Low-density lipoprotein receptor-related protein 8



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	56.98Å 93.84Å 73.46Å 90.00° 107.41° 90.00°	Depositor
Resolution (Å)	46.93 – 2.60 46.92 – 2.60	Depositor EDS
% Data completeness (in resolution range)	96.5 (46.93-2.60) 96.4 (46.92-2.60)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	0.07	Depositor
$\langle I/\sigma(I) \rangle$ ¹	5.46 (at 2.61Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.223 , 0.296 0.223 , 0.295	Depositor DCC
R_{free} test set	1135 reflections (5.45%)	DCC
Wilson B-factor (Å ²)	50.1	Xtriage
Anisotropy	0.150	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 61.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	5803	wwPDB-VP
Average B, all atoms (Å ²)	57.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.19% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, CA, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.49	0/5569	0.60	0/7578
2	B	0.47	0/293	0.64	0/394
All	All	0.49	0/5862	0.61	0/7972

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5419	0	5134	76	0
2	B	289	0	231	5	0
3	A	42	0	39	0	0
4	A	28	0	25	0	0
5	A	4	0	0	0	0
5	B	1	0	0	0	0
6	A	1	0	0	0	0
7	A	18	0	0	1	0
7	B	1	0	0	0	0
All	All	5803	0	5429	81	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 81 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:2221:LEU:HD12	1:A:2290:THR:HB	1.63	0.81
2:B:49:LYS:H	2:B:49:LYS:HD2	1.50	0.76
1:A:1985:TYR:O	1:A:2088:TRP:HD1	1.70	0.74
1:A:2055:LEU:HD11	1:A:2095:PHE:HD1	1.54	0.72
1:A:2416:LEU:HB3	1:A:2417:PRO:HD2	1.73	0.68

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	679/725 (94%)	631 (93%)	44 (6%)	4 (1%)	28	53
2	B	33/44 (75%)	28 (85%)	4 (12%)	1 (3%)	5	8
All	All	712/769 (93%)	659 (93%)	48 (7%)	5 (1%)	25	49

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	2272	ASN
1	A	2320	GLY
1	A	2421	GLU
1	A	2417	PRO
2	B	50	ASP

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	599/634 (94%)	584 (98%)	15 (2%)	53	79
2	B	35/41 (85%)	34 (97%)	1 (3%)	48	75
All	All	634/675 (94%)	618 (98%)	16 (2%)	53	79

5 of 16 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	2305	SER
1	A	2366	VAL
1	A	2522	ASN
1	A	2253	LEU
1	A	2559	CYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

2 carbohydrates are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected

value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	NAG	A	3002	1,4	14,14,15	0.55	0	15,19,21	1.10	2 (13%)
4	NAG	A	3003	4	14,14,15	0.54	0	15,19,21	0.64	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	A	3002	1,4	-	0/6/23/26	0/1/1/1
4	NAG	A	3003	4	-	0/6/23/26	0/1/1/1

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	3002	NAG	O5-C1-C2	-2.63	107.82	111.47
4	A	3002	NAG	C4-C3-C2	2.28	114.36	111.02

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.6 Ligand geometry

Of 9 ligands modelled in this entry, 6 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	NAG	A	3001	1	14,14,15	0.55	0	15,19,21	1.20	1 (6%)
3	NAG	A	3004	1	14,14,15	0.54	0	15,19,21	1.56	1 (6%)
3	NAG	A	3005	1	14,14,15	0.54	0	15,19,21	1.19	1 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	A	3001	1	-	0/6/23/26	0/1/1/1
3	NAG	A	3004	1	-	0/6/23/26	0/1/1/1
3	NAG	A	3005	1	-	0/6/23/26	0/1/1/1

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	3005	NAG	C1-O5-C5	2.52	115.65	112.17
3	A	3001	NAG	C1-O5-C5	3.54	117.05	112.17
3	A	3004	NAG	C1-O5-C5	5.41	119.62	112.17

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	685/725 (94%)	0.33	21 (3%) 49 41	49, 57, 64, 71	0
2	B	35/44 (79%)	0.86	4 (11%) 6 3	54, 57, 60, 60	0
All	All	720/769 (93%)	0.36	25 (3%) 44 36	49, 57, 64, 71	0

The worst 5 of 25 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	2299	GLU	5.6
2	B	81	CYS	5.2
2	B	50	ASP	4.4
1	A	2200	SER	3.6
1	A	1964	PHE	3.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
4	NAG	A	3002	14/15	0.84	0.17	-	44,46,47,47	0
4	NAG	A	3003	14/15	0.58	0.31	-	39,43,44,44	0

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	NAG	A	3004	14/15	0.89	0.20	0.58	63,65,67,67	0
3	NAG	A	3005	14/15	0.81	0.20	0.09	57,59,63,63	0
5	CA	A	4004	1/1	0.95	0.17	-0.24	53,53,53,53	0
5	CA	A	4003	1/1	0.91	0.14	-0.98	55,55,55,55	0
5	CA	A	4001	1/1	0.89	0.14	-1.03	52,52,52,52	0
5	CA	A	4002	1/1	0.90	0.13	-1.11	56,56,56,56	0
5	CA	B	5001	1/1	0.98	0.12	-1.25	55,55,55,55	0
6	ZN	A	4005	1/1	0.80	0.12	-	118,118,118,118	0
3	NAG	A	3001	14/15	0.83	0.18	-	50,52,53,53	0

6.5 Other polymers [i](#)

There are no such residues in this entry.